isolated data³: (1) Most of the events were observed during or after CPB. (2) Many cases were associated with the use of the curare alcuronium. (3) Some cases were observed under usual anesthesia with etomidate, fentanyl, pancuronium, and diazepam. (4) Some cases were clearly associated with protamine infusion. (5) At least in one case the vasodilatation was induced after heparin infusion. (6) The vasoplegia was also observed in noncardiac operations. (7) Our impression was that the syndrome happened periodically. (8) Many patients had diabetes. (9) Many patients were previously using the calcium antagonist diltiazem. To these observations we can add the possible effects of the angiotensin-converting enzyme inhibitors as suggested by Tuman and associates⁴ and mentioned by Yiu, Robin, and Pattison.¹

We also have used methylene blue to treat anaphylaxis. Our cumulative clinical experience includes 6 patients. The causes were iodinate radiographic contrast material (5 patients) and penicillin (1 patient). The traditional treatments (high doses of epinephrine and corticosteroids) failed to reverse the cardiocirculatory collapse, urticaria, and angioedema. Four patients were treated by a bolus infusion of 4% methylene blue, 1.5 mg/kg (120 mg), followed by continuous infusion of another 120 mg diluted in 5% glucose in water. The infusion was administered as a bolus because of the severity of shock. Two patients received only continuous infusion during 30 to 40 minutes. The anaphylactic manifestations were completely reversed in 10 to 15 minutes. In 1 patient we observed nodal cardiac rhythm over 1 minute during the infusion of the dye, which reversed spontaneously. One hypertensive patient had angina pectoris during infusion, without any electrocardiographic manifestation, and did not require coronary vasodilators to reverse the symptoms. We obtained the hemodynamic registers of 2 patients who had anaphylactic shock caused by radiographic contrast material in the cardiovascular catheterization laboratory. Cumulative experience would indicate that this dye is the treatment of choice or an alternative to the treatment of distributive circulatory shock. I am reporting, probably, the first experience treating anaphylaxis in human beings with methylene blue, and it deserves wide circulation because it is a lifesaving choice.⁵

Nitric oxide stimulates soluble guanylate cyclase and activates the production of cyclic guanosine 3',5'-monophosphate (cGMP), resulting in relaxation. Inhibition of nitric oxide synthesis is not approved for clinical use because of the lack of specificity to the inducible form of nitric oxide synthase and its association with tissue necrosis. Methylene blue inhibits guanylate cyclase, avoiding the cGMP-dependent vasorelaxant effect of nitric oxide in the smooth muscle of the vessels. Also, thiazine dye is widely used in clinical medicine to reduce methemoglobin, to treat malaria, and to cleanse the urinary tract. Otherwise, methylene blue is not associated with tissue necrosis, as caused by nitric oxide synthase inhibition, and its lethal dose is more than 40 mg/kg, as shown experimentally in sheep. Myles, Leong, and Curry⁶ failed to identify an increase in nitric oxide release after, or as a consequence of, CPB on the basis of the

serum and urine nitrite/nitrate levels. Since methylene blue blocks the soluble guanylate cyclase, which is the final effect on the nitric oxide pathway, it is hard to determine a different cGMP-dependent mechanism, and the hypothesis of direct activation of this final common pathway of nitric oxide release is very improbable. I guess it is more logical to speculate on the existence of another non-nitric cGMP-dependent pathway, with different receptors and/or different G-proteins, or else a sensitization of vascular smooth muscle cells to the action of nitric oxide, as hypothesized by Yiu, Robin, and Pattison.¹

Paulo Roberto B. Evora, MD, PhD
Division of Experimental Surgery
Ribeirão Preto School of Medicine
University of São Paulo
Department of Surgery
UNAERP Medical School
Heart Hospital of Ribeirão Preto
Rua Rui Barbosa, 367, Ap 7
14015-120 Ribeirão Preto, SP, Brazil

REFERENCES

1. Yiu P, Robin J, Pattison CW. Reversal of refractory hypotension with single-dose methylene blue after coronary artery bypass surgery. *J Thorac Cardiovasc Surg* 1999;118:195-4.
2. Andrade JCS, Batista Filho ML, Evora PRB, et al. Methylene blue administration in the treatment of the vasoplegic syndrome after cardiac surgery. *Rev Bras Circ Cardiovasc* 1996;11:107-14.
3. Evora PRB, Ribeiro PJF, Andrade JCS. Methylene blue administration in SIRS after cardiac operations [letter/comment]. *Ann Thorac Surg* 1997;63:12-3.
4. Tuman KJ, McCarthy KJ, O'Connor CJ, Holm WE, Ivankovich AD. Angiotensin-converting enzyme inhibitors increase vasoconstrictor requirements after cardiopulmonary bypass. *Anesth Analg* 1995;80:473-9.
5. Evora PRB, Roselino CH, Schiavetto PM. Methylene blue in anaphylactic shock [letter]. *Ann Emerg Med* 1997;30:240.
6. Myles PS, Leong CK, Currey J. Endogenous nitric oxide and low systemic vascular resistance after cardiopulmonary bypass. *J Cardiothorac Vasc Anesth* 1997;11:571-4.

12/8/104472

doi:10.1067/mtc.2000.104472

Reply to the Editor:

We thank Evora for adding his experience with methylene blue. In most instances of low systemic vascular resistance (SVR) syndrome necessitating pharmacologic intervention, vasopressor therapy over a number of hours is limited. Thus whether methylene blue should be the drug of choice is speculative at this stage. Nevertheless, in cases in which first-line therapy has failed, the dramatic success and long-lasting effect of this drug that we found in our case, and that Evora and his colleagues presumably found as well, are certainly impressive. This leads us to propose that early use of methylene blue may halt the progression of low SVR even in patients responsive to norepinephrine and mitigate the need for a protracted course of vasoconstrictor use. A ran-

domized clinical study to evaluate this hypothesis is justifiable.

Patrick Yiu, FRCS
Cardiothoracic Surgical Unit
Middlesex Hospital
4th Floor
Mortimer St
London W1N 8AA, United Kingdom
12/8/104473
doi:10.1067/mtc.2000.104473

Combined mitral valve repair and the Cox maze procedure for mitral valve prolapse and regurgitation and associated atrial fibrillation

To the Editor:

I read with great interest the report by Handa and associates¹ on the early and intermediate term results of combined mitral valve repair and the Cox maze procedure for patients with severe mitral regurgitation and associated atrial fibrillation. It is gratifying to note that at the Mayo Clinic the addition of the Cox maze procedure to mitral valve repair not only is safe and effective but also decreases late complications.

One of the serious complications of mitral valve prolapse, which is otherwise a common² and benign³ condition, is mitral regurgitation.^{4,5} In fact, the commonest cause of mitral regurgitation in the United States is mitral valve prolapse.^{5,6} Mitral regurgitation in mitral valve prolapse may become progressive in some patients.⁵ One of the serious complications of progressive mitral regurgitation is atrial fibrillation, which usually persists even after successful corrective surgery of the mitral valve and often recurs after pharmacologic or electric cardioversion. Although postoperative atrial fibrillation can be successfully managed by antiarrhythmic drugs and long-term anticoagulant therapy to prevent thromboembolism, these therapeutic modalities are not without side-effects, torsades de pointes and bleeding, respectively. It is gratifying that the adjunctive Cox maze procedure at the Mayo Clinic increased the restoration of sinus rhythm to 82% of their patients.

Therefore, as the authors concluded, "Although the addition of the Cox maze procedure lengthens hospital stay in patients having mitral valve repair, overall medical costs may be decreased by reducing the need for long-term anticoagulation, as well as costs associated with stroke and bleeding complications."¹

Tsung O. Cheng, MD
Professor of Medicine
The George Washington University
Washington, DC 20037

REFERENCES

1. Handa N, Schaff HV, Morris JJ, Anderson BJ, Kopecky SL, Enriquez-Sarano M. Outcome of valve repair and the Cox maze procedure for mitral regurgitation and associated atrial fibrillation. *J Thorac Cardiovasc Surg* 1999;118:628-35.
2. Cheng TO, Barlow JB. Mitral valve leaflet billowing and prolapse: its prevalence around the world. *Angiology* 1989;40:77-87.
3. Barlow JB, Cheng TO. Mitral valve billowing and prolapse. In: Cheng TO. *The International Textbook of Cardiology*, New York/Oxford: Pergamon Press; 1987. p. 497-524.
4. Cheng TO. Mitral valve prolapse. *Dis Month* 1987;33:481-534.
5. Cheng TO. Mitral valve prolapse: an overview. *J Cardiol* 1989;19 (Suppl 21):3-20.
6. Cheng TO. Mitral-valve prolapse. *N Engl J Med* 1999;341:1471-2.

12/8/104693

doi:10.1067/mtc.2000.104693

Should the bidirectional Glenn procedure be better performed through the support of cardiopulmonary bypass?

To the Editor:

We read with great interest the report of Jahangiri and associates¹ and the commentary by Jonas on the technique of bidirectional Glenn shunt through a thoracotomy without the use of cardiopulmonary bypass (CPB). We agree that clamping the superior vena cava (SVC) without decompressing the internal jugular system exposes the brain to the effects of reduced cerebral perfusion pressure. We have used electrophysiological indices of cerebral function (electroencephalography or evoked potentials) and transcranial Doppler during bidirectional Glenn shunts in patients with pulmonary atresia without support (n = 2) or through the use of CPB (n = 4). During clamping of the SVC without CPB, major reductions (>50%) in the diastolic, mean, and peak systolic blood flow velocities of the middle cerebral artery were identified, which were followed by mild electrocortical alterations as indicated by longer latencies of the cortically generated evoked potentials.² In contrast, this situation did not occur or was minimal in those cases in which clamping of the SVC was done with the support of CPB. We have learned that Doppler flow changes and electrocortical alterations may be expected as a result of acute SVC hypertension if the internal jugular venous system is not decompressed.³ In one of our cases without CPB, an intraoperative shunt into the SVC was indicated due to the presence of extremely low flow velocities and electrocortical alterations during SVC clamping. In this case, while the SVC shunt was patent, flow velocities were maintained and the latencies of evoked potentials returned to preclamping levels.⁴ In our institution, we routinely use CPB and intraoperative brain function monitoring by transcranial Doppler, near-infrared spectrophotometry, and electroencephalography for children undergoing these procedures.³ A note of caution should be made that the reported absence of gross neurologic deficits in Jahangiri and colleagues' cases is not an indicator that the brain is free of potential alterations during SVC clamping without CPB. Furthermore, under conditions of normothermic ischemia, the brain does not receive the protective benefit of hypothermia.⁵ Without brain protection, even short intervals of low cerebral perfusion could generate minor or subclinical deficits that might be detectable only through detailed cognitive testing. We agree with Jonas' comments that the demonstration of safety of this procedure will rest on the careful