ent comparison/substitute: the RA for the vein graft. Dr Lytle's comment in his editorial, "in my judgment, the RA graft is less predictable than the RITA graft in regard to patency," intrigued me. Over the past 15 years as a practicing cardiothoracic surgeon, I have become profoundly aware of the inadequacy of veins, and since my recent attendance at the enlightening and energy-invoking symposium "Arterial Conduits for Myocardial Revascularization" in Rome by Dr G. F. Possatti and Dr A. M. Calafiore, I believe the thrust should be to continue to use double ITA grafts whenever possible (especially in the young) but to substitute the RA for the vein graft. Since my return from this symposium, I have tried to do this; perhaps I did not see (or did not want to see) before, but many patients have serious venous disease of their legs precluding use of the saphenous vein. I think one of the turning points for me was when I recently (July 24, 2003) had to reoperate on an 83-year-old woman, on whom I had placed 2 grafts at age 81. Her LITA had gone down, I believe because I placed the graft inadvertently above a stenosis and her vein graft to a marginal artery occluded. If her vein graft had stayed open, she probably would not have needed reoperation at age 83. I used a sequential RA graft to the left anterior descending coronary artery and the marginal branch on-pump, and postoperatively she woke up stating that this was easier than her first operation. (Obviously this time both grafts were working!)

A second reason to substitute the vein for the RA graft could be the anticipated longer-lasting results of the drug-eluting stents used by interventional cardiologists. I know we are all interested in the same end—stamping out coronary artery disease effects—but it can be a little disconcerting for many surgeons currently in practice to see their favored coronary artery bypass grafts (LITA plus 2 veins) going the way of the dodo bird. I want to be so bold as to predict that the drug-eluting stents will rival our saphenous vein grafts (that is, by the time they figure out which drug, from which drug family, how much eluting, over what period of time, and so on, works; it might take 20 years). Although we should never be competitive with our interventional

colleagues because we have the same end point in common, we must as surgeons find something ancillary to their work. I believe total arterial grafting (be it bilateral ITA, LITA/RA, RITA, bilateral ITA/gastroepiploic artery) may well be the answer. As Dr Lytle most wisely stated at the 2003 meeting of The American Association for Thoracic Surgery in Boston (and I quote him often in this): "Did you think you were going to be doing the same operation for 100 years!?" Dr Lytle continues to be a driving force leading all our quests for the best coronary artery bypass conduit.

Teresa Kieser, MD, FRCSC, FACS
Department of Cardiac Sciences
Calgary Health Region Foothills Medical
Centre
Calgary, Alberta, Canada

Reference

1. Lytle BW. Radial versus right internal thoracic artery as a second arterial conduit for coronary surgery: early and midterm outcomes. *J Thorac Cardiovasc Surg.* 2003;126:5-6.

doi:10.1016/j.jtcvs.2003.09.015

Angiotensin-converting enzyme inhibitors predispose to hypotension refractory to norepinephrine but responsive to methylene blue

To the Editor:

We read with interest the brief communication of Grayling and Deakin¹ describing the methylene blue administration in a patient with refractory hypotension during cardiac surgery. We're currently using this drug for patients with high cardiac output and hypotension unresponsive to norepinephrine at doses similar to those suggested by Grayling and Deakin (up to 3 mg/kg in 2 different administration instead of 2 mg/kg in a single dose). We would like to suggest a possible predisposing factor of vasoplegia that responds to methylene blue: in our experience² and in the literature³⁻⁵ it is evident that most episodes of refractory hypotension are related to preoperative intake of angiotensin-converting enzyme inhibitors (ACEI); even the patient described in this report was assuming ACEI (ramipril). After experiencing a dramatic refractory hypotension in a patient on a program of ACEI,² we started using

methylene blue in case of refractory hypotension and withdrawing ACEI the day before surgery. We observed an immediate and sustained response even at doses less than 1 mg/kg.

We would like to reinforce this kind of therapeutic approach and suggest a common predisposing factor that has been so far underestimated: preoperative intake of ACEI.

Donatella Sparicio, MD

Giovanni Landoni, MD

Alberto Zangrillo, MD

Vita-Salute University of Milano, IRCCS San

Raffaele Hospital

Milano, Italy

References

1. Grayling M, Deakin CD. Methylene blue during cardiopulmonary bypass to treat refractory hypotension in septic endocarditis. *J Thorac Cardiovasc Surg.* 2003;125:426-7.
2. Pappalardo F, Landoni G, Franco A, Monaco C, Marino G, Torri G. Prolonged refractory hypotension in cardiac surgery after institution of cardiopulmonary bypass. *J Cardiothorac Vasc Anesth.* 2002;16:477-9.
3. Yiu P, Robin J, Pattison CW. Reversal of refractory hypotension with single dose methylene blue after coronary artery bypass surgery. *J Thorac Cardiovasc Surg.* 1999;118:194-5.
4. Pagni S, Austin EH. Use of intravenous methylene blue for the treatment of refractory hypotension after cardiopulmonary bypass. *J Thorac Cardiovasc Surg.* 2000;119:1297-8.
5. Berkowitz DE, Richardson C, Elliott D, Leslie JB, Schwinn DA. Hypotension resistant to therapy with alpha receptor agonists complicating cardiopulmonary bypass: lithium as a potential cause. *Anesth Analg.* 1996;82:1082-5.

doi:10.1016/j.jtcvs.2003.01.001

Modified Fontan procedure in adults

To the Editor:

At the outset I must congratulate and thank the authors of this article for this landmark work on the Fontan procedure in an adult population.¹ This is going to be a benchmark for adult Fontan procedure in years to come.

In the Western hemisphere the Fontan surgery is done in children; adult patients undergoing this procedure are exceptions, not the rule. But in developing countries adult patients for univentricular palliation, with or without prior surgical palliation, are quite common.² As the authors have addressed, the primary issue for these pa-