ent comparison/substitute: the RA for the vein graft. Dr Lytle's comment in his editorial, "in my judgment, the RA graft is less predictable than the RITA graft in regard to patency," intrigued me. Over the past 15 years as a practicing cardiothoracic surgeon, I have become profoundly aware of the inadequacy of veins, and since my recent attendance at the enlightening and energy-invoking symposium "Arterial Conduits for Myocardial Revascularization" in Rome by Dr G. F. Possatti and Dr A. M. Calafiore, I believe the thrust should be to continue to use double ITA grafts whenever possible (especially in the young) but to substitute the RA for the vein graft. Since my return from this symposium, I have tried to do this; perhaps I did not see (or did not want to see) before, but many patients have serious venous disease of their legs precluding use of the saphenous vein. I think one of the turning points for me was when I recently (July 24, 2003) had to reoperate on an 83-year-old woman, on whom I had placed 2 grafts at age 81. Her LITA had gone down, I believe because I placed the graft inadvertently above a stenosis and her vein graft to a marginal artery occluded. If her vein graft had stayed open, she probably would not have needed reoperation at age 83. I used a sequential RA graft to the left anterior descending coronary artery and the marginal branch on-pump, and postoperatively she woke up stating that this was easier than her first operation. (Obviously this time both grafts were working!)

A second reason to substitute the vein for the RA graft could be the anticipated longer-lasting results of the drug-eluting stents used by interventional cardiologists. I know we are all interested in the same end—stamping out coronary artery disease effects—but it can be a little disconcerting for many surgeons currently in practice to see their favored coronary artery bypass grafts (LITA plus 2 veins) going the way of the dodo bird. I want to be so bold as to predict that the drug-eluting stents will rival our saphenous vein grafts (that is, by the time they figure out which drug, from which drug family, how much eluting, over what period of time, and so on, works; it might take 20 years). Although we should never be competitive with our interventional

colleagues because we have the same end point in common, we must as surgeons find something ancillary to their work. I believe total arterial grafting (be it bilateral ITA, LITA/RA, RITA, bilateral ITA/gastroepiploic artery) may well be the answer. As Dr Lytle most wisely stated at the 2003 meeting of The American Association for Thoracic Surgery in Boston (and I quote him often in this): "Did you think you were going to be doing the same operation for 100 years!?" Dr Lytle continues to be a driving force leading all our quests for the best coronary artery bypass conduit.

Teresa Kieser, MD, FRCSC, FACS
Department of Cardiac Sciences
Calgary Health Region Foothills Medical
Centre
Calgary, Alberta, Canada

Reference

1. Lytle BW. Radial versus right internal thoracic artery as a second arterial conduit for coronary surgery: early and midterm outcomes. *J Thorac Cardiovasc Surg.* 2003;126:5-6.

doi:10.1016/j.jtcvs.2003.09.015

Angiotensin-converting enzyme inhibitors predispose to hypotension refractory to norepinephrine but responsive to methylene blue To the Editor:

We read with interest the brief communication of Grayling and Deakin¹ describing the methylene blue administration in a patient with refractory hypotension during cardiac surgery. We're currently using this drug for patients with high cardiac output and hypotension unresponsive to norepinephrine at doses similar to those suggested by Grayling and Deakin (up to 3 mg/kg in 2 different administration instead of 2 mg/kg in a single dose). We would like to suggest a possible predisposing factor of vasoplegia that responds to methylene blue: in our experience² and in the literature³⁻⁵ it is evident that most episodes of refractory hypotension are related to preoperative intake of angiotensin-converting enzyme inhibitors (ACEI); even the patient described in this report was assuming ACEI (ramipril). After experiencing a dramatic refractory hypotension in a patient on a program of ACEI,² we started using

methylene blue in case of refractory hypotension and withdrawing ACEI the day before surgery. We observed an immediate and sustained response even at doses less than 1 mg/kg.

We would like to reinforce this kind of therapeutic approach and suggest a common predisposing factor that has been so far underestimated: preoperative intake of ACEI.

Donatella Sparicio, MD

Giovanni Landoni, MD

Alberto Zangrillo, MD

Vita-Salute University of Milano, IRCCS San Raffaele Hospital
Milano, Italy

References

1. Grayling M, Deakin CD. Methylene blue during cardiopulmonary bypass to treat refractory hypotension in septic endocarditis. *J Thorac Cardiovasc Surg.* 2003;125:426-7.
2. Pappalardo F, Landoni G, Franco A, Monaco C, Marino G, Torri G. Prolonged refractory hypotension in cardiac surgery after institution of cardiopulmonary bypass. *J Cardiothorac Vasc Anesth.* 2002;16:477-9.
3. Yiu P, Robin J, Pattison CW. Reversal of refractory hypotension with single dose methylene blue after coronary artery bypass surgery. *J Thorac Cardiovasc Surg.* 1999;118:194-5.
4. Pagni S, Austin EH. Use of intravenous methylene blue for the treatment of refractory hypotension after cardiopulmonary bypass. *J Thorac Cardiovasc Surg.* 2000;119:1297-8.
5. Berkowitz DE, Richardson C, Elliott D, Leslie JB, Schwinn DA. Hypotension resistant to therapy with alpha receptor agonists complicating cardiopulmonary bypass: lithium as a potential cause. *Anesth Analg.* 1996;82:1082-5.

doi:10.1016/j.jtcvs.2003.01.001

Modified Fontan procedure in adults To the Editor:

At the outset I must congratulate and thank the authors of this article for this landmark work on the Fontan procedure in an adult population.¹ This is going to be a benchmark for adult Fontan procedure in years to come.

In the Western hemisphere the Fontan surgery is done in children; adult patients undergoing this procedure are exceptions, not the rule. But in developing countries adult patients for univentricular palliation, with or without prior surgical palliation, are quite common.² As the authors have addressed, the primary issue for these pa-

tients is suitability for surgery. I am unclear on a few points in the article, however, and have some questions for the authors.

This landmark work began just after introduction of the Fontan principle. Is this why initial modifications were done in a majority of patients? Or is there another reason? Do you continue to do these modifications today? If so, for what indications and why? The issue of superiority of 1 modification over others in long-term results is far from resolved. For budding cardiac surgeons like me, which type of modification do you recommend for hemodynamically adequate adult patients? In other words, what should govern my choice of modification in different sets of patients?

The fenestrated Fontan modification has widened the horizon of univentricular repair. You have elaborately described your point. Do you feel in spite of anticoagulation the incidence of thromboembolic and other side effects or complications outweighs the benefit of fenestration in decreasing the morbidity of procedure, which has been vigorously tested? Does the age of the patients also affect the decision to fenestrate? If yes, in what way?^{2,3}

What is your policy regarding 1-stage or 2-stage procedure? For example, do you perform superior cavopulmonary first and then complete the Fontan as a separate procedure, especially for patients without palliation or with systemic to pulmonary artery shunts or pulmonary artery banding? Is the principle of gradual volume unloading of the ventricle different in adults than children?

Once again, I thank the authors for this outstanding article.

Manoj Purohit
SPR, Paediatric Cardiac Surgery
Alder Hey Hospital
Liverpool, United Kingdom

References

1. Burkhart HM, Dearani JA. The modified Fontan procedure: early and late results in 132 adult patients. *J Thorac Cardiovasc Surg.* 2003;125:1252-9.
2. Airan B, Sharma R, Choudhary S, Mohanty SR. Univentricular repair: is routine fenestration justified? *Ann Thorac Surg.* 2000;69:1900-6.
3. Lemler MS, Scott WA, Steven R. Fenestration improve clinical outcome of Fontan procedure: a prospective randomized study. *Circulation.* 105:207-12.
doi:10.1016/j.jtcvs.2003.09.025

Reply to the Editor:

In the setting of single ventricle physiology, the common and accepted practice is to proceed with the Fontan procedure at 2 to 4 years of age in the majority of circumstances. In developing countries, the opportunity for staged palliation with eventual Fontan procedure in young children is less common. Some children may have undergone a palliative operation with a systemic-to-pulmonary artery shunt or cavopulmonary shunt during early childhood but were lost to follow-up or never had the opportunity for the completion Fontan procedure. Consequently, there are potentially a greater number of older patients who may be candidates for the Fontan procedure in developing countries.

In our experience with more than 1000 patients, surgical modifications to the Fontan procedure have evolved over the last 3 decades in an effort to avoid complications occurring late after operation. Currently, the technique of the Fontan procedure that is applied in our patients includes the total cavopulmonary connection with an extracardiac conduit or intra-atrial conduit. We believe this technique will minimize the incidence of late atrial arrhythmias.

To reduce the risk of thromboembolism, we eliminate blind stumps (eg, segment above the pulmonary valve) and blind intracardiac chambers (eg, vestigial right ventricle in tricuspid atresia via the ventricular septal defect) that are exposed to the systemic circulation. This is accomplished by patch or suture closure of the pulmonary valve annulus or closure of the ventricular septal defect.

When significant risk factors for early mortality are present prior to the Fontan procedure, such as severe atrioventricular valve regurgitation, severe pulmonary artery abnormalities, or severe subaortic stenosis, we would proceed with a staged surgical approach that corrects these abnormalities prior to the completion Fontan procedure. This would include repair or replacement of the atrioventricular valve, relief of pulmonary artery stenosis or distortion, and relief of subaortic stenosis (resection or Damus-Stansel-Kaye anastomosis). A concomitant bidirectional cavopulmonary anastomosis is also commonly performed at this time. The completion of the Fontan procedure would then be performed at a subsequent operation. Minor intracardiac or pulmonary artery abnor-

malities are corrected at the time of the Fontan procedure. In the absence of risk factors, we do not believe it is necessary to perform a bidirectional cavopulmonary shunt prior to the Fontan procedure.

We have used the fenestration selectively at the time of the Fontan procedure. Although we attempt to correct major abnormalities during the course of a staged surgical approach in an effort to reduce operative risk, some reversible risk factors for early or late mortality may still be present. These may include elevated preoperative pulmonary artery pressure, reversible ventricular dysfunction, increased ventricular mass, or uncorrected pulmonary artery abnormalities. Fenestration has also been used in patients with predominant right ventricular morphology.

The role of warfarin anticoagulation after the Fontan procedure has been controversial. In general, we tend to anticoagulate most adult patients, particularly those with a fenestration or residual intracardiac shunt, history of arrhythmias, intracardiac thrombus, slow circulation, and venous varices. In children and young adult patients, we may substitute aspirin as an alternative to warfarin anticoagulation.

Joseph A. Dearani, MD
Francisco J. Puga, MD
Gordon K. Danielson, MD
Division of Cardiovascular Surgery,
Mayo Clinic, Rochester, MN 55905
doi:10.1016/j.jtcvs.2003.09.026

Mitral valve surgery with extensive calcification of the annulus

To the Editor:

Feindel and coauthors¹ report results in a challenging set of patients, those with mitral annular calcification (MAC) causing mitral regurgitation. Several alternative techniques to avoid excising the calcified bar all have disadvantages. Displacing the prosthetic sewing line into the left atrium (with or without a prosthetic cuff) subjects the left atrium to high ventricular pressure. Sewing to the mitral leaflets central to the calcified bar requires sewing to fragile tissue and downsizing the valve. Sewing to the calcified bar usually leads to dehiscence, and trying to pass the sutures around the bar jeopardizes the circumflex coronary artery.