References

1. Suzuki K, Murtaza B, Heslop L, Morgan JE, Smolenski RT, Suzuki N, et al. Single fibers of skeletal muscle as a novel graft for cell transplantation to the heart. *J Thorac Cardiovasc Surg.* 2002;123:984-92.
2. Menasché P, Hagege AA, Scorsin M, Pouzet B, Desnos M, Duboc D, et al. Myoblast transplantation for heart failure. *Lancet.* 2001;357:279-80.
3. Pouzet B, Vilquin JT, Hagege A, Scorsin M, Messas E, Fiszman M, et al. Factors effecting functional outcome of autologous skeletal myoblast transplantation. *Ann Thorac Surg.* 2001;71:844-51.
4. Neumeyer AM, DiGregorio DM, Brown RH Jr. Arterial delivery of myoblasts to skeletal muscle. *Neurology.* 1992;42:2258-62.
5. Suzuki K, Murtaza B, Suzuki N, Smolenski RT, Yacoub MH. Intracoronary in fusion of skeletal myoblasts improves cardiac function in doxorubicin-induced heart failure. *Circulation.* 2001;104 Suppl 1:I213-7. doi:10.1067/mtc.2003.288

Reply to the Editor:

We appreciate the interest of Sim and colleagues in our article describing the implantation of single, isolated skeletal muscle fibers into the heart. We agree that many important questions need to be answered before further clinical trials of myoblast transplantation into the heart.

A major issue is the mechanism by which skeletal myoblast transplantation can improve cardiac function. Sim and colleagues' suggestion that the improvement in myocardial function is directly related to the number of myoblasts implanted has not been firmly established. We consider the quality of cells implanted to be as important as, or perhaps even more important than, cell number.¹ In addition, only one recent study with donor cardiac myocytes has attempted to adequately quantify graft cell survival with quantitative real-time polymerase chain reaction, and no direct correlation between numbers of surviving donor-derived cells and functional improvement was made in that study.²

It should be noted that the issues regarding the adverse effects of prolonged in vitro culture on skeletal myoblasts are far from theoretical, as suggested by Sim and colleagues. There is clear evidence for the susceptibility to infection, as well as altered immunogenicity and adverse effects on myogenicity.³ With respect to

the risk of coronary embolism after cell transplantation, we have reported that infusion of more than 2 million myoblasts in a volume of 1 mL into the coronary circulation of the rat heart results in embolization, with edema and cardiac arrest.⁴

Sim and colleagues suggest that the functional benefit of myoblast transplantation is due, at least in an ischemic heart failure model, to the formation of new muscle ("neomyogenesis") with "phenotypic features similar to skeletal and cardiac muscle." This statement is not clear, and such a mechanism would require extensive functional, electrophysiological, and molecular characterization.

The relevance of the discordant human-to-pig xenogenic model cited by Sim and colleagues is not clear, and it is unlikely that a significant degree of myoblast survival would be seen after grafting with only transient immunosuppression—although the protocol for this is not detailed. We find interpretation of the staining in the figure difficult. With respect to labeling, the promoter driving reporter *lacZ* expression is not stated; thus one cannot be absolutely certain as to the identity of the stained cells in the presented figure. The varied distribution of the X-gal stain is difficult to explain, and counterstaining with skeletal myosin-specific antibody, as reported in our own data, would be informative; an additional nuclear marker would confirm the specificity of the stain.

The key findings in our study are that myogenic precursor cells derived from the parent fibers appeared to survive, proliferate, and differentiate within the cardiac microenvironment. Further, this treatment improved cardiac function in ischemic and dilated cardiomyopathy models to a degree that was far beyond what one might expect for the amount of new muscle formed. We have proposed at least two possible causes for this remarkable effect. One is a paracrine effect resulting in neovascularization, as shown in our article; the other is the endogenous protection afforded to the resident satellite cells by the parent fiber. This latter effect may be important in enhancing graft cell survival in the most hazardous acute phase after grafting and may also deliver satellite cells that have some important qualitative differences. We consider the skeletal muscle fiber system to be

promising, although further studies are needed to investigate the effects that we have proposed.

Bari Murtaza

Ken Suzuki

Magdi H. Yacoub

Harefield Hospital Heart Science Centre

Imperial College Faculty of Medicine

London, United Kingdom

References

1. Qu-Petersen Z, Deasy B, Jankowski R, Ikezawa M, Cummins J, Pruchnic R, et al. Identification of a novel population of muscle stem cells in mice: potential for muscle regeneration. *J Cell Biol.* 2002;157:851-64.
2. Muller-Ehmsen J, Whittaker P, Kloner RA, Dow JS, Sakoda T, Long TI, et al. Survival and development of neonatal rat cardiomyocytes transplanted into adult myocardium. *J Mol Cell Cardiol.* 2002;34:107-16.
3. Rosenblatt JD, Lunt AI, Parry DJ, Partridge TA. Culturing satellite cells from living single muscle fiber explants. *In Vitro Cell Dev Biol Anim.* 1995;31:773-9.
4. Suzuki K, Brand NJ, Smolenski RT, Jayakumar J, Murtaza B, Yacoub MH. Development of a novel method for cell transplantation through the coronary artery. *Circulation.* 2000;102(Suppl):III-359-64. doi:10.1067/mtc.2003.289

Large cell neuroendocrine carcinoma of the lung: What we have to do

To the Editor:

I read with interest the excellent article of Takei and colleagues¹ about large cell neuroendocrine carcinoma (LCNC) of the lung. This series represents the largest ever published in English language literature.

The family of neuroendocrine tumors of the lung has recently been the object of study by several pathologists and surgeons. The recent World Health Organization classification criteria² consider neuroendocrine tumors of the lung as a distinct subset of neoplasm with specific morphologic, ultrastructural, immunohistochemical, and molecular characteristics. Travis and associates³ distinguish between typical and atypical carcinoid, LCNC, and small cell neuroendocrine carcinoma of the lung. Of these tumors, LCNCs appear to have an intermediate biologic behavior between atypical carcinoid and small cell carcinoma.³ LCNCs have a tendency to spread

early, yielding mediastinal lymph node metastases or distant metastases (generally in the liver or in the brain) at the time of diagnosis.

Correct preoperative diagnosis of neuroendocrine carcinoma of the lung (and also of LCNC) is really difficult to obtain either by bronchoscopy or by transthoracic biopsy; in the case of clinical N2 neuroendocrine carcinoma, a surgical approach can be considered, because these tumors do not respond to neoadjuvant therapy.⁴ This could be appropriate in case of bronchial carcinoid, but not in case of LCNC. For these tumors, I believe, neoadjuvant treatment is mandatory because of their aggressive biologic behavior.

In this study Takei and colleagues¹ pointed out the importance of the immunohistochemical assessment, with the detection of chromogranin A staining in neoplastic cells. We must remember that LCNCs, because they are neuroendocrine carcinomas, express somatostatin receptors, which can be easily detected in vitro by reverse transcriptase polymerase chain reaction.⁵ In vivo expression of these receptors is provided by indium 111-tagged diethylenetriaminepentaacetic acid pentetreotide (OctreoScan, OCT) scintigraphy.^{6,7} This procedure is widely accepted both in the preoperative evaluation of neuroendocrine carcinoma, because of its effectiveness in detecting mediastinal or distant metastases with high sensitivity, and in follow-up for early detection of tumor recurrences or distant metastases.⁸ In case of positive OCT scintigraphy result, therapy with octreotide (new generation somatostatin analog) might be considered, alone or in association with chemotherapy. I believe that such an important biologic concept might be considered in the correct multidisciplinary treatment of LCNC of the lung. Octreotide has been proved to be effective in controlling metastatic growth, well tolerated by the patient, and without important side effects.^{8,9}

In conclusion, I congratulate Takei and colleagues¹ for this excellent article on such a difficult matter. I recommend the use of OCT scintigraphy in the preoperative evaluation of these patients and in their follow-up. Furthermore the use of octreotide (in the case of positive results of OCT scintigraphy), in association with traditional adjuvant therapy, might be consid-

ered to reduce the risk of tumor recurrence or distant metastases.

Pier Luigi Filosso, MD
University of Torino Italy
San Giovanni Battista Hospital
Department of Thoracic Surgery
Via Genova, 3
10126 Torino, Italy

References

1. Takei H, Asamura H, Maeshima A, Suzuki K, Kondo H, Niki T, et al. Large cell neuroendocrine carcinoma of the lung: a clinicopathologic study of eighty-seven cases. *J Thorac Cardiovasc Surg.* 2002;124:285-92.
2. World Health Organization. Histological typing of lung and pleural tumours. 3rd ed. Geneva: The Organization; 1999.
3. Travis WD, Colby TV, Corrin B, Shimosato Y, Brambilla E. Histological typing of lung and pleural tumours. 3rd ed. Berlin: Springer Verlag; 1999.
4. Martini N, Zaman MB, Bains MS, Burt ME, McCormack PM, Rusch VW, et al. Treatment and prognosis in bronchial carcinoids involving regional lymph nodes. *J Thorac Cardiovasc Surg.* 1994;107:1-6.
5. Papotti M, Croce S, Bello M, Bongiovanni M, Allia E, Schindler M, et al. Expression of somatostatin receptor types 2, 3 and 5 in biopsies and surgical specimens of human lung tumours. *Virchows Arch.* 2001;439:787-97.
6. Kwekkeboom DJ, Kho GS, Lamberts SW, Reubi JC, Laissus JA, Krenning EP. The value of octreotide scintigraphy in patients with lung cancer. *Eur J Nucl Med.* 1994;21:1106-13.
7. Musi M, Carbone RG, Bertocchi C, Cantalupi DP, Michetti G, Pugliese C, et al. Bronchial carcinoid tumours: a study on clinicopathological features and role of octreotide scintigraphy. *Lung Cancer.* 1998;22:97-102.
8. Filosso PL, Ruffini E, Oliaro A, Papalia E, Donati G, Rena O. Long-term survival of atypical bronchial carcinoid with liver metastases, treated with octreotide. *Eur J Cardiothorac Surg.* 2002;21:913-7.
9. O'Byrne KJ, Schally AV, Thomas A, Carney DN, Steward WP. Somatostatin, its receptors and analogs, in lung cancer. *Chemotherapy.* 2001;47:78-108.
doi:10.1067/mtc.2003.321

Reply to the Editor:

We appreciate the thoughtful comments by Filosso on our recent article¹ describing the clinicopathologic aspects of large cell neuroendocrine carcinoma (LCNC) of the lung.

The histologic type of LCNC of the lung has been included as the subtype of large cell carcinoma in the revised World Health Organization classification of lung and pleural tumors in 1999.² The clinico-

pathologic features, however, have not been described in detail. Especially little is known about the response of LCNC to the chemotherapy and radiotherapy. Considering the analogy of LCNC to the small cell histologic type, a multimodal therapeutic approach might be expected to be promising.

As Filosso pointed out, we also have a great interest in the expression of somatostatin receptors in LCNC. If these receptors are expressed frequently in LCNC, octreotide scintigraphy with somatostatin analog (OctreoScan) would reasonably be of great utility in the preoperative evaluation of extent of disease of LCNC. Indeed, carcinoid tumor and small cell carcinoma have already been shown to be well analyzed by octreotide scintigraphy.^{3,4} As for LCNC, however, according to a report by Jiang and colleagues,⁵ only 9 (40.9%) of 22 cases were shown to stain positively for somatostatin, and these tumor cells do not necessarily express somatostatin receptors. Whether octreotide scintigraphy can be used depends on the rate of expression of somatostatin receptors. At present no data are available on the expression of somatostatin receptors in LCNC.

Furthermore, the preoperative establishment of a histologic diagnosis of LCNC is crucial in planning the LCNC-specific treatment strategy. However, it is actually quite difficult to get the histologic or cytologic diagnosis of LCNC simply by the evaluation of tiny preoperative biopsy specimen. Novel LCNC-specific markers ("immunohistochemically detectable") must be sought; such markers would easily enable the histologic diagnosis of LCNC.

Again, we pointed out that LCNC should be recognized as one of the poorest prognostic subgroups in lung cancer, and novel therapeutic approaches should be established. Further accumulation of clinicopathologic characteristics, especially the response to nonsurgical treatment, is the most important issue. The histologic classification of lung tumors with neuroendocrine phenotype also seems to need further revision in the near future.

Hidefumi Takei, MD
Hisao Asamura, MD
Yoshihiro Matsuno, MD
National Cancer Center Hospital
Tokyo, Japan