treated with the high-affinity tissue angiotensin-converting enzyme (ACE) inhibitor quinaprilat had significantly better recovery of regional wall motion, a lower incidence of ventricular arrhythmias, and a smaller infarct size than controls. When the low-affinity tissue ACE inhibitor enalaprilat was administered, the decrease in infarct size was less (24% enalaprilat vs 14% quinaprilat; $P < .0001$). The beneficial effects of quinaprilat's action could be explained by its superior preservation of endothelial function, since quinaprilat-treated hearts had significantly greater coronary arterial relaxation in response to bradykinin (66% quinaprilat vs 32% enalaprilat; $P < .0001$). When the bradykinin antagonist HOE140 was added to the quinaprilat-treated hearts, the protective effect on endothelial function was abolished and these hearts had a significant increase in infarct size (48% quinaprilat + HOE vs 14% quinaprilat; $P < .0001$).

I agree that ACE inhibitors enhance myocardial protection during cardioplegic arrest. However, ACE inhibitors are not all equally protective. ACE inhibitors with high tissue ACE inhibition such as quinaprilat result in better preservation of endothelial function, which decreases ischemic necrosis. This improvement in endothelial function appears to be bradykinin-mediated. I concur with Korn and coworkers that high-affinity tissue ACE inhibitors may play an important role in enhancing cardioplegic protection during coronary revascularization.^{2,3}

Harold L. Lazar, MD
Professor of Cardiothoracic Surgery
Boston Medical Center
Cardiothoracic Surgery
Boston, MA 02118

References

1. Korn P, Kröner A, Schirmer J, Hallström S, Bernecker O, Mallinger R, et al. Quinaprilat during cardioplegic arrest in the rabbit to prevent ischemia-reperfusion injury. *J Thorac Cardiovasc Surg.* 2002;124:352-60.
2. Lazar HL, Bao Y, Rivers S, Colton S, Bernard SA. High tissue affinity angiotensin-converting enzyme inhibitors improve endothelial function and reduce infarct size. *Ann Thorac Surg.* 2001;72:548-54.
3. Lazar HL, Bao Y, Rivers S, Bernard SA. Pretreatment with angiotensin-converting inhibitors attenuates ischemic reperfusion injury during coronary revascularization. *Ann Thorac Surg.* 2002;73:1522-7.
doi:10.1067/mtc.2003.337

Tracheal replacement

To the Editor:

It is encouraging that efforts continue to produce a "tissue-engineered trachea."¹ since this is about the only technique of the many attempts at tracheal replacement by a variety of approaches over more than 50 years that seems to offer any real promise.² I question, however, whether the experiments reported succeeded in forming "the structural component of a functional tracheal replacement."

Although visually resembling a trachea, the construction described failed to function (in sheep) over 5 and 7 days in 2 animals (with resultant malacia) or longer than 2 days in 4 animals (with stenosis). The authors' findings suggest that inflammatory response might be responsible for degeneration of the graft. Cell density and hydroxyproline content support this hypothesis. Conversion toward scar probably had begun, even with survival of some cartilage. The authors also allude to future advisability of assuring blood supply, which may indeed be a critical factor in this early deterioration of the implant. One of the keys to failure, even of autotransplanted free tracheal segments of any significant length, has been failure of blood supply, despite muscular or omental pedicles. Very short tracheal grafts, of course, are of no potential clinical use.

A subject that is deferred by the authors to later experimentation, perhaps with engineered epithelium, is the lack of epithelial coverage in these experiments. In time this would lead to graft obstruction by cicatrization. Epithelium in general has failed to recolonize any extended length of tracheal replacement by ingrowth from the normal tracheal ends alone.

I shall look forward with interest to future reports by this and other groups using tissue engineering in an effort to meet the challenge of fashioning a usable tracheal prosthesis for the rare occasion when it is indeed needed.

Hermes C. Grillo, MD
Division of Thoracic Surgery
Massachusetts General Hospital
32 Fruit St
Boston, MA 02114

References

1. Kojima K, Bonassar LJ, Roy AK, Vacanti CA, Cortiella J. Autologous tissue-engineered trachea with sheep nasal chondro-

cytes. *J Thorac Cardiovasc Surg.* 2002;123:1177-84.

2. Grillo HC. Tracheal replacement: a critical review. *Ann Thorac Surg.* 2002;73:1995-2004.

doi:10.1067/mtc.2003.260

Cardiac herniation after intrapericardial pneumonectomy

To the Editor:

The brief communication by Baisi and associates¹ reporting a case of cardiac herniation after intrapericardial pneumonectomy is almost identical with a case report in another journal.² Both reports fail to mention that chest tubes were deployed after the pneumonectomy. (They can clearly be seen in their respective illustrations.)

The previous case^{2,3} instigated a spirited correspondence^{4,5} centering on the role, if any, of chest tubes after pneumonectomy and their relation to cardiac herniation. The current case also fails to mention a chest tube in the left posterior thoracic gutter clearly seen on the computed tomographic scan (see Figure 1). No mention is made of its presence or mode of use. Rather the authors emphasize (without justification or substantiation in my mind) that the induction chemotherapy was to blame. Actually the initial closure of the postpneumonectomy pericardial defect failed because the sutures, placed under tension, cut through—a mechanical problem, not a neoplastic or chemotherapeutic problem.

The empty postpneumonectomy space is not to be equated with that containing variable amounts of lung tissue, which require chest tubes attached to suction. Using such continued suction in a postpneumonectomy space even for a short period of time is to court disaster produced by excessive mediastinal shift and, in some cases, cardiac herniation. Despite the opinion of Walker⁴ of the hazards of chest tubes after pneumonectomy, Cooley⁵ expressed it well when he said that he routinely used tubes after pneumonectomy but that "when not in use the tube should be clamped." [italics mine.]

Paul A. Kirschner, MD, FACS
Department of Cardiothoracic Surgery
Box 1028
Mount Sinai Medical Center
1 Gustave L. Levy Place
New York, NY 10029