could emphasize. Second is the increasing life-saving role of these devices as access to transplants decreases. Third is the fact that all LVADs are still subject to the same entry criteria that were used in the 1970s for patients with acute failure-to-wean-from-bypass. These hemodynamic entry criteria should be reassessed in view of the changing medical therapy and the fact that these implantable pumps are for patients with chronic heart failure on transplant waiting lists who experience worsening of heart failure made manifest primarily by end-organ failure and not by acute hemodynamic compromise.

O. H. Frazier, MD

Director, Cardiopulmonary Transplantation
Texas Heart Institute
PO Box 20345
Houston, TX 77225-0345

References

1. Copeland J. Multicenter bridge to transplantation with the HeartMate assist device: Evaluation from another perspective. *J Thorac Cardiovasc Surg.* 2003; 125:228-30.
2. Frazier OH, Rose EA, Oz MC, Dembitsky W, McCarthy P, Radovancevic B, et al. Multicenter clinical evaluation of the HeartMate vented electric left ventricular assist system in patients awaiting heart transplantation. *J Thorac Cardiovasc Surg.* 2001; 122:1186-95.

doi:10.1067/mtc.2003.69

Cerebral protection with retrograde cerebral perfusion

To the Editor:

In their article Bonser and colleagues¹ claim that the cerebral flow resulting from retrograde cerebral perfusion (RCP) is insufficient to make a major contribution to cerebral oxygenation and that the brain remains ischemic. In the discussion of their impressive work, the authors stated correctly that current data suggest that RCP might provide only little brain perfusion in human subjects compared with antegrade baseline cerebral flow. Should we abandon this adjunct to hypothermic circulatory arrest, or should we explore methods to maximize the benefits from its application? The effort to improve the effectiveness of RCP during hypothermic circulatory arrest should combine reduction of the metabolic activity with simultaneous increase of blood flow to fulfill the residual metabolic requirements of the brain. In a small series

my colleagues and I² demonstrated that the introduction of vasodilators and anesthetics into the retrograde perfusate tripled RCP flow without increase in pressure and with concomitant suppression of electric activity. In the present study it is not stated whether brain electric activity was monitored, and the individual RCP flow is not specified. Simple pharmacologic manipulations might augment RCP to 20% or 30% of cerebral flow instead of only 10%. The administration of potent anesthetics or other neuroprotective agents might mitigate the metabolic needs of the brain. Such maneuvers will render the brain less ischemic, allowing safer circulatory arrest.

Amir Elami, MD

Department of Cardiothoracic Surgery
Hadassah University Hospital
PO Box 12000
Jerusalem 91120, Israel

References

1. Bonser RS, Wong CH, Harrington D, Pagano D, Wilkes M, Clutton-Brock T, et al. Failure of retrograde cerebral perfusion to attenuate metabolic changes associated with hypothermic circulatory arrest. *J Thorac Cardiovasc Surg.* 2002;123:943-50.
2. Shenkman Z, Elami A, Weiss YG, Glantz L, Milgalter E, Drenger B, et al. Cerebral protection using retrograde cerebral perfusion during profound hypothermic circulatory arrest. *Can J Anaesth.* 1997;44:1096-101.

doi:10.1067/mtc.2003.246

Reply to the Editor:

We thank Elami for his comments concerning our work on retrograde cerebral perfusion (RCP).¹ We acknowledged in our discussion that RCP could contribute to metabolic support during hypothermic circulatory arrest and that this contribution could be increased by the use of vasodilators. It is also possible that the use of anesthetics or neuroprotective agents during RCP might afford neuroprotection. The series quoted by Elami, however, describes only anecdotal evidence from 3 patients undergoing differing pharmacologic interventions.² This report described no monitoring of cerebral metabolism or perfusion, therefore making inferences regarding cerebral metabolic activity and blood flow unreliable. We have previously reported that RCP can provide some brain perfusion, but we have questioned what fraction of retrograde flow provides true brain perfusion.^{3,4} It is quite possible that increased

flow achieved by means of vasodilator administration bypasses the brain through venovenous and venoarterial collaterals.⁴ Any assumption that increasing flow increases true brain perfusion and neuroprotection requires rigorous enquiry. We would agree that the jury remains out on RCP and that a beneficial clinical effect remains to be proven. Such therapeutic interventions therefore require assessment as part of randomized controlled trials if their efficacy is to be completely investigated.

Robert Bonser, FRCP, FRCS

Deborah Harrington, MRCS

Domenico Pagano, MD, FRCS

Department of Cardiothoracic Surgery
Queen Elizabeth Hospital
Edgbaston
Birmingham B15 2TH, United Kingdom

References

1. Bonser RS, Wong CH, Harrington D, Pagano D, Wilkes M, Clutton-Brock T, et al. Failure of retrograde cerebral perfusion to attenuate metabolic changes associated with hypothermic circulatory arrest. *J Thorac Cardiovasc Surg.* 2002;123:943-50.
2. Shenkman Z, Elami A, Weiss YG, Glantz L, Milgalter E, Drenger B, et al. Cerebral protection using retrograde cerebral perfusion during profound hypothermic circulatory arrest. *Can J Anaesth.* 1997;44:1096-101.
3. Pagano D, Boivin CM, Farouqi MH, Bonser RS. Retrograde perfusion through the superior vena cava perfuses the brain in human beings. *J Thorac Cardiovasc Surg.* 1996; 111:270-2.
4. Wong C, Bonser RS. Retrograde perfusion and true reverse brain blood flow in humans. *Eur J Cardiothorac Surg.* 2000;17: 597-601.

doi:10.1067/mtc.2003.245

Angiogenesis by means of endothelial cell transplantation

To the Editor:

In their article, "Angiogenesis by Endothelial Cell Transplantation," Kim and associates¹ present results of a novel and intriguing approach to angiogenic therapy. Their preliminary report that endothelial cell transplantation into a myocardial scar accelerates angiogenesis suggests a promising alternative to angiogenic gene or protein therapy. About the same time that their article was published, we² presented similar results before the American Heart Association at our annual scientific session (Anaheim, Calif) in November 2001.