Interestingly, we created our model of ischemic cardiomyopathy by placing an ameroid constrictor on the branch of the circumflex artery in 12 adult sheep.

In our experiment, 4 weeks after constriction, we randomly assigned 3 groups (4 animals per group) to transpericardial injection (3 sites per animal): sham treatment (saline solution with denatured cells), control (no injections), and endothelial cell fibrin meshwork groups (autologous mature endothelial cells, 5×10^5 per injection site harvested from jugular veins). Eight weeks after transepicardial treatment, animals were assessed for myocardial function (by means of echocardiography), myocardial blood flow (MBF), and capillary area (by means of histology).

We demonstrated improved capillary density and MBF in keeping with their speculation that, "A coronary ligation model of myocardial infarction, in which a more extensive peri-infarct border zone could harbor viable but hypoperfused cardiomyocytes and endothelial cells, might result in greater angiogenesis."

One month after constriction, MBF in the ischemic myocardium decreased to 70% of baseline value ($P < .001$). Two months after reoperation and the beginning of treatment, animals were killed, and MBF was found to have decreased to 28% versus baseline values in the control and saline solution groups ($P < .001$). MBF in the endothelial cell group significantly increased to 108%.

In normal nonischemic myocardium in all 3 groups, there were 205.54 ± 12.31 capillaries/mm². Three months after constriction (and 2 months after reoperation and beginning of treatment) in the control and saline solution groups, there was a decrease up to 51% in capillaries per square millimeter ($P < .001$ vs baseline). However, although this decreased significantly in the control and saline solution groups versus baseline values, in the endothelial cell group this increased to 146% ($P < .001$). Hence all of our results confirmed those in the report of Kim and associates.¹

However, they also concluded that "increasing blood flow by endothelial cell transplantation in this model did not improve contractility or prevent ventricular dilatation," and "we plan to perform autol-

ogous endothelial cell transplantation in our clinically relevant porcine model of occlusion of the left anterior descending coronary artery. We anticipate that angiogenesis may improve regional and global function."

In our investigation, 4 weeks after constriction, left ventricular ejection fraction (LVEF) decreased to 75% ($P < .001$). Two months later, LVEF decreased to 57% in the control and saline groups ($P < .001$ vs 4 weeks and baseline). In the endothelial cell group LVEF increased to 89% of baseline values but continued to be less than baseline value ($P < .05$). However, LVEF in this group was significantly better than in the control and saline solution groups ($P < .001$).

On the basis of our data, we think that use of endothelial cell transplantation within a 3-dimensional matrix system can be used to induce angiogenesis in patients who cannot be adequately revascularized by using standard techniques. Exploring autologous mature endothelial cell transplantation might become a useful alternative to gene and protein therapy. We recently completed studies involving a large number of animals; these results are pending.

Valeri Chekanov, MD, PhD^a

Nicholas Kipshidze, MD, PhD^b

Milwaukee Heart Institute

Aurora/Sinai Medical Center

Milwaukee, Wis^a

Lenox Hill Heart & Vascular Institute

Cardiovascular Research Foundation

Lenox Hill Hospital

New York, NY^b

References

1. Kim E-J, Li R-K, Weisel RD, Mickle DA, Jia ZQ, Tomita S, et al. Angiogenesis by endothelial cell transplantation. *J Thorac Cardiovasc Surg.* 2001;122:963-71.
2. Chekanov V, Nikolaychik V, Tchekanov G, Dargas G, Roubin G, Leon M, et al. Transplantation of autologous endothelial cells in a fibrin meshwork induces angiogenesis and improves myocardial function in an animal model of ischemic cardiomyopathy [abstract]. *Circulation.* 2001;104(suppl 17):II260.

doi:10.1067/mtc.2003.248

Reply to the Editor:

Chekanov and Kipshidze performed a series of experiments that appear to confirm our findings that endothelial cell transplan-

tation improved perfusion after myocardial injury. Recently, a variety of techniques have been used to induce angiogenesis to treat patients who are not candidates for conventional methods of revascularization. Cell transplantation has the advantage of inducing angiogenesis by a variety of mechanisms because the technique is not limited to the release of a single protein.

Chekanov and Kipshidze used an endothelial cell fibrin meshwork applied to the epicardial surface of an ischemic region. They found that this technique induced angiogenesis. The mode of delivery and the number of cells engrafted might critically affect the benefits achieved with cell transplantation. The optimal conditions have not been well defined. In addition, the cell type that induces the most beneficial response has not been adequately established. Endothelial cells induce an extensive capillary network, but they might not induce the formation of sufficient conduit vessels. Further research into this important area might establish which cell type, which cell number, and which mode of delivery are optimal to induce angiogenesis.

Ren-Ke Li, MD

Richard D. Weisel, MD

Department of Cardiovascular Surgery

Toronto General Hospital

200 Elizabeth St

Toronto, Ontario M5G 2C4, Canada

doi:10.1067/mtc.2003.249

Missing data from mesothelioma study

To the Editor:

The phase II trial of surgery and radiation of malignant mesothelioma by Rusch and colleagues¹ is an important addition to the literature, documenting contemporary efforts to control and eventually cure this so far incurable malignant process. The 8% mortality after extrapleural pneumonectomy reflects the tremendous strides in surgical techniques effected in recent decades. The rigorous staging confirms a major attempt to bring better science to this capricious disease.

Nevertheless, Rusch and colleagues¹ have excluded a host of important information from their publication. I hope that they will be able to strengthen the report by rectifying some of these omissions: