entity. There are 4.6 million people in the United States with heart failure today, and 550,000 new cases are being reported annually by the American Heart Association.⁵ Approximately 30% to 50% of patients with heart failure have a normal or nearly normal left ventricular ejection fraction.⁶ CHF is a leading cause of cardiac morbidity and mortality from cardiovascular disease. Although left ventricular diastolic dysfunction occurs in all patients with systolic dysfunction and CHF, a third of patients have CHF with isolated diastolic dysfunction. In the general population, the mortality among patients with diastolic heart failure is 4 times that among persons without heart failure but half that among patients with systolic heart failure. It is widely known that many physicians underappreciate diastolic function in heart failure.

Reinterpreting the data in the context of the discussion asks for other explanations. In the present era of advancements in echocardiography, diastolic dysfunction needs to be taken into consideration, especially when symptoms of congestive heart failure are present. If that is the case in this study, left ventricular dysfunction remains a predictor of outcome but with a change from systolic to diastolic. Finally, I ask Davierwala and colleagues to take a second look at their data in the light of this forgotten predictor variable. I would appreciate their feedback.

Jeffrey H. Shuhaiber, MD
University of Illinois at Chicago
Chicago, IL 60612

References

1. Davierwala PM, Maganti M, Yau TM. Decreasing significance of left ventricular dysfunction and reoperative surgery in predicting coronary artery bypass grafting-associated mortality: a twelve-year study. *J Thorac Cardiovasc Surg.* 2003;126:1335-44.
2. Kurki TS, Kataja M. Preoperative prediction of postoperative morbidity in coronary artery bypass grafting. *Ann Thoracic Surg.* 1996;61:1740-5.
3. Christakis GT, Ivanov J, Weisel RD, Birnbaum PL, David TE, Salerno TA. The changing pattern of coronary artery bypass surgery. *Circulation.* 1989;80(3 Pt 1):1151-61.
4. Kitzman DW, Little WC, Brubaker PH, Anderson RT, Hundley WG, Marburger CT, et al. Pathophysiological characterization of isolated diastolic heart failure in comparison to systolic heart failure. *JAMA.* 2002;288:2144-50.
5. American Heart Association. 2000 Heart and stroke statistical update. Dallas: The Association; 2000.
6. Vasan RS, Benjamin EJ. Diastolic heart failure—no time to relax. *N Engl J Med.* 2001;344:56-9.
doi:10.1016/j.jtcvs.2003.12.043

Reply to the Editor:

Dr Shuhaiber correctly points out that left ventricular (LV) diastolic dysfunction is frequently superimposed on either normal or reduced systolic function in patients undergoing coronary artery bypass grafting. The importance of diastolic dysfunction in the development and progression of congestive heart failure (CHF) has been illustrated previously¹ and is highlighted by the divergent trends in the prevalence of LV systolic dysfunction and CHF in the series on which we reported. Early echocardiographic signs of impaired LV relaxation include decreased early transmitral LV filling and greater dependence on atrial contraction.² Pseudonormalization of LV filling is observed as LV stiffness and left atrial pressures continue to increase, and finally severe diastolic dysfunction is marked by rapid early filling, extreme LV stiffness, and elevated diastolic pressures. This diastolic dysfunction has been shown to predict outcomes in patients with CHF independently of systolic function.³

Despite this, echocardiographic characterization and quantitation of LV diastolic dysfunction is not performed and recorded for all patients undergoing coronary surgery. This may be due in part to logistical issues and in part to underappreciation of its prognostic significance. Like systolic function, diastolic function may also vary significantly with time in patients with intermittent ischemia or a maturing myocardial infarction.

Our institutional database, initiated in 1982 and now encompassing approximately 40,000 surgical patients, does not as yet contain uniform data on diastolic function, nor does the much larger Society of Thoracic Surgeons database. Thus, although one may reasonably speculate that the prevalence of diastolic dysfunction in patients undergoing coronary artery bypass grafting is increasing in parallel with other risk factors, as we have reported in this series, our data do not currently permit evaluation of the independent effect of diastolic function on mortality and morbidity. These measures of diastolic dysfunction

will have to be obtained routinely for all patients undergoing surgery and incorporated into institutional and regional databases before we be able to determine whether the decreasing influence of LV systolic dysfunction on outcomes has unmasked relatively more subtle predictors, including CHF and left main stenosis as we have reported or diastolic dysfunction as Dr Shuhaiber suggests.

Piroze M. Davierwala, MD
Terrence M. Yau, MD, MSc
Division of Cardiovascular Surgery
Toronto General Hospital
Toronto, Ontario, Canada

References

1. Gaasch WH. Diagnosis and treatment of heart failure based on left ventricular systolic or diastolic dysfunction. *JAMA.* 1994;271:1276-80.
2. Rakowski H, Appleton C, Chan KL, Dumesnil JG, Honos G, Jue J, et al. Canadian consensus recommendations for the measurement and reporting of diastolic dysfunction by echocardiography: from the Investigators of Consensus on Diastolic Dysfunction by Echocardiography. *J Am Soc Echocardiogr.* 1996;9:736-60.
3. Vasan RS, Benjamin EJ, Levy D. Prevalence, clinical features and prognosis of diastolic heart failure: an epidemiologic perspective. *J Am Coll Cardiol.* 1995;26:1565-74.
doi:10.1016/j.jtcvs.2004.01.026

Extracapsular lymph node involvement in esophageal cancer and number of involved nodes

To the Editor:

We thank Dr DeMeester¹ for his valuable comments on our article "Extracapsular Lymph Node Involvement Is a Negative Prognostic Factor in T3 Adenocarcinoma of the Distal Esophagus and Gastroesophageal Junction."² We fully agree with Dr DeMeester¹ that the number and characteristics of lymph nodes are very important determinants of survival in esophageal cancer, and we endorse his plea for taking the number of involved nodes into account in the TNM staging system of esophageal cancer.

In his editorial,¹ Dr DeMeester made some points that we would like to clarify. With regard to the survival in patients without nodal involvement versus patients with intracapsular nodal involvement, there was no significant difference in survival. This

does not, however, mean that they had the same survival. Applying a Cox regression on these patients, we had a *P* value of .435 and a hazard ratio of 1.271 (95% confidence interval 0.696-2.322), which means that patients with intracapsular involved lymph nodes have a 27% higher risk of dying than did those with pN0 disease. This hazard ratio is not extreme relative to the hazard ratio of 3.37 in patients with lymph node involvement. There were sufficient patients at risk in each time interval: 33, 29, 24, 16, 15, and 12 for the pN0 group and 60, 53, 36, 25, 17, and 14 for the intracapsular group at 0, 1, 2, 3, 4, and 5 years, respectively.

The difference we found regarding the mean and median number of involved lymph nodes (3.68 and 3.0 for intracapsular vs 7.96 and 6.0 for extracapsular) was mainly caused by some extreme values in the extracapsular group, which could indicate that extracapsular involvement points to more advanced disease. However, we checked the interaction between lymph node status and number of involved lymph nodes by analysis of variance and ruled out significant interaction between these variables, so both can be considered as independent effects.

As to the subgroup of patients with exactly one intracapsular versus one extracapsular involved lymph node, we agree that a type II error is possible because of the small group. The reason we included this graph was merely to demonstrate the effect of intracapsular versus extracapsular involvement after having neutralized the effect of number of positive lymph nodes. Because of the small number of patients, we couldn't conclude that patients with one intracapsular lymph node have a better survival than patients with no involved lymph nodes. This graph merely shows the detrimental effect of extracapsular involvement on survival.

Again, we would like to highlight the importance of extracapsular involvement, which seems far greater than the effect of number of involved lymph nodes. As our model indicates, the effect of extracapsular involvement (237%) is much greater than the cumulative effect of, for example, 12 positive nodes ($12 \times 8.6\% = 103.2\%$). For this reason we believe that in addition to the number of involved lymph nodes, lymph node status (intracapsular vs extracapsular) should be routinely incorporated

in every pathology report. If our results are confirmed by other centers, then perhaps extracapsular involvement should be incorporated in the TNM staging system as well, unless indeed biologic and molecular biological markers, the Holy Grail of prognostic indicators, are able to replace the actual routine staging.

In conclusion, if you see one, ask your pathologist to look at the capsula.

T. Lerut, MD, PhD

J. Moons, CNS

S. Fieuws, MSc

Department of Thoracic Surgery

Catholic University Leuven

UZ Gasthuisberg

Herestraat 49

3000 Leuven, Belgium

References

1. DeMeester SR. Lymph node involvement in esophageal adenocarcinoma: if you see one, have you seen them all? *J Thorac Cardiovasc Surg.* 2003;126:947-9.
2. Lerut T, Coosemans W, Decker G, De Leyn P, Ectors N, Fieuws S, et al. Extracapsular lymph node involvement is a negative prognostic factor in T3 adenocarcinoma of the distal esophagus and gastroesophageal junction. *J Thorac Cardiovasc Surg.* 2003;126:1121-8.

doi:10.1016/j.jtcvs.2004.01.027

Skeletonized bilateral internal thoracic arteries in patients with diabetes: Additional advantages and concerns

To the Editor:

The article by Peterson and colleagues¹ published in the November 2003 issue of the *Journal* is clearly an important study demonstrating the beneficial impact of using skeletonized bilateral internal thoracic artery (ITA) grafts on the incidence of sternal wound infection in patients with diabetes. In terms of morbidity, sternal dehiscence as a consequence of sternal devascularization is the most worrying potential complication of bilateral ITA grafts, particularly in patients with diabetes. Patients with diabetes are those with potentially most to gain from bilateral ITA grafts, because they often have more severe, diffuse, and distal disease. Appropriate patient selection and modification of the technique for harvesting the ITA can significantly reduce the risk of impaired

wound healing in these patients. Harvesting the ITA with the traditional pedicled technique also involves harvest of surrounding parietal pleura, venae comitantes, muscle, and fascia, leaving a completely denuded and devascularized strip of chest wall. Harvesting only the ITA itself ("skeletonizing" technique) preserves intercostal collateral vessels and some sternal blood supply. Not only does the skeletonization technique result in superior sternal blood flow preservation² and reduce risk of wound healing in all patients³ and those with diabetes in particular,^{4,5} it also offers several additional advantages. These include aiding judgment of graft length, provision of extra length (an important consideration if the ITA is to be used for more than a single graft, as a sequential graft), allowing thorough visual inspection to identify spastic or damaged areas that could otherwise be obscured by perivascular fat, facilitating sequential anastomoses and composite arterial grafting (because the venae comitantes, perivascular fat, and areolar tissue do not obscure the anastomosis), and minimizing the risks of kinks or twists in the conduit.

There are, however, several points of caution to be emphasized with the skeletonization technique. It is technically more demanding than pedicle harvesting, and careful and meticulous dissection is required. Skeletonization takes longer than does the pedicle technique and has a definite learning curve. Small arterial branches can be inadvertently avulsed by excessive downward retraction, resulting in injury to the artery and compromising long-term patency. Finally, skeletonization probably induces a greater degree of spasm.

Perhaps what is missing from this particular study is the incidence of ITA injury sustained during skeletonization and the fate of any injured ITAs. Also, it would be interesting to know about the authors' experience of ITA spasm resulting from skeletonization and any measures they took to deal with it.

Shahzad G. Raja, MRCS

Department of Cardiac Surgery

Alder Hey Children's Hospital

Liverpool, United Kingdom

References

1. Peterson MD, Borger MA, Rao V, Peniston CM, Feindel CM. Skeletonization of bilateral internal thoracic artery grafts lowers the risk of sternal infection in patients with diabe-