- tes. *J Thorac Cardiovasc Surg.* 2003;126:1314-9.
2. Cohen AJ, Lockman J, Lorberboym M, Bder O, Cohen N, Medalion B, et al. Assessment of sternal vascularity with single photon emission computed tomography after harvesting of the internal thoracic artery. *J Thorac Cardiovasc Surg.* 1999;118:496-502.
 3. Calafiore AM, Vitolla G, Iaco AL, Fino C, Di Giammarco G, Marchesani F, et al. Bilateral internal mammary artery grafting: midterm results of pedicled versus skeletonized conduits. *Ann Thorac Surg.* 1999;67:1637-42.
 4. Sofer D, Gurevitch J, Shapira I, Paz Y, Matsa M, Kramer A, et al. Sternal wound infections in patients after coronary artery bypass grafting using bilateral skeletonized internal mammary arteries. *Ann Surg.* 1999;229:585-90.
 5. Matsa M, Paz Y, Gurevitch J, Shapira I, Kramer A, Pevny D, et al. Bilateral skeletonized internal thoracic artery grafts in patients with diabetes mellitus. *J Thorac Cardiovasc Surg.* 2001;121:668-74.
doi:10.1016/j.jtcvs.2003.12.044

Reply to the Editor:

We thank Dr Raja for his supportive comments and we agree with the additional advantages offered by skeletonizing the internal thoracic artery (ITA) for grafting stated in his letter. In particular, the additional length gained by skeletonization is extremely helpful when the right ITA is used as an in situ graft to obtuse marginal branches of the circumflex coronary artery.¹

Dr Raja correctly notes that skeletonization of ITAs requires more technical expertise and time than does conventional pedicled harvesting. Although the total operative times were similar for the skeletonized and conventional groups in our study,² skeletonization may add 10 to 20 minutes to the ITA harvesting time. The surgeons in our study felt proficient with skeletonization after approximately 20 ITA harvests. Injury to the ITA during this learning phase is a concern, but we did not record the frequency of ITA injury in our database. If an ITA graft is injured during harvesting, we can occasionally utilize the undamaged segment as a free graft. The surgeons who routinely perform skeletonization at our institution report a very low (<1%) incidence of ITA injury. Unfortunately, most studies in the literature do not report the incidence of injury to the ITA graft during skeletonization; however,

Nishida and colleagues³ reported an incidence of 0.7%.

Dr Raja also raises the concern that manipulation of the ITA graft during skeletonization may contribute to an increased propensity toward graft spasm. Spasm of the ITA graft has been reported in the literature, and some institutions inject the ITA with papaverine to prevent spasm. However, intraluminal injection of vasodilating agents can itself lead to graft injury by pressurizing the artery, resulting in endothelial damage or dissection. Instead, we prefer to bathe the skeletonized ITAs with topical papaverine. Although we do not actively record the incidence of postoperative ITA spasm in our institution, we do know that it is uncommon.

We believe that meticulous ITA skeletonization offers many advantages relative to pedicled harvesting, with an acceptably low risk of complications.

Mark D. Peterson, MD^{a,b}

Michael A. Borger, MD, PhD^{a,b}

Division of Cardiac Surgery

Toronto General Hospital^a

Toronto, Ontario, Canada

Department of Surgery

University of Toronto^b

Toronto, Ontario, Canada

References

1. Bonacchi M, Prifti E, Battaglia F, Frati G, Sani G, Popoff G. In situ retrocaval skeletonized right internal thoracic artery anastomosed to the circumflex system via transverse sinus: technical aspects and postoperative outcome. *J Thorac Cardiovasc Surg.* 2003;126:1302-13.
2. Peterson MD, Borger MA, Rao V, Peniston CM, Feindel CM. Skeletonization of bilateral internal thoracic artery grafts lowers the risk of sternal infection in patients with diabetes. *J Thorac Cardiovasc Surg.* 2003;126:1314-9.
3. Nishida H, Tomizawa Y, Endo M, Koyanagi H, Kasanuki H. Coronary artery bypass with only in situ bilateral internal thoracic arteries and right gastroepiploic artery. *Circulation.* 2001;104(12 Suppl 1):176-80.
doi:10.1016/j.jtcvs.2004.01.028

A lost opportunity

To the Editor:

In the article "Neurodevelopmental Status at Eight Years in Children with Dextro-transposition of the Great Arteries: The Boston Circulatory Arrest Trial," Bellinger and colleagues¹ presented a sophisticated prospective randomized study, with an ex-

tensive data collection and analysis. However, they limited their investigations to "the two major methods of vital organ support, total circulatory arrest and low-flow cardiopulmonary bypass." I respectfully disagree with Bellinger and colleagues' decision, because they wasted a good opportunity by using a tremendous amount of effort and expertise to evaluate two methods already proven to provide suboptimal results. In my opinion, comparing total circulatory arrest and low-flow cardiopulmonary bypass is like comparing two different brands of cigarettes to evaluate whether one is less dangerous than the other.

There are already too many experimental and clinical studies reported in the literature on how to reduce the negative effects of hypothermia or flow reduction: metabolic derangement; endothelial lesions; and vascular, myocardial, neurologic, hematologic, and respiratory impairment.^{2,3} Instead of investing their extraordinary resources in this type of research, Bellinger and colleagues could explore an alternative method of cardiopulmonary bypass, even if they do not consider it to be one of the "two major methods of vital organ support": normothermic high-flow cardiopulmonary bypass.^{3,4} This method has already been supported by clinical experience with more than 1600 cases of surgery for congenital heart disease, including the arterial switch operation for neonates with transposition of the great arteries.⁴

The readers involved with the surgical treatment of children with congenital heart disease should be aware that we as surgeons are not obliged to choose between suboptimal methods of cardiopulmonary bypass, much as we are not obliged to choose among different brands of cigarettes if we are not smokers. There is the proven possibility of performing cardiopulmonary perfusion much closer to the physiologic state,⁴ and research units like the one of this study could use their resources to compare the "two major methods of vital organ support" with a more physiologically compatible method.

Antonio F. Corno, MD, FRCS, FETCS
Cardiovascular Surgery

Centre Hospitalier Universitaire Vaudois
(CHUV)

46 rue du Bugnon
CH-1011, Lausanne, Switzerland

References

1. Bellinger DC, Wypij D, duPlessis AJ, Rapaport LA, Jonas RA, Wernovsky G, et al. Neurodevelopmental status at eight years in children with dextro-transposition of the great arteries: the Boston Circulatory Arrest Trial. *J Thorac Cardiovasc Surg.* 2003;126:1385-96.
2. Corno AF, von Segesser LK. Is hypothermia necessary in pediatric cardiac surgery? *Eur J Cardiothorac Surg.* 1999;15:110-1.
3. Corno AF. What are the best temperature, flow, and hematocrit levels for pediatric cardiopulmonary bypass? *J Thorac Cardiovasc Surg.* 2002;124:856-7.
4. Durandy Y, Hulin S, Lecompte Y. Normothermic cardiopulmonary bypass in pediatric surgery. *J Thorac Cardiovasc Surg.* 2002;123:194.

doi:10.1016/j.jtcvs.2004.01.032

Reply to the Editor:

We thank Dr Corno for his suggestion that normothermic high-flow cardiopulmonary bypass would be likely to improve neurodevelopmental outcome relative to circulatory arrest or hypothermic reduced flow bypass, the two techniques that we studied. We agree with Dr Corno that many perfusion variables may influence late developmental outcome, and we recognize that few of these have been studied in detail. The Boston Circulatory Arrest Study, the subject of our recent report, was designed in 1987. Since that time, our laboratory studies have led us to undertake two subsequent prospective randomized clinical trials, one studying the role of pH¹ and one studying the role of hematocrit in influencing outcome after cardiopulmonary bypass.² Both

trials confirmed our laboratory results and resulted in important changes in our technique of bypass. However, our more recent laboratory studies investigating different temperature strategies for cardiopulmonary bypass have suggested that there is a disadvantage in using a higher temperature for cardiopulmonary bypass. For example, in a study that used intravital microscopy for direct examination of the cerebral microcirculation, we observed greater activation of white cells with increased numbers of rolling and adherent white cells when cardiopulmonary bypass was conducted at 34°C than at 15°C.³ We believe that this result reflects the fact that, despite Dr Corno's assertion to the contrary, cardiopulmonary bypass remains far from being a physiologic state. It is widely known that cardiopulmonary bypass is associated with multiple systemic inflammatory effects, both humoral and cellular. Our laboratory study has confirmed that a higher temperature exacerbates this inflammatory response. In addition, the high flow rate that is required to support the greater metabolic rate associated with a higher temperature also carries many disadvantages, including increased generation of emboli, increased left heart return necessitating greater suction volumes, reduced intracardiac exposure because of the need for multiple cannulas and increased left heart return, less effective myocardial protection with need for multiple reinfusions of cardioplegia (demonstrated to be a disadvantage in the neonate), and a reduced margin for safety in the event of equipment failure. The the-

oretical disadvantages of increased infection and bleeding that might result from hypothermia have not been confirmed in the neonate and infant.

In conclusion, we hope that Dr Corno and his associates will have the opportunity in the future to undertake developmental studies of their many patients who undergo normothermic high-flow bypass. Our own clinical studies will continue to be guided by translational research undertaken in the laboratory, which to date has not indicated any advantage for normothermic high-flow bypass.

Richard A. Jonas, MD
Jane W. Newburger, MD, MPH
David C. Bellinger, PhD, MSc
Children's Hospital Boston
Harvard Medical School
Boston, MA 02115

References

1. du Plessis AJ, Jonas RA, Wypij D, Hickey PR, Riviello J, Wessel DL, et al. Perioperative effects of alpha-stat versus pH-stat strategies for deep hypothermic cardiopulmonary bypass in infants. *J Thorac Cardiovasc Surg.* 1997;114:991-1000.
2. Jonas RA, Wypij D, Roth SJ, Bellinger DC, Visconti KJ, du Plessis AJ, et al. The influence of hemodilution on outcome after hypothermic cardiopulmonary bypass: results of a randomized trial in infants. *J Thorac Cardiovasc Surg.* 2003;126:1765-74.
3. Anttila V, Hagino I, Zurakowski D, Lidov HGW, Jonas RA. Higher bypass temperature correlates with increased white cell activation in the cerebral microcirculation. *J Thorac Cardiovasc Surg.* 2004;127:1781-8.
doi:10.1016/j.jtcvs.2004.01.033