and practices and (2) the outcome concerns in women and small patients.

Robert H. Habib, PhD^{ab}
 Anwar Zacharias, MD^{ac}
 Thomas A. Schwann, MD^{ac}
 Christopher J. Riordan, MD^{ac}
 Samuel J. Durham, MD^{ac}
 Aamir Shah, MD^{ac}
 Division of Cardiovascular Surgery^a
 St Vincent Mercy Medical Center
 Toledo, Ohio
 Departments of Medicine^b and
 Surgery^c
 Medical College of Ohio
 Toledo, Ohio

References

1. Guru V, Fremes SE, Tu JV. Time-related mortality for women after coronary artery bypass graft surgery: a population-based study. *J Thorac Cardiovasc Surg.* 2004;127(4):1158-65.
2. Koch CG, Khandwala F, Nussmeier N, Blackstone EH. Gender and outcomes after coronary artery bypass grafting: a propensity-matched comparison. *J Thorac Cardiovasc Surg.* 2003;126(6):2032-43.
3. Schwann TA, Habib RH, Zacharias A, et al. Effects of body size on operative, intermediate, and long-term outcomes after coronary artery bypass operation. *Ann Thorac Surg.* 2001;71(2):521-30.
4. DeFoe GR, Ross CS, Olmstead EM, et al. Lowest hematocrit on bypass and adverse outcomes associated with coronary artery bypass grafting. Northern New England Cardiovascular Disease Study Group. *Ann Thorac Surg.* 2001;71(3):769-76.
5. Habib RH, Zacharias A, Schwann TA, Riordan CJ, Durham SJ, Shah A. Adverse effects of low hematocrit during cardiopulmonary bypass in the adult: should current practice be changed? *J Thorac Cardiovasc Surg.* 2003;125(6):1438-50.

doi:10.1016/j.jtcvs.2004.04.018

Reply to the Editor:

We thank Dr Habib and colleagues for their interest in this topic. The issue of whether gender, size, or gender in combination with size can explain the worse early mortality after coronary artery bypass graft (CABG) surgery has been an ongoing debate. There still is no comprehensive evidence of what aspects of gender contribute to this increased early mortality risk; however, a strong association to body size has been previously described.¹ It also has been shown that the gender effect of worse early outcome rates of death and low output syn-

drome remain despite adjustment for body size, even when more detailed clinical data are available.¹ Size is important, but it is not the only factor that distinguishes men from women, and this certainly deserves more research. A recent article by Koch and coworkers² used propensity analysis to match women and men using 68 covariates. As seen in this study, only 26% of women from their entire cohort were matched, which certainly limited the generalizability of the study findings for the entire population of women undergoing CABG surgery. Regardless, likely there are factors other than size for women that lead to worse early outcomes after CABG surgery. There is some evidence to show that off-pump coronary artery bypass surgery might reduce adverse outcomes for women.³ Unfortunately, off-pump coronary artery bypass trials will always have a minority of women and thus not be powered to understand the gender effect fully. Perhaps studies understanding the differential gender response to cardiopulmonary bypass and myocardial ischemia are required to more fully understand why there is a gender difference. Retrospective studies, even with detailed clinical data, will never be sufficient to conclusively identify the cause behind this observation, and size only explains part of the gender story. We can all agree that it is time for the research community to look beyond simple gender differences and work in a concerted manner to try and identify the biologic factors that contribute to gender differences in CABG surgery outcomes.

Veena Guru, MD
 Stephen E. Fremes, MD, MSc
 Jack V. Tu, MD, PhD
 Institute for Clinical Evaluative Sciences
 Divisions of Cardiovascular Surgery and
 General Internal Medicine
 Sunnybrook and Women's College Health
 Sciences Centre
 University of Toronto
 Toronto, Ontario, Canada

References

1. Christakis GT, Weisel RD, Buth KJ, et al. Is body size the cause for poor outcomes of coronary artery bypass operations in women? *J Thorac Cardiovasc Surg.* 1995;110:1344-58.
2. Koch CG, Khandwala F, Nussmeier N, Blackstone EH. Gender and outcomes after coronary artery bypass grafting: a propensity-

matched comparison. *J Thorac Cardiovasc Surg.* 2003;126:2032-43.

3. Petro KR, Dullum MK, Garcia JM, et al. Minimally invasive coronary revascularization in women: a safe approach for a high-risk group. *Heart Surg Forum.* 2000;3:41-6.
 doi:10.1016/j.jtcvs.2004.04.018

Something to expect, diagnose, and treat early

To the Editor:

We read with interest the article entitled "Evolutional aspects of children and adolescents with surgically corrected aortic coarctation: clinical, echocardiographic, and magnetic resonance image analysis of 113 patients."¹ We learned that transverse aortic arch hypoplasia should be corrected concomitantly with coarctation.

Nothing about the chest radiograph or cardiothoracic ratio was mentioned. In the authors' echocardiographic analysis, they did not comment about the dimensions or functions of the cardiac chamber, although left ventricular hyperkinesia and increased ventricular mass are found even in normotensive patients long after successful coarctation repair: increased ventricular mass can provoke rhythm disturbances and might cause an imbalance between muscle and coronary vessel growth, with decreased oxygen reserve. Their series is a select group, and not all patients undergoing operations during this 26-year period were included. Therefore, the study is subject to biases and is weak. The definition of recoarctation is to be broadened to include the systolic gradient between the right arm and leg: measured with a sphygmomanometer, Doppler echocardiography, magnetic resonance angiography, and/or angiography of the aortic isthmus and descending aorta, systolic gradients and diameters equal to or greater than 20 mm Hg at rest or a combination of a right arm-leg pressure gradient of more than 50 mm Hg after the bicycle test and more than 40% reduction in cross-sectional diameter of the aorta at the anastomotic site compared with the aorta at the level of the diaphragm.²

The recoarctation percentage after end-to-end anastomosis (14%) is not correct: it should be 12.53% rather than 14%. The patch aortoplasty figure should be 7.69% rather than 7%. However, this did not affect the order of frequency. We do not understand why the aortic ratio median behaved independently of age at coarctation correction or why there was no significant