include the bilateral mediastinal nodal regions independently by their involvement. New data from RTOG trials⁴ suggest moving the elective nodal irradiation focalizing to the gross tumor volume. In our recent published report,⁵ exploring a combined modality treatment with limited irradiation volume, we have reported a grade 3 esophageal toxicity rate of 1.1%.

Moreover, surgical-related mortality, as the authors stated, could be attributed to the type of operation, pneumonectomy, more than to induction protocol. Also, data from neoadjuvant chemotherapy support these findings.⁶

Today, lymph node clearance⁷ and pathologic downstaging have been proposed as surrogate endpoints for neoadjuvant treatment in LA-NSCLC. Data from SAKK and the University of Pennsylvania agree to group together pathologic stage I and II (or pN0-1) disease as a predictor of overall survival and event-free or local recurrence-free survival. If we look at the Memorial Sloan Kettering data on 470 patients treated with neoadjuvant therapy,⁸ the multivariate analysis shows us a statistically significant difference in term of the relative rate of deaths between pathologic stage 0 and stage II to IV disease. No differences have been reported between pathologic stage 0 and stage I disease. According to this and our analysis, we share the hypothesis to use pathologic downstaging to stage 0 to stage I as a direct indicator of the effectiveness of any multimodality approach to LA-NSCLC.

Finally, 2 more issues regarding pathologic downstaging have to be underlined: its value as a predictor of distant recurrence and disease-free survival. In our analysis we have found a direct effect of downstaging on these parameters that stress the importance of local control on lung cancer, generally considered a systemic disease, and the ability of surgical intervention to achieve an eradication of lung carcinoma.

This last consideration could directly affect the quality of life of patients according to the following syllogism: "no cancer, no treatment; no treatment, no side effect"—and no doctors, too.

Rolando M. D'Angelillo, MD
Lucio Trodella, MD
Sara Ramella, MD
Department of Radiotherapy
Università Cattolica S. Cuore
Rome, Italy

References

1. Machtay M, Lee JH, Stevenson JP, et al. Two commonly used neoadjuvant chemoradiotherapy regimens for locally advanced stage III non-small cell lung carcinoma: long-term results and associations with pathologic response. *J Thorac Cardiovasc Surg.* 2004; 127(1):108-13.
2. Eberhardt W, Wilke H, Stamatis G, et al. Preoperative chemotherapy followed by concurrent chemoradiation therapy based on hyperfractionated accelerated radiotherapy and definitive surgery in locally advanced non-small-cell lung cancer: mature results of a phase II trial. *J Clin Oncol.* 1998;16(2):622-34.
3. Albain KS, Rusch VW, Crowley JJ, et al. Concurrent cisplatin/etoposide plus chest radiotherapy followed by surgery for stages IIIA (N2) and IIIB non-small-cell lung cancer: mature results of Southwest Oncology Group phase II study 8805. *J Clin Oncol.* 1995;13(8):1880-92.
4. Emami B, Mirkovic N, Scott C, et al. The impact of regional nodal radiotherapy (dose/volume) on regional progression and survival in unresectable non-small cell lung cancer: an analysis of RTOG data. *Lung Cancer.* 2003; 41(2):207-14.
5. Trodella L, Granone P, Valente S, et al. neoadjuvant concurrent radiochemotherapy in locally advanced (IIIA-IIIB) non-small cell lung cancer: long term results according to downstaging. *Ann Oncol.* 2004;15:389-96.
6. Martin J, Ginsberg RJ, Abolhoda A, et al. Morbidity and mortality after neoadjuvant therapy for lung cancer: the risks of right pneumonectomy. *Ann Thorac Surg.* 2001; 72(4):1149-54.
7. Martin J, Ginsberg RJ, Venkatraman ES, et al. Long-term results of combined-modality therapy in resectable non-small-cell lung cancer. *J Clin Oncol.* 2002;20(8):1989-95.
8. Betticher DC, Hsu Schmitz SF, Totsch M, et al. Mediastinal lymph node clearance after docetaxel-cisplatin neoadjuvant chemotherapy is prognostic of survival in patients with stage IIIA pN2 non-small-cell lung cancer: a multicenter phase II trial. *Clin Oncol.* 2003; 21(9):1752-9.
doi:10.1016/j.jtcvs.2004.04.028

Studying the lumen in composite Y internal thoracic artery-saphenous vein grafts

To the Editor:

We congratulate Gaudino and colleagues¹ for the splendid idea advanced in their study on composite Y internal thoracic artery (ITA)-saphenous vein (SV) grafts. The study of the lumen contributes enormously to the current series of studies describing the characteristics of composite grafts. However, we do not share some of the opinions presented in this article. We believe that it is not right to use the term

"flow steal" in this specific case, because the circulatory system is pressurized, is closed, and responds well to demand. We do not agree that SV grafts are the worst conduits. It is already known that SVs in aorta-left anterior descending coronary artery grafts are patent in 70% to 80% of the patients after 10 years. We also believe that the reduced diameter in distal ITA used to revascularize coronary arteries that have a proximal moderate stenosis (<70%) is physiologic and expected. The distal ITA lumen may be reduced as well when the supplied coronary artery has a poor runoff. If there is no flow demand, the ITA graft will decrease in caliber, because ITA has a huge ability to adapt itself to flow demand. It is commonly seen that an ITA supplying coronaries with proximal severe stenosis and a good runoff is usually dilated.² The fact that SV is less reactive does not compromise the coronary irrigation to which it is anastomosed, because the blood flow in SV is proportional to the coronary runoff and not to its diameter. Nevertheless, we agree with current studies that demonstrate ITA's capacity to adapt to flow demand.³ We strongly believe that SV segments do not jeopardize the composite graft efficiency and have several advantages when used in such configuration: they are smaller, they suffer less pressure and shear stress, and they receive substances produced by ITA's endothelium (nitrous oxide, for example). These advantages are believed to increase SV graft durability.⁴⁻⁶ One must also question results from an experiment that did not use a group control; in addition, the small sample does not allow for definite conclusions, indicating the need for more research in this area.

José Glauco Lobo Filho, MD, MM, PhD
Maria Claudia de Azevedo Leitão, MD
Antonio Jorge de Vasconcelos Forte

ICORP
Clinical Research
Fortaleza, Brazil

References

1. Gaudino M, Alessandrini F, Pragliola C, Luciani N, Trani C, Burzotta F, et al. Composite Y internal thoracic artery-saphenous vein grafts: short-term angiographic results and vasoreactive profile. *J Thorac Cardiovasc Surg.* 2004;127:1139-44.
2. Ochi M, Hatori N, Bessho R, Fujii M, Saji Y, Tanaka S, et al. Adequacy of flow capacity of bilateral internal thoracic artery T graft. *Ann Thorac Surg.* 2001;72:2008-12.

3. Lobo Filho JG, Leitão MC, Lobo Filho HG, Silva AA, Machado JJ, Forte AJ, et al. Myocardial revascularization surgery using composite Y-graft of the left internal thoracic artery: blood flow analysis. *Rev Bras Cir Cardiovasc.* 2004;19:1-8.
4. Cox JL, Chiasson DA, Gotlieb AI. Stranger in a strange land: the pathogenesis of saphenous vein graft stenosis with emphasis on structural and functional differences between veins and arteries. *Prog Cardiovasc Dis.* 1991;34:45-68.
5. Nwasokwa ON. Coronary artery bypass graft disease. *Ann Intern Med.* 1995;123:528-45.
6. Tarr F, Dudas G, Tarr M, Racz R, Sasvari M, Tomcsanyi I. Analysis of the stable metabolite of endothelium-derived nitric oxide of internal mammary artery bypass grafts at the venous drainage system of the recipient coronary artery: morphologic implications and consequences. *Orv Hetil.* 2002;143:2549-52.

doi:10.1016/j.jtcvs.2004.05.012

Reply to the Editor:

We thank Filho and colleagues for their comments. Although the capacity of the internal thoracic artery (ITA) to adapt its caliber to the flow demand is well known, in our series of composite Y ITA-saphenous vein (SV) grafts the proportion of cases with clear graft malfunction (string or occlusion) that resulted in ischemia in the grafted territory was not negligible. Reversibility of the string sign has been anecdotally reported but cannot be accepted for sure, and in our series it was shown to be unlikely by the scintigraphic evidence of inducible ischemia. For these reasons the term “flow steal” and our word of caution regarding the adoption of these type of hybrid grafts seem substantiated.

The hypothesis that SV graft durability can be increased by the vasoactive sub-

stances produced in the proximal tract of the ITA is totally speculative and requires objective investigation before consideration. Moreover, the fact that the SV is usually anastomosed to the extreme proximal tract of the ITA minimizes the amount of ITA-produced substances to which the SV graft is exposed.

Finally, because this type of graft configuration is used at our institution only in complex technical cases as last resort, recruitment of a control group would have been ethically unacceptable.

Mario Gaudino, MD
Gianfederico Possati, MD
Department of Cardiac Surgery
Catholic University
Rome, Italy

doi:10.1016/j.jtcvs.2004.05.013