systematic review of studies comparing bilateral and single internal mammary arteries. *Lancet*. 2001;358:870-5.

5. Nataf P, Hadjiisky P, Bourbon A, Peuchmaud M, Leprince P, Regan M, et al. Morphometric and metabolic profile of the distal segment of the internal mammary artery. *Eur J Cardiothorac Surg*. 1996;10:965-70.

doi:10.1016/j.jtcvs.2004.06.025

Reply to the Editor:

We read with great interest the comments of Dr Pocar and colleagues about the article we recently published in *The Journal of Thoracic and Cardiovascular Surgery*.¹

It is mandatory for everyone involved in cardiac surgery to know everything about the reasons that have been mentioned in the first and second paragraphs of their comments. We have previously reported and underlined our policy regarding internal thoracic artery (ITA) harvesting [. . . In order to optimize the internal thoracic artery length, and avoid the use of the most distal part of internal thoracic artery (often small in diameter and prone to spasm), we try to make the internal thoracic artery graft as short as possible. . .].² According to this policy, it was not a very distal part of the ITA that was used for coronary-coronary bypass grafting (CCBG). A crossclamp time of 41 minutes to perform 4 anastomoses on coronary arteries is not too much, and for the current level of myocardial protection, 10 minutes more means nothing.

As to the third reason, let us suppose that ITA has been used to revascularize the distal third of the left anterior descending coronary artery (LAD), although it did not have enough length as a pedicled graft to reach the anastomotic site. An aggressive form of atherosclerotic disease would progress with time, no matter that the coronary artery was grafted with the ITA, because coronary artery disease progress even if the targeted artery was grafted with an arterial graft (although to a lesser extent than if it was grafted with vein graft).³ A new proximal lesion on the LAD great enough to jeopardize distal flow (through a coronary-coronary bypass graft) would probably be the reason to attempt the revascularization (percutaneous coronary intervention or redo coronary artery by-

pass grafting), even in the absence of a coronary-coronary graft on the LAD, as well as in the case that the ITA (eg, in skeletonized fashion) was used in the first operation.

I absolutely agree with Dr Pocar and associates that anastomoses should ideally be constructed to bypass the most distal lesion, but it is obvious that with CCBG over distal and single stenosis on the LAD, we have done exactly what they have underlined. Free ITA grafts yield excellent patency rates of approximately 90% at 5 years,⁴ as well as at 7 years.⁵ We know that a major problem influencing early free ITA graft patency was related to construction of a proximal anastomosis on the ascending aorta. In the case of CCBG, we have a situation that a segment of the free ITA graft at the site of proximal anastomosis matches better with the lower dp/dt of the LAD artery than the ascending aorta. That is the reason that we can expect an even better patency rate over the years. There have been only 2 cases of late angiography in patients with coronary-coronary free ITA grafts on the LAD (3 and 8 years after surgical intervention) recording patent coronary-coronary grafts.⁶

I agree with Dr Pocar and associates that CCBG might at times be useful, and that was in the reported case. We have performed more than 25,000 cardiac procedures at our institution over 25 years. I was the attending surgeon in more than 2500 cardiac procedures. Of more than 1600 procedures performed per year (70% being coronary artery operations), we have selected only 2 patients for CCBG of a single distal lesion of the large LAD (in the patients with multiple-vessel disease). The other patient's single distal LAD lesion was grafted with a short segment of the radial artery in CCBG fashion.⁷ It means that our candidates for CCBG procedure were selected very carefully.

As to the fourth reason, when you have to perform coronary surgery because of multivessel disease, it does not make sense to increase cost with implantation of a drug-eluting stent, even in the countries with great economic potential.

Finally, our intention was not to challenge the guidelines for coronary revascularization but to underline that once in a

blue moon we can meet the patient who does not fit in a guideline frame.

Duško Nežić, MD, PhD, FETCS
Department of Cardiac Surgery
Dedinje Cardiovascular Institute
Belgrade, Serbia and Montenegro

References

1. Nežić D, Knežević A, Borović S, Čirković M, Milojević P. Coronary-coronary free internal thoracic artery graft on a single, distal, left anterior descending artery lesion. *J Thorac Cardiovasc Surg*. 2004;127:1517-8.
2. Nežić D, Perić M, Knežević A, Čirković M, Jović M, Bojčić M. Tension-free left ITA graft—the pericardial strip technique. *Cardiovasc Surg*. 2001;9(2):177-8.
3. Manninen HI, Jaakkola P, Suhonen M, Rehnberg S, Vuorenniemi R, Matsi PJ. Angiographic predictors of graft patency and disease progression after coronary artery bypass grafting with arterial and venous grafts. *Ann Thorac Surg*. 1998;66:1289-94.
4. Woo YJ, Gardner TJ. Myocardial revascularization with cardiopulmonary bypass. In: Cohn LH, Edmunds LH Jr, editors. *Cardiac surgery in adult*. New York: McGraw-Hill; 2003. p. 581-607.
5. Tatoulis J, Buxton B, Fuller J. Patencies of 2127 arterial to coronary conduits over 15 years. *Ann Thorac Surg*. 2004;77:93-101.
6. Barbosa G, Rusticali F. Proximal internal mammary in situ graft and distal coronary-coronary graft. *Tex Heart Inst J*. 2000;27:70-1.
7. Nežić D, Čirković M, Knežević A, Mangovski LJ. Coronary-coronary radial artery graft for single, distal LAD lesion. *Ann Thorac Surg*. 2004. In press.

doi:10.1016/j.jtcvs.2004.06.026

Repair of the Ebstein anomaly To the Editor:

I read with great interest the article by Chen and colleagues¹ "Early and Medium-Term Results for Repair of Ebstein Anomaly." The results are superb, without hospital mortality and with improvement of the functional condition. The technique proved to be reproducible.

It seems from previous studies² that the Ebstein anomaly is a combination of ventricular and valvular disease. This concept has led my group to develop the use of an associated bidirectional cavopulmonary shunt in severe cases.³

I was surprised by the preoperative right ventricular ejection function, which was normal in all cases except 1 in Dr Chen's series, without quantified data.

My first concern is to know what is meant by the right ventricle. Is it the effec-

tive right ventricle below the attachment of the valve or the right ventricle with the atrialized right ventricle? In most of the patients on whom my own group has operated, the right anterior wall was dilated and hypokinetic.⁴ I am interested to know whether this aspect was present in Chen and colleagues' series.¹

The postoperative decrease of right ventricular ejection fraction could be due to the section of abnormal muscular trabeculations. It is in my mind an additional reason to decrease the preload of the right ventricle with a partial Glenn procedure.

Sylvain Chauvaud, MD
Department of Cardio-Vascular Surgery
Hôpital Européen Georges Pompidou
20 rue Leblanc
75015 Paris, France

References

1. Chen JM, Mosca RS, Altmann K, Printz BF, Targoff K, Mazzeo PA, et al. Early and medium-term results for repair of Ebstein anomaly. *J Thorac Cardiovasc Surg.* 2004;127:990-9.
2. Celermajer DS, Bull C, Till JA, Cullen S, Vassilikos V, Sullivan ID, et al. Ebstein's anomaly: presentation and outcome from fetus to adult. *J Am Coll Cardiol.* 1994;23:170-6.
3. Chauvaud S, Fuzellier JF, Berrebi A, Lajos P, Marino JP, Mihaileanu S, et al. Bi-directional cavopulmonary shunt associated with ventriculo and valvuloplasty in Ebstein's anomaly: benefits in high risk patients. *Eur J Cardiothorac Surg.* 1998;13:514-9.
4. Chauvaud S, Berrebi A, d'Attellis N, Mousseaux E, Hernigou A, Carpentier A. Ebstein's anomaly: repair based on functional analysis. *Eur J Cardiothorac Surg.* 2003;23:525-31.
doi:10.1016/j.jtcvs.2004.06.041

Reply to the Editor:

My coauthors and I are in agreement with Dr Chauvaud's impression that the Ebstein anomaly is a combination of ventricular and valvular disease. In our study, the right ventricle, as analyzed by transthoracic echocardiography, was considered to be the effective right ventricle below the attachment of the valve. In several patients—most notably the adults—the right anterior wall was indeed hypokinetic. Care certainly must be taken not to assign hypokinetic areas of atrialized ventricle to the true right ventricle. We have not routinely used a Glenn cavopulmonary shunt in our Ebstein repair, but we recognize it as a viable strategy in those patients for whom reduc-

tion in right-sided volume loading may be beneficial.

Jonathan M. Chen, MD
Pediatric Cardiac Surgery
Columbia University College of Physicians and Surgeons
New York, NY 10032
doi:10.1016/j.jtcvs.2004.06.042

Pleural effusion and off-pump Fontan procedure

To the Editor:

We read with great interest the study by Gupta and colleagues¹ "Risk Factors for Persistent Pleural Effusions After the Extracardiac Fontan Procedure," published in the June 2004 issue. Prolonged and excessive pleural drainage after the Fontan procedure is still the subject of debate. In their series of 100 patients, Gupta and colleagues¹ reported prolonged duration of pleural drainage in 37% and increased volume of pleural drainage in 30% after the extracardiac Fontan operation.¹ Lower preoperative oxygen saturation, presence of postoperative infection, smaller graft size, and longer cardiopulmonary bypass time were significantly associated with prolonged and increased pleural drainage.

We have used an off-pump technique for the bidirectional Glenn shunt and the extracardiac Fontan operation in patients without intracardiac anomalies.^{2,3} Prolonged pleural effusion was seen in 2 of 30 patients undergoing the bidirectional Glenn shunt (6.6%). Among 10 patients undergoing the off-pump extracardiac Fontan operation, only 1 (10%) had pleural drainage for longer than 2 weeks.

Lower preoperative oxygen saturation and increased preoperative pulmonary arterial pressure may be consequences of increased pulmonary vascular resistance. We surmise that cardiopulmonary bypass causes prolonged and excessive pleural effusion with increase of pulmonary vascular resistance and pulmonary arterial pressure. In patients with cyanosis, the blood flow of the major aortopulmonary collateral arteries may increase during cardiopulmonary bypass, which causes prolonged pleural effusion in some patients.

In conclusion, cardiopulmonary bypass is significantly associated with increased volume of pleural drainage after both the Glenn shunt and the extracardiac Fontan procedure. We believe that pleural drain-

age will decrease with the use of an off-pump technique.

[Response declined]

Ali Kubilay Korkut, MD
Gurkan Cetin, MD
Ilksen Soyler, MD
Emin Tireli, MD
Department of Cardiovascular Surgery
Istanbul University
Istanbul, Turkey

References

1. Gupta A, Daggett C, Behera S, Ferraro M, Wells W, Starnes V. Risk factors for persistent pleural effusions after the extracardiac Fontan procedure. *J Thorac Cardiovasc Surg.* 2004;127:1664-9.
2. Tireli E, Basaran M, Kafali E, Harmandar B, Camci E, Dayioglu E, et al. Perioperative comparison of different transient external shunt techniques in bidirectional cavo-pulmonary shunt. *Eur J Cardiothorac Surg.* 2003;23:518-24.
3. Tireli E. Extracardiac Fontan operation without cardiopulmonary bypass: how to perform the anastomosis between inferior vena cava and conduit. *Cardiovasc Surg.* 2003;11:225-7.

doi:10.1016/j.jtcvs.2004.06.044

The difference is meaningful: Anatomic coronary-coronary bypass or physiologic coronary-coronary bypass?

To the Editor:

It was with great interest that I read the communication of Nežić and colleagues.¹ The authors have renewed interest in coronary-coronary bypass (CCB) grafting by highlighting its complementary technical issues, especially in the setting of multiple left anterior descending artery (LAD) stenosis, which might require a synchronous multisided revascularization. The authors have advanced the presumed physiologic advantages related to CCB, aiming thereby to promote their current surgical alternative. However, the current case points out the vigilance that should be paid in distinguishing anatomic CCB from physiologic CCB.

The physiologic advantage of CCB has been highlighted by Biglioli and associates.² However, these authors have imputed this physiologic advantage directly to the diastolic coronary flow rather than to its systolic fraction. Biglioli and associates used the initial portion of the right coronary artery as a donor site for the saphenous graft proximal anastomosis providing the LAD. From a physiologic point of view, Biglioli and asso-