included in the investigation. For example, in the study of Fakler and associates,¹ new formulas with new coefficients could be calculated using 70 patients included in their investigation and could be validated against the remaining 73 patients for comparing measured and estimated VO_2 .

Fakler and associates¹ reported that the routine use of assumed VO_2 when calculating cardiac output with the direct Fick principle might frequently result in large errors in the estimation of cardiac output and dependent parameters. To obtain reliable cardiac output values by the direct Fick equation, VO_2 should be measured as concluded by Fakler and associates¹ or estimated with a new formula calculated using the population included in the study.

Habib Benallal

Experimental Toxicology

Department-Modeling Group

INERIS (National Institute for industrial

Environment and RISks)

60550 Verneuil en Halatte, France

References

1. Fakler U, Pauli C, Henning M, Sebening W, Hess J. Assumed oxygen consumption frequently results in large errors in the determination of cardiac output. *J Thorac Cardiovasc Surg.* 2005;130:272-6.
2. Krovetz LJ, Goldbloom S. Normal standards for cardiovascular data. I. Examination of the validity of cardiac index. *Johns Hopkins Med J.* 1972;130:174-86.
3. LaFarge CG, Miettinen OS. The estimation of oxygen consumption. *Cardiovas. Res.* 1970;4: 23-30.
4. Laitinen PO, Räsänen J. Measured versus predicted oxygen consumption in children with congenital heart disease. *Heart.* 1998; 80:601-5.

doi:10.1016/j.jtcvs.2005.08.069

Reply to the Editor:

We thank Benallal for his comments on our article and appreciate his remarks.

Our study was specifically designed to assess the importance of measured oxygen consumption (VO_2) in the preoperative diagnostic hemodynamic evaluation during cardiac catheterization and to assess whether published formulas allow a reliable calculation of VO_2 . We did not intend to develop a new formula to assume VO_2 .

Our results emphasize that both the Krovetz-Goldbloom formula,¹ which had been used to this point in our institution, and the more commonly used formula published by LaFarge and Miettinen² are not

feasible in all patients to assume VO_2 and thereby calculate cardiac output by the Fick principle.

Li and colleagues³ presented the important conclusion that other predictive equations published by Lundell and associates,⁴ Wessel and colleagues,⁵ and Lindahl⁶ also do not accurately estimate VO_2 . All these formulas were developed to offer a suitable and reliable method to assume VO_2 in order to simplify the calculation of hemodynamic parameters. However, these formulas cannot be used in all patients and settings.

As Benallal pointed out, these formulas include coefficients depending on characteristics of the population investigated in these studies. The difference between measured and assumed VO_2 might be due to a population difference. In addition, it is also influenced essentially by other conditions such as the mode of general anesthesia and relaxation used in different institutions.

It is difficult to identify feasible coefficients to develop a formula fitting to all conditions of hemodynamic evaluation during cardiac catheterization.

Moreover, we are convinced that our population of 143 patients is too small to identify and validate suitable coefficients for a new formula. LaFarge and Miettinen derived their most commonly used formulas from a multivariate analysis of covariance applied to data on a series of 879 patients.

U. Fakler, MD

J. Hess, MD

German Heart Center

Department of Pediatric Cardiology and

Congenital Heart Disease

Technische Universität München

Munich, Germany

References

1. Krovetz LJ, Goldbloom S. Normal standards for cardiovascular data. I. Examination of the validity of cardiac index. *Johns Hopkins Med J.* 1972;130:174-86.
2. LaFarge CG, Miettinen OS. The estimation of oxygen consumption. *Cardiovasc Res.* 1970;4: 23-30.
3. Li J, Bush A, Schulze-Neick I, Penny DJ, Redington AN, Shekerdemian LS. Measured versus estimated oxygen consumption in ventilated patients with congenital heart disease: the validity of predictive equations. *Crit Care Med.* 2003;31:1235-40.
4. Lundell BPW, Casas ML, Wallgren CG. Oxygen consumption in infants and children during heart catheterization. *Pediatr Cardiol.* 1996;17:207-13.

5. Wessel HU, Rorem D, Muster AJ, Acevedo RE, Paul MH. Continuous determination of oxygen uptake in sedated infants and children during cardiac catheterization. *Am J Cardiol* 1969;24:376-85.
6. Lindahl SGE. Oxygen consumption and carbon dioxide elimination in infants and children during anaesthesia and surgery. *Br J Anaesth.* 1989;62:70-6.

doi:10.1016/j.jtcvs.2005.11.005

Linking gene expression, nuclear factor kappa B, remote ischemic preconditioning, and transplantation: A quest for an elusive Holy Grail or a road to an amazing discovery?

To the Editor:

We read with interest the recent paper of Ishiyama and colleagues¹ reporting that inhibition of nuclear factor kappa B (NF- κ B) by I κ B suppressor gene transfer ameliorated ischemia-reperfusion (IR) injury after experimental lung transplantation. A few comments seem appropriate.

Is activation of NF- κ B before transplantation destructive or protective? Two murine studies provided seemingly contradictory results. In the study of Ishiyama and colleagues,¹ inhibition of NF- κ B rendered protection against the IR injury, yet in another study² a targeted deletion of the p105 subunit of the NF- κ B abolished the protective effect of the remote ischemic preconditioning (rIPC), a novel clinically applicable mode of protection against IR injury, whereby the organs can be protected by a brief ischemia applied *remotely* (eg, limb ischemia). Gene expression appears to play an important role in rIPC. In our own studies, the rIPC stimulus modified NF- κ B regulated gene expression in leukocytes³ and the target organ,⁴ and when the rIPC was applied to the recipient in porcine model it rendered myocardial protection *after* the orthotopic heart transplantation.⁵

It is widely known that the transcription factor NF- κ B can be activated by a variety of stimuli, for example, tumor necrosis factor (TNF)- α , bacterial lipopolysaccharide (LPS), and heat shock protein (HSP). Stimulation of the cell may lead to phosphorylation of I κ B and translocation of NF- κ B into the nucleus, resulting in gene expression, including those genes responsible for exocytosis of adhesive molecules and cytokine production in leukocytes, but also for production of protective substances eg, manganese superoxide dismutase (Mn-SOD) and inducible nitric oxide synthase

Figure 1. Innate immunity pathways and NF- κ B. Remote ischemic preconditioning stimulus upregulates (red) and downregulates (green) gene expression of intracellular inflammatory pathways. *HSP*, Heat shock protein; *LPS*, lipopolysaccharide; *LBP*, LPS binding protein; *TLR*, toll-like receptor; *IRAK*, interleukin (IL)-1 receptor associated kinase; *TIR*, toll/IL-1/plan R homology domain; *IKK*, I- κ B kinase; *MnSOD*, manganese superoxide dismutase. (From Konstantinov IE, Arab S, Kharbanda RK, Li J, Cheung MM, Cherepanov V, et al. The Remote Ischemic Preconditioning Stimulus Modifies Inflammatory Gene Expression in Humans. *Physiol Genomics*. 2004;19:143-50; used with permission of The American Physiological Society.)

(iNOS)² (Figure 1). It is perhaps less known, however, that the “classic” NF- κ B exists as a p50-p65 heteromer. The p50 derives from the precursor protein p105, which appears to be indispensable for rIPC.² Suppression of gene expression of NF- κ B p65 subunit by the rIPC stimulus observed in our study³ was of special interest because this might be a potential site of uncoupling of production of cytokines and MnSOD/iNOS synthesis (Figure 1). It appears that the initial TNF- α signaling pathway activation by rIPC induces protective MnSOD synthesis, but yet suppresses gene expression responsible for subsequent cytokine production and TNF- α signaling pathway restoration. Taking into consideration that general nonspecific inhibition of transcription by actinomycin-D abolishes

protection of ischemic preconditioning, it seems likely that a brief activation of NF- κ B might be protective, while its sustained activation may have detrimental results. Or, perhaps, a unit-specific activation of the NF- κ B is more important?

Further research of NF- κ B involvement in IR injury, particularly in transplantation, could be an area where the major discoveries in cardiothoracic surgery may be made.

Igor E. Konstantinov, MD, PhD

Andrew N. Redington, MD

Divisions of Cardiovascular Surgery and

Cardiology

Hospital for Sick Children

University of Toronto

555 University Ave

Toronto, M5G 1X8, Ontario, Canada

References

1. Ishiyama T, Dharmarajan S, Hayama M, Moriya H, Grappenhuis K, Patterson GA. Inhibition of nuclear factor κ B by I κ B superrepressor gene transfer ameliorates ischemia-reperfusion injury after experimental lung transplantation. *J Thorac Cardiovasc Surg*. 2005;130:194-201.
2. Li G, Labruto F, Sirsjo A, Chen F, Vaage J, Valen G. Myocardial protection by remote preconditioning: the role of nuclear factor kappa-B p105 and inducible nitric oxide synthase. *Eur J Cardiothorac Surg*. 2004;26:968-73.
3. Konstantinov IE, Arab S, Kharbanda RK, et al. The remote ischemic preconditioning stimulus modifies inflammatory gene expression in humans. *Physiol Genomics*. 2004;19:143-50.
4. Konstantinov IE, Arab S, Li J, Coles JG, Boscarino C, Mori A, et al. The remote ischemic preconditioning stimulus modifies

gene expression in mouse myocardium. *J Thorac Cardiovasc Surg.* 2005;130:1326-32.

5. Konstantinov IE, Li J, Cheung MM, Shimizu M, Stokoe J, Kharbanda RK, et al. Remote ischemic preconditioning of the recipient reduces myocardial ischemia-reperfusion injury of the denervated donor heart via a Katp channel-dependent mechanism. *Transplantation.* 2005;79:1691-5.

doi:10.1016/j.jtcvs.2005.07.043

From mesothelioma to cardiovascular protection via the phosphoinositide-3 kinase pathway: A new vista in cardiothoracic surgery

To the Editor:

We read with interest the article by Rascoe and colleagues¹ on the role of tyrosine kinase receptors and the phosphoinositide-3 kinase (PI3K) pathway in mesothelioma. The authors demonstrated differential involvement of 3 growth factor (GF) receptors—epidermal growth factor (EGF), insulin-like growth factor (IGF), and platelet-derived growth factor (PDGF) receptors—in PI3K-mediated survival of the malignant mesothelioma cells.

Survival of many cells under stress, particularly of myocytes, is dependent on PI3K pathway activation.² It is not often appreciated, however, that the PI3K pathway can be activated not only via tyrosine kinase receptors by GFs, but also via toll-like receptor by heat shock protein, bacterial lipopolysaccharide, and by tumor necrosis factor (Figure 1). Although the precise molecular mechanisms by which the GF receptors interact with PI3K pathway are unknown, they appear to have different properties. For example, PDGF appears to play a crucial role in the pathogenesis of atherosclerosis.³ We agree with Rascoe and colleagues that PDGF receptor inhibition does not negatively affect cellular survival. In fact, PDGF inhibition results in significant reduction in oxidative stress³ and is observed after ischemic preconditioning.⁴ Indeed, we have previously demonstrated that intravenous infusion of IGF improves oxygen transport by reducing oxygen consumption as well as increasing cardiac output and oxygen delivery after ischemia-reperfusion (IR) injury.⁵ Further-

more, cardiovascular protection from IR injury is abrogated when PI3K pathway is inhibited by LY294002.² Thus, systemic inhibition of IGF or PI3K may result in detrimental clinical side effects.

While local inhibition of PI3K signaling either directly or via EGF receptor inhibition may appear a promising treatment for mesothelioma, it is possible that sustained systemic inhibition of PI3K, either directly by LY294002 or indirectly via tyrosine kinase receptor inhibition, could lead to rapid progression of vascular disease and increased myocardial vulnerability to IR syndromes in susceptible patients. Whatever the clinical implications of the study by Rascoe and colleagues might be, any therapeutic intervention must be applied cautiously and balanced carefully against potential risks.

In summary, we congratulate Rascoe and colleagues on their stimulating and thought-provoking study. It is clear that contemporary academic cardiothoracic surgery must be performed within the realms of an understanding of the burgeoning field of molecular

Figure 1. Schematic of phosphoinositide-3 kinase (PI3K) signaling pathway. Activation of growth factor receptor results in phosphorylation (P) on tyrosine residues. PI3K is activated by binding to tyrosine residues of growth factor receptor with its regulatory subunit (p85). The lipid product of PI3K, phosphatidylinositol-3,4,5-triphosphate (PIP3), recruits serine-threonine kinase (Akt) via pleckstrin homology (PH) domain to the membrane, where it is activated. Akt phosphorylation of the apoptosis inducing protein BAD prevents BAD from binding to Bcl-2, thus releasing it for survival response. Direct binding to Ras protein further stimulates PI3K activity. Stimulation of toll-like receptor (TLR) 4 by bacterial lipopolysaccharide (LPS) or heat shock protein (HSP) results in PI3K activation via the myeloid differentiation factor (MyD) 88. Tumor necrosis factor (TNF) produces PI3K/Akt-mediated phosphorylation of p65 subunit of nuclear factor kappaB (NF-κB) via the alpha subunit of IkappaB kinase (IKK). Translocation of NF-κB to the nucleus results in gene expression and synthesis of cytokines, manganese superoxide dismutase (MnSOD), and endothelial nitric oxide synthase (eNOS) that are protective during acute ischemia.