this response. The rapid transactivation of EGFR occurs by an intracellular Src-mediated event, but not through the release of an extracellular epidermal growth factor-like ligand. These results suggest that the early effects of COX-2-derived PGE₂ are in part mediated by EGFR, and this transactivation is responsible for subsequent downstream effects, including the stimulation of cell migration and invasion.⁵ In the light of this information, we propose that celecoxib may inhibit downstream signaling of EGFR by decreasing the COX-2 protein and PGE₂ levels, resulting in tumor growth inhibition in NSCLC.

Mehmet Ates, MD^a
Ozden Altundag, MD^b
Serdal Aktolga, MD^c
Kadri Altundag, MD^d

Department of Cardiovascular Surgery
Siyami Ersek Thoracic and
Cardiovascular Education and Research
Hospital
Istanbul, Turkey^a
8181 Fannin St
No. 728
Houston, TX 77054^b
Department of Internal Medicine
Marmara University Medical School
Istanbul, Turkey^c
Department of Medical Oncology
Hacettepe University Medical School
Ankara, Turkey^d

References

1. Sievers EM, Bart RD, Backhus LM, Lin Y, Starnes M, Castanos R, et al. Evaluation of cyclooxygenase-2 inhibition in an ortho-

topic murine model of lung cancer for dose-dependent effect. *J Thorac Cardiovasc Surg.* 2005;129:1242-9.

2. Gupta RA, Tejada LV, Tong BJ, Das SK, Morrow JD, Dey SK, et al. Cyclooxygenase-1 is overexpressed and promotes angiogenic growth factor production in ovarian cancer. *Cancer Res.* 2003;63:906-11.
3. Mendelsohn J, Baselga J. Status of epidermal growth factor receptor antagonists in the biology and treatment of cancer. *J Clin Oncol.* 2003;21:2787-99.
4. Sheng H, Shao J, Washington MK, DuBois RN. Prostaglandin E₂ increases growth and motility of colorectal carcinoma cells. *J Biol Chem.* 2001;276:18075-81.
5. Buchanan FG, Wang D, Bargiacchi F, DuBois RN. Prostaglandin E₂ regulates cell migration via the intracellular activation of the epidermal growth factor receptor. *J Biol Chem.* 2003;278:35451-7.

doi:10.1016/j.jtcvs.2005.07.046

Reply to the Editor:

We thank Ates and colleagues for their thoughtful comments. We agree that there is an interrelationship between the cyclooxygenase-2 (COX-2) enzyme (and prostanoïd products) and the epidermal growth factor receptor (EGFR) family. In an important article, Pai and colleagues¹ reported that COX-2/prostaglandin E₂ transactivates the EGFR receptor, with resultant activation of downstream signaling.¹ Similarly it has been proposed that EGFR signaling can increase COX-2 activity. These findings have given cause for initiation of human trials of combination therapy with both EGFR receptor inhibitors and celecoxib in patients with non-small cell cancer, with some exciting initial results. However, the interaction between the two is complex and as yet incompletely understood.

Recently, Richardson and colleagues² found that COX-2 protein levels are independent of EGFR expression or activation in patients with non-small cell cancer.² We have found that combination therapy (EGFR receptor blocker together with COX-2 blockade by celecoxib) are occasionally antagonistic (at least in one cell line) in an animal orthotopic lung cancer model. When understanding the interaction, though, it is important to realize that levels of the COX-2 enzyme (both messenger RNA and protein) are variably affected by celecoxib, but the activity of the enzyme is profoundly blocked even by low doses of the drug. With regard to final sentence of the letter, we would agree that celecoxib decreases PGE₂ levels (but not COX-2 levels) and that interactions with the EGFR cascade probably account for some of the antineoplastic activity of the compound.

Eric Sievers, MD
R. M. Bremner, MD, PhD
Department of Cardiothoracic Surgery
Keck School of Medicine
Los Angeles, CA 90033

References

1. Pai R, Soreghan B, Szabo IL, Pavelka M, Baatar D, Tarnawski AS. Prostaglandin E₂ transactivates EGF receptor: a novel mechanism for promoting colon cancer growth and gastrointestinal hypertrophy. *Nat Med.* 2002; 8:289-93.
2. Richardson CM, Richardson D, Swinson DE, Swain WA, Cox G, O'Byrne KJ. Cyclooxygenase-2 protein levels are independent of epidermal growth factor receptor expression or activation in operable non-small cell lung cancer. *Lung Cancer.* 2005;48:47-57.
doi:10.1016/j.jtcvs.2005.08.002