The options available to treat pediatric airway obstruction include surgical resection and anastomosis (including slide tracheoplasty), silicone (eg, Dumon) stents, Montgomery T-tubes, and SEMS. Recently, there has been a disturbing trend of using SEMS as the first option in treating benign tracheobronchial pathology, primarily because of the ease of deployment and the dramatic relief of obstruction in the short term.² We feel that it needs to be reiterated strongly that SEMS should remain the last resort in patients with benign stenosis or those with a surgically correctable airway obstruction. SEMS have a number of complications that are difficult to treat long-term. The ideal indication for SEMS is a patient with terminal malignancy with airway obstruction in whom the expected low survival from the primary malignancy precludes the occurrence of stent-related complications. SEMS cause intense granulation tissue, which often causes the same symptoms for which the stent was deployed; subsequent attempts to alleviate this problem with laser, electrocautery, and so on usually have disappointing results. Moreover, contrary to the authors' views, removal of these stents is difficult and fraught with danger. Single and short-segment tracheal stenosis should be treated by primary surgery in centers experienced in pediatric tracheal surgery. Stenting should be reserved for patients in an acute situation (where silicone stents may be deployed as a temporary measure) before referral to a center specializing in tracheal surgery.

Although not practical in infants and very small children, Dumon stents and Montgomery T-tubes are invaluable in stenting lesions that are too long for safe reconstruction.³ More often than not, the stent stabilizes the airway, and decannulation is possible in most cases.⁴ Moreover, the ease with which silicon stents can be removed if definitive surgery is being considered makes it an attractive option to tide over an acute situation. The pediatrician, pulmonologist, or thoracic surgeon involved in managing complex airway problems needs to be aware of the basic underlying pathology and intelligently apply the right solution to a specific problem.

C. S. Pramesh, MS, FRCS
Rajesh C. Mistry, MS
Division of Thoracic Surgery
Tata Memorial Hospital
Mumbai, India

References

1. Vinograd I, Keidar S, Weinberg M, Silbiger A. Treatment of airway obstruction by metallic stents in infants and children. *J Thorac Cardiovasc Surg.* 2005;130:146-50.
2. Wood DE, Liu YH, Vallieres E, Karmy-Jones R, Mulligan MS. Airway stenting for malignant and benign tracheobronchial stenosis. *Ann Thorac Surg.* 2003;76:167-74.
3. Grillo HC. Stents and sense. *Ann Thorac Surg.* 2000;70:1142.
4. Pereszlenyi A, Igaz M, Majer I, Harustiak S. Role of endotracheal stenting in tracheal reconstruction surgery—retrospective analysis. *Eur J Cardiothorac Surg.* 2004;25:1059-64. doi:10.1016/j.jtcvs.2005.08.025

Reply to the Editor:

In my article I emphasized that the use of intratracheal stents in children is dangerous and carries the risk of severe complications.

With regard to the use of self-expandable metallic stents (SEMS), I completely agree that their use should be avoided. There are enough data in the literature to support this.¹ The SEMS in this study were made from nitinol, an alloy with a "shape memory effect." In animal studies, this stent, because of its unique biochemical property, could be fixed and removed easily, even in very small airways.² In children, their use was associated with an excessive granulation, and although the extraction proved to be easy and smooth, their application is currently suspended.

Currently available silicone or Dumon stents have an external diameter that prevents their use in infants and small children.

The Montgomery and I-tubes could be an option in tracheal strictures, but they are not practical in bronchial or distal airway problems.

I would like to thank Mr Pramesh for his remarks.

Itzhak Vinograd, MD
Department of Pediatric Surgery
DANA Children's Hospital
Sackler Faculty of Medicine
Tel Aviv University
Tel Aviv, Israel

References

1. Giaissert H, Grillo H, Wright C, Donahue D, Wain J, Mathisen D. Complication of benign tracheobronchial strictures by self-expanding metal stents. *J Thorac Cardiovasc Surg.* 2003;126:744-7.
2. Vinograd I, Klin B, Brosh T, Weinberg M, Flomenblit Y, Nevo Z. A new intratracheal stent made from nitinol, an alloy with "shape

memory effect." *J Thorac Cardiovasc Surg.* 1994;107:1255-61.

doi:10.1016/j.jtcvs.2005.08.040

Molecular changes occurring with aneurysm formation: Possible impact of sampling error

To the Editor:

We congratulate Cotrufo and colleagues¹ for exploring messenger RNA expression and extracellular matrix protein content in the ascending aorta associated with bicuspid aortic valves (BAVs), with particular attention to differences between the inner and outer curvatures. Our laboratory is also exploring the molecular changes occurring with aneurysm formation, and we have been concerned about the potential impact of sampling error on experimental results.

Cotrufo and colleagues¹ demonstrated differences in levels of collagens I, III, and IV, as well as fibronectin and laminin protein, by western blot analysis when comparing samples taken from the concave and convex portions of dilated aortas resected from patients with BAV. Additionally, they reported differences between patients with functionally regurgitant versus stenotic valves. Importantly, Cotrufo and colleagues¹ also reported significant differences in protein levels between dilated aortas (regardless of functional valve pathology) and normal control aortas. Indeed, these differences were as striking as those observed between concavity and convexity or between stenotic and regurgitant specimens.

We too are intrigued by the observations made on aneurysmal tissues, but we urge caution in interpreting the comparisons made between dilated aortas and the control group. A marked difference in age may have profoundly impacted results. Cotrufo and colleagues¹ reported mean ages for the regurgitation, stenosis, and control groups as 56 ± 9 , 59 ± 12 , and 33 ± 8 years, respectively. The 95% confidence intervals for these groups (from data provided in the article's on-line Table E1¹) were 51.2-61.4, 50.9-66.4, and 24.6-41.4 years, respectively. The difference in collagen expression led Cotrufo and colleagues¹ to conclude that "collagen degradation by proteinases could prevail over collagen synthesis" and that smooth muscle cells "could react differently to different types and degrees of mechanical stress and could favor, in turn, a predominant production of either ECM [extracellular matrix] proteins or proteolytic enzymes." An alternative hypothesis

is that age-related changes explain these differences. A proportional decrease in aortic collagen with advancing age has been previously described.² In our laboratory, we have looked at messenger RNA expression in dilated aortas with microarray gene chip analysis. Looking at expression of tenascin, for example, we have found similar expressions in tissues from patients with aneurysms and older control subjects (mean ages 63.3 ± 6.4 years, 95% confidence interval 59.8-66.9 years, and 48.8 ± 4.9 years, 95% confidence interval 35.3-62.3 years, respectively). We too have struggled to obtain appropriately age-matched control tissue for obvious reasons; however, appreciation of these age-related changes may be crucial to understanding the pathophysiology and genetic basis of BAV and aortic aneurysms and should not be overlooked.

Again, Cotrufo and colleagues are to be commended for their comparative study of expression of extracellular matrix proteins within dilated aortas among patients with BAV. Their comparisons between regurgitant and stenotic BAV and between the concavity and convexity of dilated aortas are valuable contributions to the field and should be in the foreground of discussions, whereas the comparisons between patients with disease and control subjects should remain in the background.

Stephen H. McKellar, MD

Sami Takieddine, MD

Ramanath Majumdar, PhD

Thoralf M. Sundt III, MD

Division of Cardiovascular Surgery

Mayo Clinic

200 First Street SW

Rochester, MN 55905

References

1. Cotrufo M, Della Corte A, De Santo LS, Quarto C, De Feo M, Romano G, et al. Different patterns of extracellular matrix protein expression in the convexity and concavity of the dilated aorta with bicuspid aortic valve: preliminary results. *J Thorac Cardiovasc Surg.* 2005;130:504-11.
2. Cattell MA, Anderson JC, Hasleton PS. Age-related changes in amounts and concentrations of collagen and elastin in normotensive human thoracic aorta. *Clin Chim Acta.* 1996; 245:73-84.

doi:10.1016/j.jtcvs.2005.09.014

Reply to the Editor:

We appreciate the comments made by McKellar and colleagues regarding our study on extracellular matrix (ECM) proteins in the dilated ascending aorta of patients with bicuspid aortic valve (BAV) disease.¹ We agree with their words of caution about the comparison between patients and control subjects unmatched for age. Unfortunately, the only setting in which we could obtain normal aortic tissue was multiorgan harvesting, and heart donors are usually younger than patients with aortic aneurysms.

Scarce information is available concerning age-related changes in aortic ECM composition in healthy subjects: in rats, the collagen amount per square millimeter decreases with age,² and similarly in human aortas, according to the study of Cattell and associates³ cited by McKellar and colleagues,³ the collagen amount per sample (in this case the amount in a tissue surface area 1 cm in diameter) decreased by 80% between the ages of 14 and 90 years. However, both in rats³ and in human beings,⁴ the concentration of collagen—that is, the amount per unit of tissue mass—increases with age, because of a predominant loss of other media constituents. In our study, as acknowledged in the Methods section,¹ the presence of collagen in both normal and diseased aortas was not quantified as amount per sample, which could have been affected by the differences in specimen thickness between diseased (thinner wall) and normal vessel, but in terms of the amount of collagen separated from a constant mass of total proteins. Therefore the difference in collagen between patients with BAV and control subjects could have been rather reduced than emphasized by a possible interference of age. Nevertheless, the actual impact of age differences cannot be discerned, and we are trying in our ongoing research to obtain study groups as properly matched for age as possible.

Indeed, our suggestion that the synthesis and degradation of each ECM protein may be differently regulated by different types and degrees of rheologic and mechanical disturbances¹ was derived from the comparison between patients with stenotic and regurgitant BAVs (with

comparable mean ages) and not that between patients and control subjects. Our most important finding, as recognized by McKellar and colleagues, was the different expressions of ECM proteins in the convexity and the concavity of dilated aortas, with typical signs of the “flow-induced vascular remodeling” process prevailing at the convexity. If further confirmed, this evidence could exclude that a congenital defect of some wall component may be the prime mover for aortic dilation in BAV disease, because a genetic disorder capable in itself of causing dilatation should be equally expressed along the circumference of a given tract of the vessel. On the contrary, wall stress, which is known to promote vascular remodeling,¹ has been found to unevenly involve the vessel when the valve is bicuspid, with maximal burden at the convex aspect of the ascending tract.⁵

Alessandro Della Corte, MD

Stefania Montagnani, MD

Maurizio Cotrufo, MD

Department of Cardiothoracic and Respiratory Sciences

Second University of Naples

V Monaldi Hospital

Via L Bianchi, 80131 Naples

Italy

References

1. Cotrufo M, Della Corte A, De Santo LS, Quarto C, De Feo M, Romano G, et al. Different patterns of extracellular matrix protein expression in the convexity and concavity of the dilated aorta with bicuspid aortic valve: preliminary results. *J Thorac Cardiovasc Surg.* 2005;130:504-11.
2. Bruel A, Oxlund H. Changes in biomechanical properties, composition of collagen and elastin, and advanced glycation endproducts of the rat aorta in relation to age. *Atherosclerosis.* 1996;127:155-65.
3. Cattell MA, Anderson JC, Hasleton PS. Age-related changes in amounts and concentrations of collagen and elastin in normotensive human thoracic aorta. *Clin Chim Acta.* 1996; 245:73-84.
4. Marque V, Kieffer P, Atkinson J, Lartaud-Idjouadiene I. Elastic properties and composition of the aortic wall in old spontaneously hypertensive rats. *Hypertension.* 1999;34: 415-22.
5. Robicsek F, Thubrikar MJ, Cook JW, Fowler B. The congenital bicuspid aortic valve: how does it function? Why does it fail? *Ann Thorac Surg.* 2004;77:177-85.

doi:10.1016/j.jtcvs.2005.09.034