tories of Dr Elefteriades and his colleagues.¹ Their work continues to broaden our fundamental understanding of this lethal but treatable disease. The recently reported work demonstrates a remarkable correlation between the predicted behavior of the aorta based on mechanical properties and the natural history of this disorder as observed previously by this same group of investigators. In the course of their discussion of the significance of their work, the authors graciously referenced our previous work touching on the mechanical properties of the ascending aorta, with particular emphasis on bicuspid aortic valve disease and changes with aging.² An additional finding from our study relevant to the author's work, however, perhaps should be made. As one might have anticipated, we observed changes in the intrinsic mechanical properties of the aorta, as defined by the elastic properties determined by means of biaxial testing. Of greater interest, perhaps, were our observed changes with age in the opening angle as a reflection of residual strain.

Residual stress and strain are critical concepts in understanding the mechanical properties of the arterial wall. Absent residual strain, intraluminal pressure within a thick-walled tube will result in an uneven distribution of stress across the wall, with the greatest stress born by the inner portion and the least stress by the outer. The pioneering work by Chuong and Fung³ demonstrated over 2 decades ago that the distribution of stresses and strains within a vessel wall are nonuniform, both circumferentially and longitudinally. Calculations based on the assumption that the unloaded vessel is at zero stress result in remarkably high stress concentrations at the inner wall. The unloaded vessel, however, is not physiologic. The vessel wall exhibits residual strain to effect a more even distribution of stress across the thickness of media and adventitia in the loaded state. Just as a concrete beam can be prestressed in anticipation of the loaded state, the effect of residual strain in a vessel wall is to make more uniform the distribution of circumferential wall stress under physiologic conditions with intraluminal pressure. Experimental observations confirm this phenomenon in vivo. Variations in residual strain have been observed in differing segments of the aorta in the rat, just as one would anticipate given differences in pressure wave forms.⁴ Liu

and Fung⁵ further demonstrated that residual strain is subject to remarkably rapid changes with cellular remodeling caused by altered hemodynamics.

The existence of residual strain is easily demonstrated in the operating room by using a test that is, in fact, familiar to all cardiovascular surgeons. If one resects a tubular segment of a vessel and then cuts the resulting ring open, the vessel will spring open (the opening angle) to a zero-stress state.⁶ This reflects residual tensile stress in the outer wall, while the inner wall has residual compressive stress. Accordingly, the ring will tend to turn itself inside out.

Of what possible clinical relevance is this phenomenon? Aneurysm rupture and dissection represent structural failures of the aortic wall. We also know that, despite arguments over the relative contributions of cellular apoptosis and fiber degradation, all would agree that rather dramatic changes in the structure of the wall itself can be observed with aging or against the backdrop of connective tissue disease. One can well imagine that concomitant changes in residual strain and other material properties might occur, as well as changes in vessel diameter and thickness in response to aging and hemodynamic forces. Could these alterations affect stress distribution across the wall of the aorta? It is tempting to hypothesize that alterations in vessel wall structure leading to disproportionate loading of the outer portion of the aortic wall could lead to primary failure of the subadventitia. Could this be an alternative explanation for the phenomenon of intramural hematoma? Could primary mechanical failure of the adventitia lead to creation of a potential space that fills with blood as a secondary phenomenon?

Although this theory is unsubstantiated by experimental work, it demonstrates the potential importance of accounting for residual strain among the mechanical properties of the aortic wall. In this regard we must respectfully disagree with the authors' statement that "a full profile of the mechanical properties of the aorta can be gleaned by measuring six specific physical characteristics: blood pressure (systolic and diastolic), aortic diameter (systolic and diastolic), and thickness of the aortic wall (systolic and diastolic)." We believe there is more to it than that. The intrinsic properties of this highly complex amalgam of

connective tissue and cellular elements must enter into the equation.

Thoralf M. Sundt, III, MD
Division of Cardiovascular Surgery
Mayo Clinic and Foundation
200 First St, SW
Rochester, MN 55905
E-mail: sundt.thoralf@mayo.edu

References

1. Koullias G, Modak R, Tranquilli M, Korkolis DP, Barash P, Elefteriades JA. Mechanical deterioration underlies malignant behavior of aneurysmal human ascending aorta. *J Thorac Cardiovasc Surg.* 2005;130:677-83.
2. Okamoto ref Okamoto RJ, Xu HD, Kouchoukos NT, Moon MR, Sundt TM. The influence of mechanical properties on wall stress and distensibility of the dilated ascending aorta. *J Thorac Cardiovasc Surg.* 2003;126:842-50.
3. Chuong CJ, Fung YC. Three-dimensional stress distribution in arteries. *J Biomech Eng.* 1983;105:268-74.
4. Liu SQ, Fung YC. Zero-stress states of arteries. *J Biomed Eng.* 1988;110:82-4.
5. Liu SQ, Fung YC. Relationship between hypertension, hypertrophy, and opening angle of zero-stress state of arteries following aortic constriction. *J Biomed Eng.* 1989;111:325-35.
6. Chuong CJ, Fung YC. On residual stress in arteries. *J Biomed Eng.* 1986;108:189-92. doi:10.1016/j.jtcvs.2005.12.070

Reply to the Editor:

Our team wishes to thank Dr Sundt for his extremely kind comments regarding our scientific investigations at the Center for Thoracic Aortic Disease at Yale University. We agree entirely with Dr Sundt that "residual strain" represents yet another facet of the complex mechanical behavior and measurable properties of the aortic wall. He and his colleagues, in the important work they cite in their letter, have measured this phenomenon in human aortas. From an engineering standpoint, residual strain is considered more important in thick-walled tubes (ratio of wall thickness to vessel diameter >1:10). The aneurysmal aorta is often thin-walled (thickness of 2-3 mm in a vessel with a diameter $\geq$ 5 cm) and often becomes even more so as the aneurysmal process advances (Figure 1). Thus the effect of residual strain, although an important inherent design characteristic of the human aorta, might diminish as the aneurysmal state advances.

We also have observed regional differences in stress distribution and resultant

Figure 1. The opened wall of a large ascending aortic aneurysm. Note that the wall of the aorta has become so thin that the markings on a ruler can be read right through the aortic wall.

strain. In our echocardiographic studies on normal and aneurysmal ascending aortas, we have consistently found that the anterior wall of the aorta (the “greater curve,” so to speak) expands more than the posterior wall. These data were recently presented at the American Society of Anesthesiologists meeting in Las Vegas (October 23-27, 2004). This finding agrees with the mechanical concept that an unsupported tissue layer (the anterior wall) experiences more stress compared with a supported layer (the posterior wall, which is supported by the pulmonary artery and the associated central mediastinal fibrous tissues). These findings validate and support the importance of regional variations in mechanical properties of the aorta that Dr Sundt and colleagues have detected.

We share Dr Sundt’s skepticism regarding ruptured vasa vasorum as the cause of intramural hematoma. In fact, prompted by an insightful observation by Dr Kouchoukos some years ago, we have scrutinized the internal lining of the widely opened aorta during deep hypothermic arrest in cases of intramural hematoma. In most, but not all, cases one can find an intimal tear that was not appreciated even on high-quality preoperative radiographic images. In a small minority of cases, no intimal tear exists. It is specifically to explain such no-tear intramural hematomas that the concept of rupture of the vasa vasorum has been postulated. Also, our

echocardiographic measurements have confirmed Dr Sundt’s hypothesis that stress on the adventitia is greater than stress on the media. We agree with Dr Sundt that uneven distribution of wall stress across the layers of the aortic wall might play a role in the genesis of intramural hematoma. This thesis awaits scientific confirmation.

George Koullias, MD

Raj Modak, MD

Paul Barash, MD

John A. Eleftheriades, MD

*Section of Cardiovascular Surgery
and Department of Anesthesia*

Yale University School of Medicine

333 Cedar St (121 FMB)

New Haven, CT 06510

E-mail: john.eleftheriades@yale.edu

doi:10.1016/j.jtcvs.2006.01.050

Correlation of autologous skeletal myoblast survival with changes in left ventricular remodeling in dilated ischemic heart failure

To the Editor:

McConnell and colleagues¹ present a well-designed study with implantation of skeletal myoblasts into an infarcted area. The infarction was created by interventional closing of the circumflex artery, and this resulted in dilated ischemic heart failure. The results, however, were somewhat disappointing. As assumed in the editorial by

Tang,² the small number of surviving myoblasts could be one reason for this outcome. Of course, there might be several reasons for cell loss of transplanted cells, but one important issue has not been addressed by both authors: ischemia. The cardiomyocytes of the target area were lost because of insufficient blood supply. There is no reason why the transplanted cells should be more tolerant to ischemia than the native myocardial cells! Therefore the restoration of blood supply to an infarcted area is possibly one important measure to improve survival of transplanted cells. Additionally, further cell division and differentiation also depend on sufficient oxygen and nutrition supply.

Helmut Gulbins, MD

Department of Cardiac Surgery

University Hospital Ulm

Steinhoevelstr. 9

Ulm D-89070

Germany

References

1. McConnell PI, del Rio CL, Jacoby DB, Pavlicova M, Kwiatkowski P, Zawadzka A, et al. Correlation of autologous skeletal myoblast survival with changes in left ventricular remodeling in dilated ischemic heart failure. *J Thorac Cardiovasc Surg.* 2005;130:1001.
2. Tang GH, Fazel S, Weisel RD, Verma S, Li RK. Optimizing cardiac cell therapy: from processing to delivery. *J Thorac Cardiovasc Surg.* 2005;130:966-8.
doi:10.1016/j.jtcvs.2005.11.050

Intercostal muscle flap without increase of pain and blood loss after lung surgery

To the Editor:

We read with interest the article by Cerfolio and colleagues.¹ This report demonstrated that harvesting the intercostal muscle flap (ICM) reduces the pain of thoracotomy. We also have some data about the pain in comparison of cases with ICM harvesting and other cases without ICM harvesting (non-ICM), as in the study performed in our institution. ICM is very mobile and vascular rich. Therefore the ICM was used in the reinforcement of perforated esophagus or bronchopleural fistula.² In our series of patients, performing lobectomy with reinforcement of ICM to bronchial stump never resulted in bronchopleural fistula over the decade.

We had 35 patients who had ICM flap with lobectomy (ICM group) and 51 pa-