Figure 2. Intraoperative transesophageal echocardiogram still frame illustrating a well-deployed low-profile device (arrow). There is a mild (1+) residual paravalvular leak. LA, Left atrium; LV, left ventricle; PVL, paravalvular leak.

occur as a result of gradual weakening of annular tissue, deterioration of suture material, technical mistakes, or acute infection.¹ Surgical intervention to repair the leak improves symptoms of heart failure, decreases the need for blood transfusion, and is an independent predictor of long-term survival when compared with medical therapy alone.^{2,5} The choice of operation involves either direct suture repair of the PVL or rereplacement of the valve and is indicated to correct worsening left ventricular function or symptoms of heart failure and hemolysis.¹

Although surgical intervention has been the gold standard for the repair of PVLs, there has been a growing interest in the attempt to close PVLs through the use of interventional cardiologic techniques. Several reports have documented the successful percutaneous repair of PVLs with closure devices similar to those applied

for the percutaneous treatment of septal defects.³ Percutaneous closure has several advantages, including the elimination of the risk of a redo operation and shortening the length of hospital stay. Nevertheless, the procedure is complex, time consuming, and not always successful, as illustrated in this case. Successful intraoperative device closure of muscular ventricular septal defects has previously been reported.⁶ However, to our knowledge, the case presented herein is the first reported use of the intraoperative placement of a percutaneous occluder device for the surgical repair of a PVL. This novel approach for the repair of a mitral PVL can be applied in the context of extremely difficult mitral prosthesis exposure, such as the presence of a concurrent aortic prosthesis, or to avoid lengthy redo operations in very high-risk patients.

References

1. Kirali K, Mansuroglu D, Yilmaci B, Omeroglu SN, Basaran Y, Ipek G, et al. Paravalvular leakage after mitral valve replacement: is left atrial enlargement an additional indication for reoperation? *J Heart Valve Dis.* 2001;10:418-25.
2. Akins CW, Bitondo JM, Hilgenberg AD, Vlahakes GJ, Madsen JC, MacGillivray TE. Early and late results of the surgical correction of cardiac prosthetic paravalvular leaks. *J Heart Valve Dis.* 2005;14:792-800.
3. Pate GE, Thompson CR, Munt BI, Webb JG. Techniques for percutaneous closure of prosthetic paravalvular leaks. *Catheter Cardiovasc Interv.* 2006;67:158-66.
4. Genoni M, Franzen D, Tavakoli R, Seiffert B, Graves K, Jenni R, et al. Does the morphology of mitral paravalvular leaks influence symptoms and hemolysis? *J Heart Valve Dis.* 2001;10:426-30.
5. Genoni M, Franzen D, Vogt P, Seifert B, Jenni R, Kunzli A, et al. Paravalvular leakage after mitral valve replacement: improved long-term survival with aggressive surgery? *Eur J Cardiothorac Surg.* 2000;17:14-9.
6. Bacha EA, Cao QL, Starr JP, Waight D, Ebeid MR, Hijazi ZM. Percutaneous device closure of muscular ventricular septal defects on the beating heart: technique and results. *J Thorac Cardiovasc Surg.* 2003;126:1718-23.

Feasibility of preservation of subvalvular apparatus in mitral valve replacement with the On-X mechanical valve

Hiroshi Tanaka, MD, Teruo Yamashita, MD, Kenji Okada, MD, Keitaro Nakagiri, MD, Yujiro Kawanishi, MD, and Yutaka Okita, MD, Kobe, Japan

Preserving subvalvular apparatus in mitral valve replacement brings a significant long-term advantage by preserving left ventricular function. The possibility of prosthetic valve entrapment of the retained subvalvular apparatus, left

ventricular outflow tract obstruction, and implantation of a smaller-sized prosthesis has been of major concern to investigators advocating this procedure. Recently introduced, the On-X mechanical valve (Medical Carbon Research Institute, Austin, Tex) has the advantage of reduced anticoagulation and a specific structure of leaflet flare for protection from retained subvalvular apparatus and pannus encroachment. We operated on 6 patients requiring mitral valve replacements using the On-X mechanical mitral valve and obtained good results without complications associated with this procedure.

Clinical Summary

From October 2005 to March 2006, 6 patients underwent mitral valve replacement for mitral valve stenosis using the On-X mitral valves with the techniques of preserving mitral papillary muscle continuity.

From Kobe University Hospital, Department of Cardiovascular, Thoracic, and Pediatric Surgery, Kobe, Japan.

Received for publication July 15, 2006; accepted for publication Aug 8, 2006.

Address for reprints: Hiroshi Tanaka, MD, Kobe University Hospital, Department of Cardiovascular, Thoracic, and Pediatric Surgery, 7-5-2, Kusunoki-cho, Chuo-ku, Kobe, Japan (E-mail: hirot@ams.odn.ne.jp).

J Thorac Cardiovasc Surg 2006;132:1470-1

0022-5223/\$32.00

Copyright © 2006 by The American Association for Thoracic Surgery

doi:10.1016/j.jtcvs.2006.08.030

TABLE 1. Techniques for preserving subvalvular apparatus and pre- and postoperative electrocardiography data

Case	Technique for preserving mitral-papillary muscle continuity	Preop		Postop		
		LVdD/Ds (mm)	EF (%)	LVdD/Ds (mm)	EF (%)	LVOT PG (mm Hg)
1	Artificial chordae ×4	43/30	70	42/36	64	5.4
2	PML preserved	38/23	64	38/28	71	6.7
3	Artificial chordae ×4	49/37	40	48/36	50	11.4
4	PML and lateral part of AML preserved; artificial chordae ×1	53/35	48	55/35	50	16.6
5	Artificial chordae ×4	47/28	64	48/25	63	12.9
6	PML preserved; artificial chordae ×2	53/40	42	55/40	45	4.8

LVdD, Left ventricular dimension diastole; Ds, dimension systole; EF, ejection fraction; LVOT PG, pressure gradient across the left ventricular outflow tract; PML, posterior mitral leaflet; AML, anterior mitral leaflet.

Figure 1. Comparing the profiles of prosthetic valves. a, Stentless bioprosthesis; b, Carbomedics standard valve; c, St Jude Medical Standard valve; d, the On-X valve.

Methods

After cardiopulmonary bypass is established, the aorta is clamped and cardioplegic arrest is obtained. The left atrium is opened from the right side after dissecting the groove. A Cosgrove retractor is inserted, and the mitral valve is examined. Basically, autologous subvalvular apparatus is meant to be preserved. However, in case of severe calcified subvalvular apparatus that cannot be preserved, we reconstruct the chordae tendineae with polytetrafluoroethylene sutures (Gore-Tex suture; W. L. Gore & Associates, Inc, Flagstaff, Ariz) as artificial chordae. If the patient's left ventricle is so small that the retained subvalvular apparatus may cause left ventricular outflow obstruction, preserving the continuity on the anterior mitral leaflet side is abandoned. The On-X valve is implanted with 2-0 Ethibond everting mattress sutures (Ethicon, Inc, Somerville, NJ) in the antianatomic position.

Results

No patients died in the hospital or during the follow-up period. Echocardiography was performed 1 month after the operation. The preservation techniques and preoperative and postoperative echocardiographic data are shown in Table 1. Left ventricular function was maintained or improved in all cases. No instances of prosthetic valve dysfunction and no significant pressure gradients across the left ventricular outflow tract were recorded.

Comments

Chordal preservation in mitral valve replacement has been widely accepted because of the advantage of preserving left ventricular function and decreasing the risk for ventricular rupture in mitral valve replacement.¹ This technique should incorporate several principles²: (1) The procedure must not cause left ventricular outflow obstruction. (2) Retained subvalvular structures should not interfere with the prosthetic valve mechanism. (3) Sufficient tissue should be resected to allow for implantation of a suitably sized valve.

Currently, various kinds of prosthetic valves are available in mitral valve surgery. The St Jude Medical mitral valve (St Jude Medical, Inc, St Paul, Minn), whose pivot is located in the left atrial portion of the

housing and whose leaflets open within the housing profile, has been believed to be a most suitable valve for chordal-preserving mitral valve replacement. However, several cases in which the preserved subvalvular apparatus were entrapped have been reported.³ The Carbomedics Standard mitral valve (Sulzer Carbomedics, Inc, Austin, Tex), which has been reported to cause few instances of thrombus or pannus formation,⁴ is designed so that the leaflets open beyond the housing profile, which might cause entrapment of retained subvalvular apparatus. With regard to stented bioprosthetic valves, a high-profile valve could cause left ventricular outflow obstruction in the subvalvular apparatus preserving mitral valve surgery as previously reported⁵ (Figure 1).

The On-X mitral valve is designed to prevent pannus ingrowth by high-profile inlet flare, although it is not so high as stented bioprostheses. Our experience showed the feasibility of chordal-preserving mitral valve replacement consisting of various techniques using the On-X valve, although it included a small number of cases. This technique using the On-X valve could bring better long-term outcome in patients undergoing mitral valve replacement.

References

- David TE, Uden DE, Strauss HD. The importance of the mitral apparatus in left ventricular function after correction mitral regurgitation. *Circulation*. 1983;68:76-82.
- Reardon MJ, David TE. Mitral valve replacement with preservation of the subvalvular apparatus. *Curr Opin Cardiol*. 1999;14:104-15.
- Agostini F, Click RL, Mulvagh SL, Abel MD, Dearani JA. Entrapment of subvalvular mitral tissue causing intermittent failure of a St Jude mitral prosthesis. *J Am Soc Echocardiogr*. 2000;13:1121-3.
- Konagai N, Yano H, Makimura S, Nishida K, Iida Y, Sato M, et al. A surgical case for a Carbomedics stuck valve due to thrombotic pannus formation in the mitral position. *Ann Thorac Cardiovasc Surg*. 2005; 11:186-9.
- Jett GK, Jett MD, Bosco P, van Rijk-Swicker GL, Clark RE. Left ventricular outflow tract obstruction following mitral valve replacement: effect of strut height and orientation. *Ann Thorac Surg*. 1986;42: 299-303.