everyone would prefer the less-invasive one. This issue is not quantifiable; on the other hand, suggesting the superiority of the OPCABG over PCI for a single-vessel disease on the basis of a non-updated analysis may be misleading for those physicians aiming at explaining a complete “state of the art.”

Luca Testa, MD, PhD^a

William van Gaal, MD^a

Giuseppe G. Biondi-Zoccai, MD^b

Antonio Abbate, MD^c

Ravinay Bhindi, MD, PhD^d

Pierfrancesco Agostoni^e

Institute of Cardiology^a

John Radcliffe Hospital

Oxford, United Kingdom

Division of Cardiology^b

University of Turin

Turin, Italy

Department of Medicine^c

Virginia Commonwealth University

Richmond, Va

Institute of Cardiology^d

Catholic University

Rome, Italy

Antwerp Cardiovascular Institute Middelheim^e

AZ Middelheim

Antwerp, Belgium

References

1. Bainbridge D, Cheng D, Martin J, Novick R; Evidence-based Peri-operative Clinical Outcomes Research (EPiCOR) Group. Does off-pump or minimally invasive coronary artery bypass reduce mortality, morbidity, and resource utilization when compared with percutaneous coronary intervention? A meta-analysis of randomized trials. *J Thorac Cardiovasc Surg.* 2007;133:623-31.
2. Smith SC, Feldman TE, Hirshfeld JW, et al. ACC/AHA/SCAI 2005 Guideline Update for Percutaneous Coronary Intervention: A Report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (ACC/AHA/SCAI Writing Committee to Update the 2001 Guidelines for Percutaneous Coronary Intervention). Available at: www.cardiosource.org.
3. Center for Overview, Meta-analysis and Evidence-based medicine Training (COMET). Available at: <http://www.comet.gs>. doi:10.1016/j.jtcvs.2007.03.053

Fixed subaortic stenosis

To the Editor:

We read with great interest the article entitled “Geometry of the Left Ventricular Outflow Tract in Fixed Subaortic Stenosis and Intact Ventricular Septum: An Echo-

cardiographic Study in Children and Adults.”¹ We have a few points to discuss.

First, we agree with the authors in choosing the wider mitral-aortic separation, steeper aortoseptal angle, smaller left ventricular outflow tract width, aortic valve dextroposition, increased left ventricle wall thickness, increased septal thickness, and indexed left ventricular end-diastolic and end-systolic diameters as the echocardiographic parameters to check in cases of subaortic stenosis.

Second, we think that left ventricular outflow tract obstruction in cases of Shone complex is a true challenge. We prefer not to intervene early, giving the heart and body time to grow, which is contrary to the general preference to intervene early in the other cases of subaortic stenosis. Discrete subaortic stenosis can be cured in most patients by membranectomy associated with myotomy or myectomy. Because the anatomic substrate is not addressed by these surgical techniques, however, recurrences are likely during long-term follow-up, particularly in patients who have undergone previous operations for an aortic coarctation and in patients with less than optimal relief of the left ventricular outflow tract gradient. In this subset of patients, the optimal surgical technique remains to be described. Intraoperative recording of the left ventricle-aorta gradient by transesophageal echocardiography or pressure measurement remains an important tool for more aggressive subaortic resection in case of a residual gradient greater than 30 mm Hg. According to the size and function of the aortic valve, the Ross-Konno procedure or the modified Konno procedure by patch septoplasty seems to be the appropriate surgical technique. Tunnel subaortic stenosis represents a more severe and challenging cause of left ventricular outflow tract obstruction, particularly when symptoms start early in life. Although multivariate analysis did not demonstrate this anatomic form as an independent risk factor for overall mortality and recurrence rates, in univariate analyses it was associated with a higher overall mortality rate and reoperation rate, along with other factors of the Shone complex (ie, hypoplastic aortic anulus, mitral stenosis, and existence of coarctation).²

Third, we have a remark on Table 3. We expected the indexed left ventricular end-systolic dimension to decrease postoperatively especially in group 1, which was not

the case. The left ventricular end-systolic dimension increased in group 1, although it decreased in groups 2 and 3. Do you have an explanation?

Sameh Ibrahim Sersar, MD

Ahmed A. Jamjoom, MD

Ghassan M. Baslaim, MD

Division of Cardiothoracic Surgery

Department of Cardiovascular Diseases

King Faisal Specialist Hospital and

Research Center

Jeddah, Saudi Arabia

References

1. Barkhordarian R, Wen-Hong D, Li W, Josen M, Henein M, Ho SY. Geometry of the left ventricular outflow tract in fixed subaortic stenosis and intact ventricular septum: an echocardiographic study in children and adults. *J Thorac Cardiovasc Surg.* 2007;133:196-203.
2. Serraf A, Zoghby J, Lacour-Gayet F, Houel R, Belli E, Galletti L, et al. Surgical treatment of subaortic stenosis: a seventeen-year experience. *J Thorac Cardiovasc Surg.* 1999;117:669-78.

doi:10.1016/j.jtcvs.2007.03.032

Reply to the Editor:

We thank Drs Sersar, Jamjoom, and Baslaim for their comments on our study.¹

Shone complex was originally described as comprising 4 coexisting obstructive lesions: Supravalvar ring of the left atrium, “parachute” deformity of the mitral valve, subaortic stenosis, and coarctation of the aorta are considerably more challenging than is an isolated subaortic stenosis.

On the basis of mean values, there are some differences in the left ventricular end-systolic dimension between groups, despite being statistically insignificant. This may be attributable to differences in left ventricular compliance to changes in loading conditions. This remains to be reexamined in a larger group of patients.

Reza Barkhordarian, MBBS, MRCS^a

Wei Li, MD, PhD^b

Michael Henein, MSc, PhD, FESC, FACC, FAHA^c

Department of Cardiac Morphology^a

Royal Brompton Hospital and Imperial College

Department of Cardiothoracic Surgery

St Bartholomew's Hospital

London, UK

Department of Cardiology^b

Royal Brompton Hospital

London, UK

Department of Cardiology^c
West Middlesex University Hospital and
Royal Brompton Hospital
London, UK

Reference

1. Barkhordarian R, Wen-Hong D, Li W, Josen M, Henein M, Ho SY. Geometry of the left ventricular outflow tract in fixed subaortic stenosis and intact ventricular septum: an echocardiographic study in children and adults. *J Thorac Cardiovasc Surg.* 2007;133:196-203.

doi:10.1016/j.jtcvs.2007.03.033

Familial recurrence of discrete membranous subaortic stenosis

We read with interest the article by Fatimi and colleagues¹ published in a recent issue of the *Journal*.

The authors report on 2 siblings affected by discrete membranous subaortic stenosis (DMSS), who were born to unaffected consanguineous parents. Literature reports of familial recurrence of DMSS were reviewed, and conclusions were that "review of the pedigrees and the information provided in the published reports does not reveal a clear inheritance pattern."

Nevertheless, we believe that Mendelian autosomal recessive inheritance may be considered as the main mechanism involved in the cause of concordant recurrence of DMSS in families on the basis of several observations. First, at least 8 families with recurrence of DMSS in siblings are reported in the literature,¹⁻⁵ including 2 sisters with DMSS born from unaffected consanguineous parents described by our group in 1993² (Figure 1, a). Second, parental consanguinity was observed in 4 families with recurrence of DMSS in siblings, including ours.^{1,2,4} Third, the recurrent cardiac defect is anatomically concordant in affected siblings from pedigrees with horizontal familial recurrence.¹⁻⁵

Obviously, genetic heterogeneity is implicated in the cause of DMSS, and the autosomal dominant inheritance with incomplete penetrance may be hypothesized in several instances.¹ In fact, parent-to-child transmission or segregation in uncle and nephew may be explained in this matter. In our 6-year (1993–1999) investigation of family history in patients affected by nonsyndromic DMSS, 3 additional pedigrees with familial recurrence of cardiac

defect have been detected (Figure 1, b-d). Cardiac defect was concordant in 1 family (Figure 1, b). In addition, a mother with DMSS had a child with aortic coarctation and valvar aortic stenosis (Figure 1, c), and a proband with DMSS had a cousin affected by tetralogy of Fallot (Figure 1, d). A multifactorial model of inheritance, suggesting that several chromosomal loci could interact together in association with environmental factors, may also be involved, particularly in sporadic cases and in families with recurrence of discordant cardiac defect.

Although DMSS is usually considered an "acquired" cardiac defect with delayed clinical presentation, the possibility of familial recurrence suggests a genetic substrate at least in some cases.

The genetic basis of DMSS is still unknown. In the future, molecular studies will provide an important aid in the interpretation of familial recurrence of DMSS. Genetic heterogeneity of DMSS probably includes the involvement of different genes. Similarly to that documented in some families segregating conotruncal heart defects, some genes may act accordingly to the autosomal recessive mechanism of inheritance,⁶ whereas others may follow the autosomal dominant model with incomplete penetrance.^{7,8}

Gerardo Piacentini, MD^a

Bruno Marino, MD^a

Maria Cristina Digilio, MD^b

Pediatric Cardiology, Department of Pediatrics^a
University "La Sapienza,"

Rome, Italy

Medical Genetics, Bambino Gesù Hospital^b

Rome, Italy

Figure 1. a-d, Four pedigrees with recurrent cardiac defect from our series of patients with DMSS. DMSS, Discrete membranous subaortic stenosis.

References

1. Fatimi SH, Ahmad U, Javed MA, Shamin S, Ahmad R. Familial membranous subaortic stenosis: review of familial inheritance patterns and case report. *J Thorac Cardiovasc Surg.* 2006;132:1484-6.
2. Digilio MC, Giannotti A, Marino B, Obregon MG, Dallapiccola B. Discrete membranous subaortic stenosis in siblings. *Eur J Pediatr.* 1993;152:622.
3. Gale AW, Cartmill TB, Bernstein L. Familial subaortic membranous stenosis. *Aust N Z J Med.* 1974;4:576-81.
4. Urbach J, Glaser J, Balkin J, Rosenmann D, Levy R, Marin G, et al. Familial membranous subaortic stenosis. *Cardiology.* 1985;72:214-7.
5. Abdallah H, Toomey K, O'Riordan AC, Davidson A, Marks LA. Familial occurrence of