the myocardium (reflected, for instance, by a downregulation of respiratory chain enzymes),³ with obvious beneficial effects on cell survival. These effects are probably most pronounced in the mechanically and biologically overburdened infarct border zone. Therefore, it does not seem surprising that the authors found less apoptosis and more markers of vital cells—including c-kit- and Sca-1-expressing cells—right there. To study true regeneration processes, however, unloading of the hearts later after myocardial infarction might produce more meaningful data. Here, the effects of myocardial rescue by unloading have abated, enabling the researcher to concentrate on true regeneration processes instead of salvage of pre-existing cells. We expect that the differences in regenerative capacity between unloaded and loaded hearts would then be much smaller, similar to the clinical observation that weaning from left ventricular assist device support during acute infarction or in nonischemic cardiomyopathy is sometimes possible,⁴ but very rarely in patients with chronic ischemic heart failure.

Taken together, the data presented by Suzuki and colleagues add nicely to the evidence supporting the concept of rapid hemodynamic load reduction in acute infarction. Whether unloading alone is able to also initiate true myocardial regeneration, especially in the clinical setting, remains to be seen.

Christof Stamm, MD
Boris Nasser, MD
Roland Hetzer, MD
Deutsches Herzzentrum Berlin
Cardiothoracic and Vascular Surgery
Berlin, Germany

References

1. Suzuki R, Li TS, Mikamo A, Takahashi M, Ohshima M, Kubo M, et al. The reduction of hemodynamic loading assists self-regeneration of the injured heart by increasing cell proliferation, inhibiting cell apoptosis, and inducing stem-cell recruitment. *J Thorac Cardiovasc Surg.* 2007;133:1051-8.
2. Potapov EV, Weng Y, Hausmann H, Kopitz M, Pasic M, Hetzer R. New approach in treatment of acute cardiogenic shock requiring mechanical circulatory support. *Ann Thorac Surg.* 2003;76:2112-4.
3. Bugger H, Leippert S, Blum D, Kahle P, Barleon B, Marme D, et al. Subtractive hybridization for differential gene expression in mechanically unloaded rat heart. *Am J Physiol Heart Circ Physiol.* 2006;291:H2714-22.

4. Dandel M, Weng Y, Siniawski H, Potapov E, Lehmkuhl HB, Hetzer R. Long-term results in patients with idiopathic dilated cardiomyopathy after weaning from left ventricular assist devices. *Circulation.* 2005;112(9 Suppl):I37-45.

doi:10.1016/j.jtcvs.2007.05.065

Reply to the Editor:

We greatly appreciate the letter from Stamm and colleagues with its well-directed comments on our article. As they suggest, we need to clarify many problems about hemodynamic unloading for myocardial repair, including “regeneration versus rescue,” “acute versus chronic,” and “clinical patient versus experimental model.”

First, rescue and regeneration are very different biologic processes. The former involves time-limited work to rescue the surviving myocardium soon after damage, whereas the latter is a slow and long-term process of regenerating new myocardium for repairing the injured heart. We recognize that the regenerative potency of the injured heart is poor because of the very limited proliferating potency of cardiomyocytes.¹ Inasmuch as we removed hemodynamic loading soon after infarction in this study, it is possible that hemodynamic unloading rescued much of the surviving myocardium within the injured heart and contributed to the dramatic improvement in the infarcted area and the left ventricular wall thickness, as shown in Figure 1 of our article.² However, the increased number of Ki-67-positive cells indicated an accelerated biologic process of regeneration in the unloading infarcted heart. Furthermore, although we did not follow their fate, the increased number of stem cells in the unloading infarcted heart also played a role in myocardial repair. On the basis of this evidence, we concluded that the reduced hemodynamic loading assists self-regeneration of the injured heart.

Second, compared with the acute phase, the chronically injured heart was characterized by compensatory hypertrophy of the surviving myocardium, the senescence of cardiac stem cells,³ and many other neurohormonal changes. Thus, it is worth investigating whether hemodynamic unloading would assist the regeneration of a chronically injured heart and the capacity and speed of this regeneration.

The final problem relates to the self-regenerative capacity of the injured heart in both patients and experimental animals. In

contrast to healthy young animals, many of the patients with heart failure were older and had coexisting systemic diseases, such as diabetes, hyperlipidemia, and hypertension, which contributed to the senescence of stem cells. Naturally, the proliferating potency of cardiomyocytes and the number of cardiac stem cells in older patients with heart failure would be much weaker than those in healthy young animals.⁴

In all, the functional recovery of the failing heart achieved by left ventricular assist device support involves complex mechanisms, which will be affected by many factors.⁵ Although our experimental data suggest that hemodynamic unloading assists self-regeneration of the injured heart, further study is warranted to estimate the functional contribution of myocardial regeneration in patients subjected to left ventricular assist device support.

Tao-Sheng Li, MD

Ryo Suzuki, MD

Kimikazu Hamano, MD

Department of Surgery and Clinical Science
Yamaguchi University
Graduate School of Medicine
Ube, Yamaguchi, Japan

References

1. Beltrami AP, Urbanek K, Kajstura J, Yan SM, Finato N, Bussani R, et al. Evidence that human cardiac myocytes divide after myocardial infarction. *N Engl J Med.* 2001;344:1750-7.
2. Suzuki R, Li TS, Mikamo A, Takahashi M, Ohshima M, Kubo M, et al. The reduction of hemodynamic loading assists self-regeneration of the injured heart by increasing cell proliferation, inhibiting cell apoptosis, and inducing stem-cell recruitment. *J Thorac Cardiovasc Surg.* 2007;133:1051-8.
3. Urbanek K, Torella D, Sheikh F, De Angelis A, Nuszynska D, Silvestri F, et al. Myocardial regeneration by activation of multipotent cardiac stem cells in ischemic heart failure. *Proc Natl Acad Sci U S A.* 2005;102:8692-7.
4. von Harsdorf R, Poole-Wilson PA, Dietz R. Regenerative capacity of the myocardium: implications for treatment of heart failure. *Lancet.* 2004;363:1306-13.
5. Wohlschlaeger J, Schmitz KJ, Schmid C, Schmid KW, Keul P, Takeda A, et al. Reverse remodeling following insertion of left ventricular assist devices (LVAD): a review of the morphological and molecular changes. *Cardiovasc Res.* 2005;68:376-86.

doi:10.1016/j.jtcvs.2007.06.030