Francesca di Marco, MD, PhD
Gino Gerosa, MD
Department of Cardiological
Thoracic and Vascular Sciences
Division of Cardiac Surgery
Padua University Medical School
Padua, Italy

References

1. Vincentelli A, Wautot F, Juthier F, Fouquet O, Corseaux D, Marechaux S, et al. In vivo autologous recellularization of a tissue-engineered heart valve: are bone marrow mesenchymal stem cells the best candidates? *J Thorac Cardiovasc Surg.* 2007;134:424-32.
2. Menasche P, Hagege AA, Scorsin M, Pouzet B, Desnos M, Duboc D. Myoblast transplantation for heart failure. *Lancet.* 2001;357:279-80.
3. Bertipaglia B, Ortolani F, Petrelli L, Gerosa G, Spina M, Pauletto P, et al. Cell characterization of porcine aortic valve and decellularized leaflets repopulated with aortic valve interstitial cells: the VESALIO Project (Vitalitate Exornatum Succedaneum Aorticum Labore Ingenioso Obtenibitur). *Ann Thorac Surg.* 2003;75:1274-82.

doi:10.1016/j.jtcvs.2007.10.075

Reply to the Editor:

We thank Drs Di Marco and Gerosa for considering that injection of mesenchymal stem cells (MSCs) in decellularized scaffolds before in vivo implantation represents an innovative concept in the valvular field, and we completely agree that this concept requires a careful validation. However, they expressed some concern about our conclusion that bone marrow-derived mononuclear cell injection induced a structural deterioration of the scaffold and that MSCs induced a protective effect.

We believe that the observation of a deleterious effect of bone marrow-derived mononuclear cells is an interesting point because these cells had never been tested in the field of valve tissue engineering, whereas they are currently used in clinical assays in the field of myocardial failure. Also, we agree that endothelialization and scaffold recolonization are 2 distinct processes that do not always take place together. However, Bertipaglia and colleagues¹ have demonstrated that 2 weeks after in vitro interstitial cell seeding into decellularized leaflets, grafted cells were found penetrating the decellularized scaffold and expressed various immunologic differentiation patterns, such as fibroblasts, myofibroblasts, smooth muscle cells, and

also endothelial cells. Similarly, we observed a significant recolonization after injection of MSCs, and moreover, as stated in our article, pulmonary leaflets exhibited a typical organization in 3 layers (ie, fibrosa, spongiosa, and ventricularis) in each animal of the MSC group.²

Finally, several lines of evidence indicate that in vivo recellularization of valve scaffolds is no longer a “nonvalidated route.” In a previous work we demonstrated evidence of spontaneous recellularization of a decellularized valve.³ Such results were emphasized by Dohmen and coworkers⁴ comparing in vitro seeded and nonseeded valves implanted in sheep. After 6 months, valves from both groups exhibited full recolonization of the leaflets, with comparable hemodynamic behavior and valve remodeling.

We recognize that further studies are needed to compare spontaneous recolonization of an appropriated scaffold and the effects of in situ injection of autologous cells. Also, we have to investigate how injected autologous MSCs improve in situ recolonization by myofibroblasts and endothelial cells or improve host cell migration. However, the challenge seems now to understand and to favor in vivo, rather than in vitro, valve scaffold recolonization.

André Vincentelli, MD, PhD^{a,b}

Francis Juthier, MD^{a,b}

Brigitte Jude, MD, PhD^{a,b}

Centre Hospitalier Régional^a

Universitaire de Lille

Pôle de Chirurgie Cardiovasculaire

Institut d'Hématologie Transfusion

Lille, F-59037 France;

Université de Lille 2^b

EA 2693

Faculté de Médecine

Lille, F-59045 France

References

1. Bertipaglia B, Ortolani F, Petrelli L, Gerosa G, Spina M, Pauletto P, et al. Cell characterization of porcine aortic valve and decellularized leaflets repopulated with aortic valve interstitial cells: the VESALIO Project (Vitalitate Exornatum Succedaneum Aorticum Labore Ingenioso Obtenibitur). *Ann Thorac Surg.* 2003;75:1274-82.
2. Vincentelli A, Wautot F, Juthier F, Fouquet O, Corseaux D, Marechaux S, et al. In vivo autologous recellularization of a tissue-engineered heart valve: are bone marrow mesenchymal stem cells the best candidates? *J Thorac Cardiovasc Surg.* 2007;134:424-32.
3. Juthier F, Vincentelli A, Gaudric J, Corseaux D, Fouquet O, Calet C, et al. Decellularized heart

valve as a scaffold for in vivo recellularization: deleterious effects of granulocyte colony-stimulating factor. *J Thorac Cardiovasc Surg.* 2006;131:843-52.

4. Dohmen PM, da Costa F, Yoshi S, Lopes SV, da Souza FP, Vilani R, et al. Histological evaluation of tissue-engineered heart valves implanted in the juvenile sheep model: is there a need for in-vitro seeding? *J Heart Valve Dis.* 2006;15:823-9.

doi:10.1016/j.jtcvs.2007.12.030

Percutaneous aortic valve replacement with the CoreValve bioprosthesis

To the Editor:

We read with interest the recent article by Marcheix and colleagues¹ in the *Journal* dealing with percutaneous aortic valve implantation. They reported their experience using the second generation (21F) of the CoreValve prosthesis (CoreValve, Inc, Paris, France) in 10 consecutive high-risk surgical patients (median logistic EuroSCORE of 32%). The prosthesis was successfully delivered in all cases, with a doubling of the aortic valve area (from 0.57 ± 0.19 to 1.2 ± 0.35 cm²) and a significant decrease in transvalvular gradient.

We would like to comment on these results and also ask for some clarifications regarding their data:

Prosthesis sizing is an important issue. On one hand, undersizing may be responsible for valve migration (owing to poor anchoring), perivalvular leak, or patient-prosthesis mismatch. On the other hand, oversizing may theoretically lead to coronary obstruction or favor atrioventricular block. There is actually only one available size of the CoreValve prosthesis, with a 21-mm bioprosthesis implanted within the stent frame. This valve is at the present time indicated when the aortic annulus diameter ranges from 20 to 27 mm.² Although the authors did not report the annulus size in their group of patients, we do not understand how proper sizing could be achieved with a single-sized bioprosthesis.

Aortic bicuspidy is observed in half of cases of severe aortic stenosis.³ In our experience, misdeployment of a valved stent is likely to occur within a bicuspid aortic valve,^{4,5} which may alter its long-term durability. Did the authors face a case with a bicuspid aortic valve or have they systematically preoperatively screened the patients and excluded for the endovascular procedure those with a bicuspid aortic valve?

Although this study summarizes an initial experience, we thank the authors for sharing with us their results and for their contribution to this important new field of interventional therapy.

Rachid Zegdi, MD, PhD^{a,b}

Paul Achouh, MD, PhD^a

Jean-Noël Fabiani, MD^{a,b}

Assistance Publique-Hôpitaux de Paris AP-HP

Service de Chirurgie Cardiovasculaire

Hôpital Européen Georges Pompidou

Paris, France^a

Université René Descartes

Paris, France^b

References

1. Marcheix B, Lamarche Y, Berry C, Asgar A, Laborde J-C, Basmadjian A, et al. Surgical aspects of endovascular retrograde implantation of the aortic CoreValve bioprosthesis in high-risk older patients with severe symptomatic aortic stenosis. *J Thorac Cardiovasc Surg*. 2007;134:1150-6.
2. Grube E, Schuler G, Buellesfeld L, Gerckens U, Linke A, Wenaweser P, et al. Percutaneous aortic valve replacement for severe aortic stenosis in high-risk patients using the second- and current third-generation self-expanding CoreValve prosthesis. *J Am Coll Cardiol*. 2007;50:69-76.
3. Roberts WC, Ko JM. Frequency by decades of unicuspid, bicuspid, and tricuspid aortic valves in adults having isolated aortic valve replacement for aortic stenosis, with or without associated aortic regurgitation. *Circulation*. 2005;111:920-5.
4. Zegdi R, Khabbaz Z, Ciobotaru V, Noghin M, Deloche A, Fabiani JN. Calcific bicuspid aortic stenosis: a questionable indication for endovascular valve implantation? *Ann Thorac Surg*. In press.
5. Zegdi R, Ciobotaru V, Noghin M, Sleilaty G, Lafont A, Latrémouille C, et al. Is it reasonable to treat all calcific stenotic aortic valves with a valved stent? Results from a human anatomic study in adults. *J Am Coll Cardiol*. In press.
doi:10.1016/j.jtcvs.2007.11.054

Reply to the Editor:

We appreciate the authors' concerns about the quality of our work published the *Journal of Thoracic and Cardiovascular Surgery*. We totally agree with the two underlined points and we would like to simply answer the two questions.

It is true that prosthesis sizing is an important issue. All the reported patients had an aortic annulus diameter ranging from 20 to 23 mm. Smaller and larger annuli were considered as contraindicated so as to

allow correct anchorage of the prosthesis in the annulus.

All the patients had the echographic aspect of a tricuspid aortic valve. We cannot answer Zegdi and colleagues because we do not have any experience with percutaneous aortic valve implantation in bicuspid valves. Bicuspid valves are generally more calcified than tricuspid valves and frequently come with a moderate dilatation of the aorta. They are less favourable for currently used percutaneous aortic valves.

Bertrand Marcheix, MD

Raymond Cartier, MD

Montreal, Quebec, Canada

doi:10.1016/j.jtcvs.2008.03.008

Does the method of lung preservation influence outcome after transplantation? An analysis of 681 consecutive procedures

To the Editor:

We read with great interest the study by Ganesh and colleagues¹ that compared various graft preservation methods in lung transplantation. The authors rigorously assessed the association between the lung preservation method and the posttransplant survival up to 3 years in a large cohort of patients; they failed to detect any association. However, we think that the way the graft ischemic time was calculated and included in the analysis may have biased the results. Graft ischemic time has been shown to be associated with survival after lung transplantation in several studies.^{2,3} In these studies, in case of double-lung transplantation, graft ischemic time was defined as the time from donor crossclamping to reperfusion of the second lung. In their study, Ganesh and colleagues considered ischemic time as the time from donor crossclamping to reperfusion of the first lung only. This could lead to incorrect adjustment. Second, we have found that the relationship between graft ischemic time and survival could be of exponential form, the graft ischemic time altering the survival of recipients only when it exceeds 4 to 6 hours.³ Unfortunately, the authors did not report on the distribution of graft ischemic time by the preservation method. Given the incidence of local donors in their cohort, we suspect that graft ischemic time was rather short in most patients.

This point is essential, because the efficacy of different graft preservation solutions may prove similar for short graft ischemic times and different for longer ischemic times. We think that the analysis of adjusted survival by preservation solution for patients with a long ischemic time would have yielded more insight in the efficacy of preservation solutions. However, the number of patients in each group would have probably been low, precluding powerful statistical analysis. The authors report no interaction between the graft ischemic time and the preservation solution used; however, interaction testing is not powerful enough. We agree that the way to definitely answer this question is to conduct a randomized controlled trial, and we think that such a trial should stratify patients by the expected graft ischemic time or the surgical procedure (single vs bilateral lung transplantation).

Gabriel Thabut, MD^a

Yves Castier, MD^b

Hervé Mal, MD^a

Service de pneumologie B et transplantation pulmonaire^a

Service de chirurgie thoracique et vasculaire^b
Hôpital Bichat et Université Paris 7,
Paris, France

References

1. Ganesh JS, Rogers CA, Banner NR, Bonser RS. Does the method of lung preservation influence outcome after transplantation? An analysis of 681 consecutive procedures. *J Thorac Cardiovasc Surg*. 2007;134:1313-21.
2. Novick RJ, Bennett LE, Meyer DM, Hosenpud JD. Influence of graft ischemic time and donor age on survival after lung transplantation. *J Heart Lung Transplant*. 1999;18:425-31.
3. Thabut G, Mal H, Cerrina J, Darteville P, Dromer C, Velly JF, et al. Graft ischemic time and outcome of lung transplantation: a multicenter analysis. *Am J Respir Crit Care Med*. 2005;171:786-91.
doi:10.1016/j.jtcvs.2007.11.059

Reply to the Editor:

We appreciate the comments made by Thabut and colleagues on our article on lung preservation.¹ We agree that graft ischemia time may be an important predictor to survival after lung transplantation, particularly if this is particularly prolonged. The study by Novick and colleagues² did not find any influence of graft ischemia time alone on survival, but there was a significant