RIFLE criteria in aortic arch surgery: The further role of surgical subgroup

To the Editor:

I read with great interest the recent article by Arnaoutakis and colleagues¹ detailing their application of the RIFLE (risk, injury, failure, loss, end stage) criteria in aortic arch surgery. Arnaoutakis and colleagues¹ clearly demonstrated a significant relationship between degree of renal injury and operative mortality. Their series of 267 patients, however, was heterogeneous with respect to aortic pathology: 35.6% of the cases (95/267) were aortic dissection, with 74.7% (71/95) being acute and necessitating emergency surgery with deep hypothermic circulatory arrest.

The clinical presentation of acute aortic dissection significantly determines operative risk and mortality, particularly when it is associated with organ ischemia.² Thus this confounding effect of an aortic arch subgroup with distinctive presentations ideally merits a de novo analysis in a large acute aortic dissection series. This goal would require many years to achieve for a single center, however, even a single, high-volume experienced center. As a result, current single-center series, including our own,³ have included acute aortic dissection as a subgroup and thus been unable to escape the confounding effects of the mixed aortic arch cohort.³ I look forward to future multicenter trials with adequate power to examine the RIFLE criteria in aortic arch surgical cohorts stratified by aortic pathology.

I congratulate Arnaoutakis and colleagues¹ again on their important contribution. I look forward to their comments about this aspect of trial design.

John G. T. Augoustides, MD, FASE
Assistant Professor, Department of
Anesthesiology and Critical Care
Hospital of the University of Pennsylvania
Philadelphia, PA 19104

References

1. Arnaoutakis GJ, Bihorac A, Martin TD, Hess PJ, Klodell CT, Ejaz AA, et al. RIFLE criteria for acute kidney injury in aortic arch surgery. *J Thorac Cardiovasc Surg.* 2007; 134:1554-61.
2. Geirsson A, Szeto WY, Pochettino A, McGarvey, Keane MG, Woo YJ, et al. Significance of malperfusion syndromes prior to contemporary surgical repair for acute type A dissection: outcomes and need for additional revasculari-

zations. *Eur J Cardiothorac Surg.* 2007;32: 255-62.

3. Augoustides JG, Pochettino A, Ochroch EA, Cowie D, Weiner J, Gambone AJ, et al. Renal dysfunction after thoracic aortic surgery requiring deep hypothermic circulatory arrest: definition, incidence, and clinical predictors. *J Cardiothorac Vasc Anesth.* 2006;20: 673-7.

doi:10.1016/j.jtcvs.2007.12.050

Reply to the Editor:

We appreciate the interest of Dr Augoustides in our article,¹ and he raises a valid point that the subgroup of patients with acute aortic dissection are at even higher risk for acute kidney injury because of their emergency presentation and variable malperfusion. Patients with acute dissection accounted for 71 of the 97 emergency operations in our series. In the multivariate analysis, emergency surgery (and accordingly acute dissection) was not identified as a risk factor for either acute kidney injury or mortality. As Dr Augoustides noted, however, it would require many years for a single center to examine this question.

The RIFLE criteria provide standardized outcome definitions to study acute kidney injury in patients undergoing arch aortic surgery—a group known to be at significant risk for renal injury. Standardized definitions are the first step if we are to compare outcomes at aortic centers interested in acute kidney injury.

To take the next step—examination of renal protective strategies—we concur with Dr Augoustides that patients should be stratified according to aortic pathology and also that it will take the cooperation of several centers with large aortic experiences. The question then arises, which protective strategies could we study? We and others have been intrigued by the potentially favorable renal effects of nesiritide (Natrecor; Scios Inc, Fremont, Calif).^{2-5*}

It is important to note that even less severe kidney injury after cardiac surgery has been found to be independently associated with in-hospital and long-term mortality.^{6,7} Accordingly, with interest from the center of Dr Augoustides, our own, and others, perhaps it is now time to ask the National Institutes of Health to support a randomized, multicenter trial adequately powered to determine whether nesiritide (or other potential agents) can afford high-risk patients undergoing thoracic aortic surgery protection against acute kidney injury.

Thomas M. Beaver, MD, MPH

Azra Bihorac, MD

Ahsan A. Ejaz, MD

Division of Cardiovascular Surgery

University of Florida

Gainesville, FL 32610

*Disclosure: Authors have received research support from Scios Inc.

References

1. Arnaoutakis GJ, Bihorac A, Martin TD, Hess PJ Jr, Klodell CT, Ejaz AA, et al. RIFLE criteria for acute kidney injury in aortic arch surgery. *J Thorac Cardiovasc Surg.* 2007; 134:1554-61.
2. Chen HH, Sundt TM, Cook DJ, Heublein DM, Burnett JC Jr. Low dose nesiritide and the preservation of renal function in patients with renal dysfunction undergoing cardiopulmonary-bypass surgery: a double-blind placebo-controlled pilot study. *Circulation.* 2007; 116(11 Suppl):1134-8.
3. Beaver TM, Winterstein AG, Shuster JJ, Gerhard T, Martin T, Alexander JA, et al. Effectiveness of nesiritide on dialysis or all-cause mortality in patients undergoing cardiothoracic surgery. *Clin Cardiol.* 2006;29:18-24.
4. Mentzer RM Jr, Oz MC, Sladen RN, Graeve AH, Hebel RF Jr, Luber JM Jr, et al. Effects of perioperative nesiritide in patients with left ventricular dysfunction undergoing cardiac surgery: the NAPA Trial. *J Am Coll Cardiol.* 2007;49:716-26.
5. Ejaz AA, Heinig ME, Kazory A, Bihorac A, Hobson CE, Beaver TM. The rise and fall of natriuretic peptides in acute kidney injury: a misunderstood relationship? *Rev Cardiovasc Med.* 2007;(8 Suppl 5):S32-7.
6. Thakar CV, Worley S, Arrigain S, Yared JP, Paganini EP. Influence of renal dysfunction on mortality after cardiac surgery: modifying effect of preoperative renal function. *Kidney Int.* 2005;67:1112-9.
7. Loeff BG, Epema AH, Smilde TD, Henning RH, Ebels T, Navis G, et al. Immediate postoperative renal function deterioration in cardiac surgical patients predicts in-hospital mortality and long-term survival. *J Am Soc Nephrol.* 2005;16:195-200.

doi:10.1016/j.jtcvs.2008.03.012

Reply to the Editor:

I thank Dr Pocar for promoting a discussion about our article and technical proposition. In fact, the juxtaposition of the right free wall over the septum is not new and was applied for the first time by Boer and Boer in 1998;¹ they called it “septal stabilization.”

The authors^{2,3} cited by Dr Pocar made an intelligent association of 2 techniques, that is, septal stabilization¹ and infarction exclusion,⁴ with good results.

In our study, we proposed the juxtaposition of both the right and left free walls over the septum, and this is the originality of the