looks quite different depending on one's perspective. From the aortic or ventricular side, the AML clearly begins immediately below the semilunar attachment, whereas from the atrial side, because of the thickness of the LA itself, the base of the AML seems to be well below the attachment of the aortic valve cusp. The muscle of the LA essentially folds down over the base of the AML.

Again we thank Drs Nežić, Knežević, and Borović for their interest and welcome the debate. What could be more fun than spirited disagreement among colleagues!

Jayesh Dhareshwar, MD^a
Dylan V. Miller, MD^b
Thoralf M. Sundt, III, MD^a
Division of Cardiovascular Surgery^a
Department of Anatomic Pathology^b
Mayo Clinic, Rochester, MN 55905

References

1. Nicks R, Cartmill T, Bernstein L. Hypoplasia of the aortic root. The problem of aortic valve replacement. *Thorax*. 1970;25:339-46.
2. Anderson RH. Demystifying the anatomic arrangement of the aortic valve. *Eur J Cardiothorac Surg*. 2006;29:1006-7.
3. Anderson RH. Clinical anatomy of the aortic root. *Heart*. 2000;84:670-3.
4. Sutton JP 3rd, Ho SY, Anderson RH. The forgotten interleaflet triangles: a review of the surgical anatomy of the aortic valve. *Ann Thorac Surg*. 1995;59:419-27.
5. Nuñez L, Gil Aguado M, Pinto AG, Larrea JL. Enlargement of the aortic annulus by resecting the commissure between the left and noncoronary cusps. *Tex Heart Inst J*. 1983;10:301-3.

doi:10.1016/j.jtcvs.2008.03.009

Endovascular treatment of mycotic aortic aneurysms

To the Editor:

We read with interest the article by Hsu and Lin¹ in which they retrospectively reviewed their patients who received surgical treatment for infected aortic aneurysms at the aortic arch. Mycotic aortic aneurysm treatment is a challenging issue. Since the beginning of modern cardiovascular surgery, surgeons have searched for better treatment options because of the high mortality and morbidity rates of the classic surgical methods. The authors' results also confirm this.¹

With the increasing experience and refinements in graft technology, endovascular grafts are frequently used in the treatment of

aneurysms at critical segments of the aorta, such as thoracic and arch levels,² and even to the ascending aorta by means of hybrid therapy or alone.³

On the other hand, although it seems completely opposite to the nature of the treatment of infected aneurysms, endoluminal stent grafting is also applied for the treatment of mycotic aneurysms. Although the long-term results are not presented, the short and mid-term outcomes of this treatment modality have been promising when compared with conventional surgical techniques (ie, extensive debridement and artificial graft replacement or extra-anatomic bypass procedures).^{4,5}

We believe that endovascular stent graft repair of mycotic aortic aneurysms, especially at the critical segments of the aorta and in debilitated patients, is an attractive alternative option with low mortality and morbidity rates in the midterm. However, long-term results are definitely required before this becomes the standard therapy. It would be helpful if the authors would have commented on this issue in their article.

Dilek Erer, MD^a
Maciej Banach, MD^b
Murat Ugurlucan, MD^c
Department of Cardiovascular Surgery^a
Gazi University Medical Faculty
Ankara, Turkey
Department of Cardiology^b
Lodz University Medical Faculty
Lodz, Poland
Department of Cardiovascular Surgery^c
Istanbul University Istanbul Medical Faculty
Istanbul, Turkey

References

1. Hsu RB, Lin FY. Surgery for infected aneurysm of the aortic arch. *J Thorac Cardiovasc Surg*. 2007;134:1157-62.
2. Ugurlucan M, Akyol Y, Guven K, Poyanli A, Alpogut U, Rozanes I, et al. Treatment of chronic type B aortic dissection by endovascular grafting in a previously CABG patient. *Acta Chir Belg*. 2007;107:419-23.
3. Mussa FF, LeMaire SA, Bozinovski J, Coselli JS. An entirely endovascular approach to the repair of an ascending aortic pseudoaneurysm. *J Thorac Cardiovasc Surg*. 2007;133:562-3.
4. Ugurlucan M, Alpogut U. Endoluminal stenting of thoracic aorta mycotic aneurysms. *Eur J Cardiothorac Surg*. 2007;32:945-6.
5. Kpodonu J, Williams JP, Ramaiah VG, Diethrich EB. Endovascular management of a descending thoracic mycotic aneurysm: mid-term follow-up. *Eur J Cardiothorac Surg*. 2007;32:178-9.

doi:10.1016/j.jtcvs.2007.11.060

Reply to the Editor:

We appreciate Dr Ugurlucan's comments on the issue of mycotic aortic aneurysms treated by endovascular approach. We believe that the endovascular stent graft is superior in some ways to open surgical repair for the treatment of mycotic aortic aneurysms, especially in the emergency setting for debilitated patients with critical aortic segment involvement. In a recent published literature review by Kan and colleagues,¹ the endovascular treatment strategy was noted to have 30-day and 2-year survivals of 89.6% and 82.2%, respectively. To date, however, there is still no level I evidence to convince us that endovascular treatment is the criterion standard for mycotic aortic aneurysm.

In Taiwan, the abdominal aortic stent graft was approved by Department of Health for clinical use only after July 2005, and the thoracic stent graft was approved after November 2006. In our published article,² patients were treated through 2006, before the endovascular stent graft era in Taiwan. Since 2006, we have performed endovascular stent graft placement in patients with ruptured mycotic aortic aneurysm. In our preliminary data, there have been 4 patients with ruptured mycotic abdominal and thoracic aortic aneurysms treated by the endovascular technique. Of these, 1 patient required a total visceral artery debranching procedure for the ruptured suprarenal mycotic aneurysm simultaneously with the stent graft placement. The pathogen is similar to our published database² as nontyphoid *Salmonella*. There was one in-hospital death with new development of aortoduodenal fistula. There was another death later than 30 days from the non-aneurysm-related cause of uncontrolled septic shock. The other 2 patients require long-term oral antibiotic therapy. There were no cerebral vascular events or stent graft-related deaths among these 4 patients.

This seems to be a result comparable to that of open surgical repair. We are still not sure, however, whether for those with aorto-esophageal aneurysm, aortobronchial aneurysm, or residual abscess in the image study further débridement and elective open surgical grafting will be necessary after stent graft placement, once the patient condition has been stabilized. In addition, for mycotic aneurysms involving the aortic arch or visceral arterial segment, the hybrid debranching procedure or fenestrated

branching technique for the stent graft placement still carries some morbidity and mortality risk. The optimal use of endovascular stent graft for mycotic aortic aneurysm needs to be clarified in further studies.

I-Hui Wu, MD

Ron-Bin Hsu, MD

Fang-Yue Lin, MD, PhD

Department of Surgery

National Taiwan University Hospital

National Taiwan University College of Medicine

Taipei, Taiwan

References

1. Kan CD, Lee HL, Yang YJ. Outcome after endovascular stent graft treatment for mycotic aortic aneurysm: a systematic review. *J Vasc Surg.* 2007;46:906-12.
2. Hsu RB, Lin FY. Surgery for infected aneurysm of the aortic arch. *J Thorac Cardiovasc Surg.* 2007;134:1157-62.

doi:10.1016/j.jtcvs.2008.03.013

Grey zone area for Glenn and future Fontan candidates

To the Editor:

We enjoyed the very good article by Scheurer and colleagues.¹ We have a few comments.

Inhaled nitric oxide produces significant reduction in Glenn pressures and improvement in systemic perfusion and pulmonary gas exchange in patients with marked increases of Glenn pressures after bidirectional Glenn anastomosis. Patients who do not respond should be investigated for an anatomic lesion.²

Glenn shunt mortality was higher with surgical intervention before age 120 days but was not affected by surgical intervention during the respiratory virus season. Operations in the youngest age groups might adversely affect mortality.³

There were no differences in morbidity and mortality outcomes after the bidirectional Glenn operation between the Norwood-Right ventricle to pulmonary artery and Norwood-Blalock-Taussig Shunt groups. Despite younger age at presentation, the Norwood-Right ventricle to pulmonary artery patients had better growth rate, which might have contributed to the similar postoperative outcomes.⁴

Atrial septal defects should be left open for all patients with severe right heart hypoplasia and in patients with postbypass right atrial pressures of more than 12 mm Hg.⁵

The advantages of incorporating a hypoplastic ventricle to partly support the pulmonary circulation are the following: (1) ability to increase the cardiac output; (2) adaptation to exercise; (3) maintenance of pulsatile flow in the pulmonary circulation; (4) flexibility to increased pulmonary vascular resistance; (5) circulation at low venous pressure in the Inferior Vena Cava system; and (6) capability of a hypoplastic right heart to adequately handle the reduced preload.⁶

There is a major debate for the gray-zone high-risk patients who do not fulfill the Choussat commandments. The current protocol in those patients is first to review all the data in detail. Full consideration is given as to whether there are any surgically (or interventional) correctable lesions, such as atrioventricular valve regurgitation amenable to repair or isolated stenoses or hypoplasia within the central pulmonary arteries. Ventricular function, if impaired, might be optimized by altering medical management (vasodilators, diuretic therapy, and trial of oral enoximone, if appropriate). Evidence of increased pulmonary vascular resistance is treated with a course of pulmonary vasodilator therapy (sildenafil or bosentan). The patients are then reassessed for suitability for Fontan completion. The evidence would suggest that only 2 of the original 10 commandments of Choussat carry significant weight for both early and late outcome over the Fontan procedure and that they are preoperative ventricular function and preoperative pulmonary artery pressures of greater than 15 mm Hg.⁷

We found that right ventricular morphology; single ventricle; intact septum; semilunar valve abnormalities; diastolic dysfunction; myocardial hypertrophy, dilatation, or both; older age; bilateral superior vena cavae; pulmonary airway disease; and hypotonia all carry a worse prognosis and a tough postoperative course.

To be right ventricular morphology; single ventricle; intact septum; semilunar valve abnormalities; diastolic dysfunction; myocardial hypertrophy, dilatation, or both; older age; bilateral superior vena cavae; Interrupted IVC; pulmonary airway disease; and hypotonia all carry a worse prognosis and a tough postoperative course.

Sameh Ibrahim Sersar, MD

Division of Cardiothoracic Surgery

Department of Cardiovascular Diseases

King Faisal Specialist Hospital and

Research Center

Jeddah, Saudi Arabia

References

1. Scheurer MA, Hill EG, Vasuki N, Maurer S, Graham EM, Bandisod V, et al. Survival after bidirectional cavopulmonary anastomosis: analysis of preoperative risk factors. *J Thorac Cardiovasc Surg.* 2007;134:82-9.
2. Agarwal HS, Churchwell KB, Doyle TP, Christian KJ, Drinkwater DC, Byrne DW, et al. Inhaled nitric oxide use in bidirectional Glenn anastomosis for elevated Glenn pressures. *Ann Thorac Surg.* 2006;81:1429-34.
3. Nicolas RT, Hills C, Moller JH, Huddleston CB, Johnson MC. Early outcome after Glenn shunt and Fontan palliation and the impact of operation during viral respiratory season: analysis of a 19-year multi-institutional experience. *Ann Thorac Surg.* 2005;79:613-7.
4. Lai L, Laussen PC, Cua CL, Wessel DL, Costello JM, del Nido PJ, et al. Outcomes after bidirectional Glenn operation: Blalock-Taussig shunt versus right ventricle-to-pulmonary artery conduit. *Ann Thorac Surg.* 2007;83:1768-73.
5. Chowdhury UK, Airan B, Sharma R, Bhan A, Kothari SS, Saxena A, et al. One and a half ventricle repair with pulsatile bidirectional Glenn: results and guidelines for patient selection. *Ann Thorac Surg.* 2001;71:1995-2002.
6. Stellina G, Vidaa VL, Milanese B, Rubinoa, Padalinoa, Secchierib S, et al. Surgical treatment of complex cardiac anomalies: the "one and one half ventricle repair." *Eur J Cardiothorac Surg.* 2002;22:431-7.
7. Hosein RBM, Clarke AJB, McGuirk SP, Griselli M, Stumper O, Giovanni JVD, et al. Factors influencing early and late outcome following the Fontan procedure in the current era. The "two commandments"? *Eur J Cardiothorac Surg.* 2007;31:344-53.

doi:10.1016/j.jtcvs.2007.07.074

Hypertension in coarctation

To the Editor:

I read with great interest the article by Hager and associates¹ on coarctation repair. The authors note that the majority of patients remained hypertensive in the long term, even after coarctation repair. This very interesting phenomena, however, has to be examined in the mirror of the little known fact that most patients are often hypertensive *below the level of the coarctation even before the coarctation is repaired!* While pressure measurements indeed show significantly higher pressures above the coarctation, the mean arterial pressure below the coarctation is also higher than that of an average normal subject. Thus, arterial hypertension