idea that, especially in the absence of a measurement of CSF pressure (ie, in those patients without spinal cord drainage), central venous pressure (CVP) can perhaps be used as a substitute, in this context, for CSF pressure. We think it is unlikely that CSF pressure can ever be less than CVP, but the exact relationship between them may not be entirely straightforward despite the observation that CSF and CVP correlate closely during normal cardiopulmonary circulation.^{1,2} The relevance of a high CVP in the context of spinal cord perfusion is explained by the fact that the arterial pressure of the inflow to the spinal cord after segmental artery sacrifice is only a fraction of systemic mean arterial pressure (MAP), as explained next.

We think that establishing an index for assessing the adequacy of the postoperative SCPP is an excellent idea. However, among the parameters readily available in the clinical setting (in the operating room and intensive care unit), we believe the most important variable to adequately assess SCPP is missing: the actual inflow arterial pressure in the collateral system supplying the most vulnerable region of the spinal cord. To obtain this pressure, we have recently adapted a method we developed in our laboratory: we insert a small catheter directly into the stump of a thoracic or lumbar segmental artery during surgery to measure the pressure in the collateral circulation during the operation, and for a variable interval postoperatively when possible.

After extensive segmental artery occlusion, when inflow is provided predominantly by the collateral network, the arterial input for SCPP routinely decreases to levels as low as 20% of MAP, according to our experimental studies.^{3,4} We have confirmed similarly low levels of inflow SCPP (20% of MAP) transiently in patients after segmental artery sacrifice using a catheter placed in L1. In patients with an L1 catheter and CSF drainage, an accurate SCPP can be calculated as $SCPP = L1 \text{ pressure} - CSF \text{ pressure}$.

The acceptable minimum SCPP—the value recorded at the end of operation in the presence of intact function as documented by SSEP/MEP monitoring—could be used as a baseline when calculating an adequate flow for postoperative follow-up: $SCPP \text{ index} = \text{current SCPP} - \text{minimum SCPP}$.

Such an SCPP index could be monitored along with the usual vital signs, and changes in hemodynamics instituted if it de-

creases below zero. A negative SCPP index would indicate that spinal cord perfusion is below safe levels and trigger corrective measures, such as increasing MAP and decreasing CSF or CVP to improve perfusion; the target SCPP index should be zero or more at all times. An additional CONSIDERATION is that the patient may warm significantly after surgery, and increases in metabolic rate in the cord, as well as other tissues supplied by the collateral flow, THAN AT BASELINE. For calculation of A SCPP index in the absence of CSF pressure measurements, CVP could be substituted.

In the absence of direct monitoring, however, we think that routine consideration of AN INDIVIDUAL'S preoperative MAP would be a simplified way for personnel charged with the care of patients after aneurysm surgery which poses a risk of paraplegia to monitor the minimum systemic pressures necessary to provide adequate spinal cord perfusion without their being required to have a sophisticated understanding of spinal cord perfusion requirements. In general, we think that a CVP less than 10 mm Hg would be acceptable, but a high CVP may have to be treated more aggressively if arterial pressure is low. To raise the arterial pressure, we would—in view of the need to keep the CVP low—advocate the use of pharmacologic agents such as epinephrine or norepinephrine rather than using volume infusions.

We thank Dr Augoustides for raising these issues and the editorial board for the opportunity of discussing them.

Christian D. Etz, MD
Randall B. Griepp, MD
Mount Sinai School of Medicine
New York, NY

References

1. Grum DF, Svensson LG. Changes in cerebrospinal fluid pressure and spinal cord perfusion pressure prior to cross-clamping of the thoracic aorta in humans. *J Cardiothorac Vasc Anesth*. 1991;5:331-6.
2. Huynh TT, Miller CC 3rd, Estrera AL, et al. Correlations of cerebrospinal fluid pressure with hemodynamic parameters during thoracoabdominal aortic aneurysm repair. *Ann Vasc Surg*. 2005;19:619-24.
3. Etz CD, Homann TM, Plestis KA, et al. Spinal cord perfusion after extensive segmental artery sacrifice: can paraplegia be prevented? *Eur J Cardiothorac Surg*. 2007;31:643-8.

4. Etz CD, Homann TM, Luehr M, et al. Spinal cord blood flow and ischemic injury after experimental sacrifice of thoracic and abdominal segmental arteries. *Eur J Cardiothorac Surg*. 2008;33:1030-8. Epub 2008 Apr 11.

doi:10.1016/j.jtcvs.2008.05.008

Cardiac stem cells in the real world

To the Editor:

We read with great interest the recent article by Pouly and colleagues.¹ These authors suggest that c-kit⁺ cells retrieved out of human endomyocardial biopsies do not express the classic markers of stemness but are actually mast cells. By immunofluorescent staining on paraffin slices they show that c-kit⁺ cells are CD45⁺ and CD105⁻. Furthermore they found that these c-kit⁺ cells also coexpress tryptase, a granule-derived serine proteinase, and as a consequence appear to be mast cells.

In our experiments, right atrial appendages were removed during routine cardiac surgery and cultured as previously described.² Immunofluorescent staining was performed on cells growing out of the explants after 2 weeks. Primary labeled antibodies directed against c-kit (phycoerythrin), CD45 (fluorescein isothiocyanate [FITC]), and CD105 (FITC) were used. A small percentage of CD45⁺ cells could be detected. These CD45⁺ cells did not stain positive for c-kit (Figure 1, A). In contrast, a subpopulation of CD45⁻ cells expressed c-kit (Figure 1, A). This is in contradiction with the results published by Pouly and colleagues.¹ Double staining with CD105 (FITC) and c-kit (phycoerythrin) revealed that all cells in the outgrowth are CD105⁺ and that a subpopulation of these cells coexpress c-kit (Figure 1, B). These findings refute that c-kit⁺ cells would be mast cells (CD45⁺, tryptase⁺, CD105⁻, and c-kit⁺). Our findings are in accordance with those published by Smith and colleagues² and Bearzi and colleagues.³ Their immunofluorescence and flow cytometric analysis also show that c-kit⁺ cells are CD45⁻ but do express CD105.

Pouly and colleagues¹ state that care has to be taken that phenotypic changes could have occurred by passaging the cells. According to the authors, this could explain why differences in the antigen expressing profile are detected. In our setting, cells were not passaged or processed; immunofluorescence was performed directly on the cell

Figure 1. Immunofluorescent staining on cells grown out of cardiac tissue. Cells incubated with CD45-FITC (A, lower left) and C-kit-phycoerythrin (A, upper right). 4,6-diamino-2-phenylindole (A and B, upper left) was used to stain nuclei. A, Outgrown c-kit⁺ cells do not coexpress CD45 and c-kit. B, Cells are incubated with CD105-FITC (lower left) and c-kit-phycoerythrin (upper right); 95% of the cells do express CD105 and a subpopulation coexpresses c-kit. DAPI, 4,6-diamino-2-phenylindole.

outgrowth. So the difference in the expression profile between our c-kit⁺ cells and that detected by Pouly and colleagues¹ cannot be the result of cell processing.

Our findings indicate that c-kit⁺ cells, present in right atrial appendages, coexpress CD105 but are CD45⁻. It is therefore unlikely that these cells are mast cells because our data indicate that they are probably cardiac progenitor cells.

Remco Koninckx, MSc

Karen Hensen, PhD

Jean-Luc Rummens, MD

Marc Hendriks, MD

Virga Jesse Hospital, Hasselt, Belgium and

University of Hasselt, Diepenbeek, Belgium

References

1. Pouly J, Bruneval P, Mandet C, Proksch S, Peyrard S, Amrein C, et al. Cardiac stem cells in the real world. *J Thorac Cardiovasc Surg*. 2008;135:673-8.
2. Smith R, Brile L, Cho H, Leppo M, Hare J, Messina E, et al. Regenerative potential of cardiosphere-derived cells expanded from percutaneous endomyocardial biopsy specimens. *Circulation*. 2007;115:896-908.
3. Bearzi C, Rota M, Hosoda T, Tillmanns J, Nascimbene A, De Angelis A, et al. Human cardiac stem cells. *Proc Natl Acad Sci U S A*. 2007;104:14068-73.

doi:10.1016/j.jtcvs.2008.05.019

Reply to the Editor:

We thank Koninckx and colleagues for their comments. As stated in our article, data were obtained from both endomyocardial biopsies and atrial appendages, and these 2 sampling sites yielded concordant data. However, the major difference is that we performed in situ detection and characterization of cells, whereas Koninckx and colleagues cultured cells for 2 weeks before immunostaining. Such a time interval can change

the cell phenotype and delete some cell populations that do not survive under these conditions. The latter phenomenon could explain why Koninckx and colleagues did not find any mast cell in their myocardial tissue cultures, whereas it is well established that the myocardium does contain such cells. Because a minor component of the c-kit-positive cells could have represented a subset of cells different from mast cells, we also tested them for other markers of stemness (CD105, islet-1, and MDR1). However, in our hands, these markers remained negative. The data of Koninckx and colleagues suggest that after a period of culture, c-kit-positive cardiac "stem" cells can be identified, but it would be clinically relevant that they provide a quantitative estimate of these cells to assess whether this number allows one to reasonably envision their use for therapeutic purposes.

Julia Pouly, MD

Patrick Bruneval, MD

Philippe Menasche, MD, PhD

Hopital Europeen Georges Pompidou

Department of Cardiovascular Surgery

Paris, France

doi:10.1016/j.jtcvs.2008.05.020

Aprotinin; An economy of truth? To the Editor:

I write to comment on the editorials by Drs Sundt¹ (April 2008) and Westaby² (March 2008). Dr Sundt is perceptive but fails to consider that the observational studies suggesting a danger with aprotinin may have had bias in the analysis.

At the advisory committee meeting of the Food and Drug Administration (FDA) held on September 12, 2007, Dr Mangano allowed the FDA access to the McSPI (Multicenter Study of Perioperative Ischemia) data set. Whereas Mangano and col-

leagues³ used a propensity score that was based on likelihood of bleeding, the FDA reanalysis of these data used stratification according to risk of adverse outcome. The FDA analysis showed no increases in relative risks (RRs) for death (RR 0.91, 95% confidence interval [CI] 0.54–1.53), heart failure (RR 1.05, 95% CI 0.75–1.47), myocardial infarction (RR 1.10, 95% CI 0.88–1.39), or renal dysfunction (RR 1.26, 95% CI 0.76–2.11) when data from 1222 aprotinin-treated patients were compared with those of 1307 patients who did not receive the drug.⁴

At the same FDA meeting, Dr Karkouti, who used matching of pairs of data, showed that the inclusion into the model of cardiopulmonary bypass variables (time and circulatory arrest) and transfusion (>4 units of red blood cells and fresh-frozen plasma) removed the statistical effects of aprotinin on renal function.⁴ The Toronto data have never shown any other mortality or morbidity risk. Dr Funary also presented the North West Consortium analysis, which showed that any apparent effect of aprotinin on adverse outcome is lost when red blood cell transfusion numbers are included as a confounding variable.⁵

In the article from Shaw and colleagues⁶ of the Duke University Medical Center, the populations of patients receiving aprotinin or ε-aminocaproic acid (EACA) were hugely different. No matter how clever the statistical modeling, clinicians will recognize that there must be differences in management and outcome between a patient with isolated myocardial ischemia undergoing primary, elective revascularization (given EACA) and one undergoing a nonelective reoperation for heart failure associated with valve pathology (who would likely receive aprotinin in about 70%-80% of cases worldwide). Despite this, Shaw and colleagues⁶