describing their experience of a transcranial Doppler system (Embo-Dop; DWL Elektronische Systeme GmbH, Singen, Germany) to discriminate between gaseous and particulate emboli after aortic valve replacement. The main finding of their study was that solid microemboli occurred more frequently after mechanical aortic valve replacement compared with biological valve replacement at 3 months after surgery. The inability to distinguish the difference between solid and gaseous microemboli has limited the use of standard transcranial Doppler when both coexist, for example, during open carotid endarterectomy. As the authors state, this particular Doppler technique was first assessed by Brucher and Russell² in a 2-part study comprising both in vitro and in vivo models. The in vivo part used 45 patients with carotid artery stenosis (39 symptomatic), producing a relatively low number of 32 embolic signals, of which 30 were classified as solid.

The assumption in validation studies is that all patients with carotid artery stenosis produce only solid emboli. One arm of a clinical validation study performed by Markus and Punter³ used a cohort of patients with symptomatic carotid artery stenosis to generate 145 presumed solid embolic signals. Of these 145 signals, the Embo-Dop system classified 72 as gaseous emboli. This gave a sensitivity of 50.3% and specificity of 96.5% for detecting solid embolic signals. Markus and Punter concluded that although the Embo-Dop system allowed better discrimination than simple intensity threshold alone, it was not accurate enough for clinical or research studies. Unless the specificity is approximately 100%, a significant number will be classified wrongly as potentially significant solid emboli. We believe that Embo-Dop has to be used with caution in distinguishing gaseous from solid emboli.

The authors comment on no antiplatelet therapy after mechanical versus biological valve replacement. Our

group conducted a UK survey of consultant practice and literature review of this subject.⁴ We concluded that the incidence of thromboembolism after aortic bioprosthesis is highest in the first 3 months but then diminishes. Whether this is related to endothelialization of prosthetic material or specifically to antiplatelet therapy is unknown. Therefore, their observation that the proportion of particulate emboli at 3 months was in the biological valve replacement cohort receiving aspirin but not in the mechanical group receiving only warfarin could indeed be explained by the antiplatelet function of aspirin.

Justin Nowell, MRCS
Marjan Jahangiri, FRCS (CTh)
St George's Hospital
London, United Kingdom

References

1. Guerrieri Wolf L, Choudhary BP, Abu-Omar Y, Taggart DP. Solid and gaseous cerebral microembolization after biologic and mechanical aortic valve replacement: investigation with multirange and multifrequency transcranial Doppler ultrasound. *J Thorac Cardiovasc Surg.* 2008;135:512-20.
2. Brucher R, Russell D. Automatic online embolus detection and artifact rejection with the first multifrequency transcranial Doppler. *Stroke.* 2002;33:1969-74.
3. Markus HS, Punter M. Can transcranial Doppler discriminate between solid and gaseous microemboli? Assessment of a dual-frequency transducer system. *Stroke.* 2005;36:1731-4.
4. Nowell J, Wilton E, Markus H, Jahangiri M. Antithrombotic therapy following bioprosthetic aortic valve replacement. *Eur J Cardiothorac Surg.* 2007;31:578-85.

doi:10.1016/j.jtcvs.2008.05.065

Reply to the Editor:

We thank Nowell and Jahangiri for their interest in our recent article¹ using a novel transcranial Doppler system (Embo-Dop; DWL Elektronische Systeme GmbH, Singen, Germany) to investigate the number and nature of cerebral microemboli in recipients of biological and mechanical aortic valve prostheses. The 2 important issues highlighted by the authors in their

Letter to the Editor were, in fact, both discussed specifically in our article.

The first issue is the potentially low sensitivity and specificity of the Embo-Dop system as suggested in the clinical validation study by Markus and Punter.² With specific regard to that study, we wrote in the discussion section of our article: "More recently, Markus and Punter, while confirming the superior discrimination achievable with this new Embo-Dop transcranial Doppler system compared with conventional transcranial Doppler ultrasound, reported it to have a sensitivity of only 50% but a specificity of 96% for detecting solid microemboli in vivo. If confirmed, that finding will only serve to emphasize the already significantly greater solid microembolic load detected with mechanical rather than biological prostheses." In other words, because the sensitivity of the Embo-Dop system is much lower than its specificity in detecting solid microemboli, this will result in an underestimation of the solid cerebral microembolic load.

The second issue raised by Nowell and Jahangiri is that the reduction of solid microemboli over time in the biological prosthesis recipients, but not in the mechanical prosthesis recipients, could be explained by the use of aspirin in the first group. Again, we discussed that issue in our article stating, "Although biological recipients were all receiving aspirin, mechanical recipients received warfarin without any antiplatelet agents. This could explain the reduction in the proportion of particulate microemboli over time in the biological valve group but not in the mechanical group." We subsequently suggested, "However, the results could also be affected by the use of aspirin in the biologic valve group, which would explain the reduction of microemboli detected over time in group B (biological) but not in group M (mechanical). Indeed, using aspirin instead of warfarin after a biological aortic valve implantation is the current practice among the

majority of cardiac surgeons, and consequently our findings reflect a 'real-world' situation."

Lorenzo Guerrieri Wolf, MD

Bikram P. Choudhary, MRCS

Yasir Abu-Omer, MRCS

David P. Taggart, MD (Hons), PhD,

FRCS

Department of Cardiothoracic Surgery

John Radcliffe Hospital

Oxford, United Kingdom

References

1. Guerrieri Wolf L, Choudhary BP, Abu-Omar Y, Taggart DP. Solid and gaseous cerebral microembolization after biological and mechanical aortic valve replacement: investigation with multirange and multifrequency transcranial Doppler ultrasound. *J Thorac Cardiovasc Surg.* 2008;135:512-20.
2. Markus HS, Punter M. Can transcranial Doppler discriminate between solid and gaseous microemboli? Assessment of a dual-frequency transducer system. *Stroke.* 2005;36:1731-4.

doi:10.1016/j.jtcvs.2008.07.003

CONTEGRA AND EXTRACARDIAC FONTAN

To the Editor:

We read with the greatest interest the article entitled "Feasibility of the Extracardiac Conduit Fontan Procedure in Patients Weighing Less than 10 Kilograms."¹ After congratulating the authors for their excellent work, we have a few comments.

As a Fontan Conduit, they used polytetrafluoroethylene (Gore-Tex; WL Gore & Associates Inc, Flagstaff, Ariz) (14–20 mm in diameter) King Faisal Specialist Hospital and Research Center in Jeddah, used Contegra valved conduits in 4 patients for inferior vena cava–right pulmonary artery connection (Fontan). They speculated that an additional benefit of using the Contegra xenograft could be that it provides a competent valve in the Fontan circulation, which might help by reducing the reversal of flow and maintaining better forward flow into the pulmonary circulation. The concern was whether the valve, in a nonpulsatile, passive Fontan circulation, would facilitate smooth antegrade flow into the pulmonary circula-

tion or act as a source of resistance to the flow. The findings that none of their patients had hepatomegaly or ascites on physical examination and that the post-operative echocardiograms showed excellent antegrade flow in the inferior vena cava, which also was not dilated, and minimal flow reversal in the hepatic veins support the facilitation of smooth forward flow. They presented a new option for constructing an extracardiac Fontan connection with the Contegra xenograft that has not been previously published. Their early echocardiographic results supported the idea of minimizing the reversal of infradiaphragmatic venous flow, decreasing hepatic congestion, and possibly reducing the development of ascites.² They do have a way of assessing the conduit size by using Rowlatt and colleagues' chart.³

The authors routinely administered 5 mg/kg ticlopidine as antiplatelet therapy. Warfarin was not used even though they had a patient with arteriovenous valve replacement. How did they control the thrombogenic potential here?

Measurement of the lower lobe index is an important adjunct to evaluate candidates for the Fontan operation. The predictive value of the Nakata index is less informative than the lower lobe index, which was introduced to optimize the preoperative selection of Fontan candidates. Lower lobe index may be more predictive to evaluate the adequacy of the pulmonary vascular tree, given that central pulmonary arteries used to calculate the Nakata index can be enlarged before or during the Fontan operation.⁴

Sameh Ibrahim Sersar, MD

Ahmed A. Jamjoom, MD

Division of Cardiothoracic Surgery

Department of Cardiovascular

Diseases

King Faisal Specialist Hospital and

Research Center

Jeddah, Saudi Arabia

References

1. Ikai A, Fujimoto Y, Hirose K, Ota N, Tosaka Y, Nakata T, et al. Feasibility of the extracardiac conduit Fontan procedure in patients weighing less

than 10 kilograms. *J Thorac Cardiovasc Surg.* 2008;135:1145-52.

2. Baslaim G, Hussain A, Kouatli A, Jamjoom A. Bovine valved xenograft conduits in the extracardiac Fontan procedure. *J Thorac Cardiovasc Surg.* 2003;126:586-8.
3. Rowlatt JF, Rimaldi JMA, Lev M. The quantitative anatomy of the normal child's heart. *Pediatr Clin North Am.* 1963;10:499-588.
4. Ovrutski S, Alexi-Meskishvili V. Does the Nakata index predict outcome after Fontan operation? *Eur J Cardiothorac Surg.* 2008;33:951.

doi:10.1016/j.jtcvs.2008.05.067

Reply to the Editor:

We appreciate Dr Sersar's comments on our recent article, "Feasibility of the Extracardiac Conduit Fontan Procedure in Patients Weighing Less than 10 Kilograms." These comments raise several questions concerning the following: (1) choice of conduit type, (2) anticoagulation therapy after the Fontan procedure, and (3) examination of the pulmonary arteries for evaluating the pulmonary vascular tree.

In our article, we describe our strategy for the extracardiac Fontan operation without fenestration. Our strategy aimed to eliminate cyanosis as soon as possible. Fenestration might reduce the duration of pleural drainage but cannot eliminate cyanosis even after the Fontan procedure. We mentioned in the article that peritoneal drainage might be able to reduce the duration of pleural drainage in small children in particular. In addition to the obvious purpose of the drainage of ascites itself, drainage of ascites also results in the drainage of the inflammatory factors present in the ascites.¹ In our series at least, there were no patients with recurrent ascites after the cessation of the initial peritoneal drainage. We have a chance to use a Contegra conduit² if this conduit is an attractive option for reducing ascites and pleural effusion. However, the use of this conduit for the Fontan operation is not permitted in Japan. The fate of the conduit as a xenograft, especially in terms of obstruction of conduit and durability of valve function, over the long term should be investigated.