documented in the literature in the context of simple transposition of the great vessels but was not discussed further in this study. The arterial switch component of the double-switch procedure carries all the known risk hazards as in any arterial switch and long-term complications, including fatal arrhythmias, neo-aortic regurgitation, and left ventricular dysfunction.³ These risks can be higher among those with a single orifice coronary artery.

Training is a relative term and is tailored to the individual patient. Excessive training leading to LV pressures more than 75% systemic is a well-known association with late ventricular dysfunction. Although the study states that LV pressures were 65% to 80%, it would be more informative to the reader if the authors explained the target pressures for the 24 patients after PA banding in the nontrained subgroup. Furthermore, approximately 46% (n = 5) of the trained subgroup had more than 1 PA band to properly prepare the morphologic LV (mLV) for the double switch. It would be important to know whether such patients' ventricles fared better than the application of singleton pressure load PA banding, especially when not all LV training is detrimental in older patients.

It is interesting to note that the mLV/mRV pressure ratio was similar in the trained and nontrained groups at the time of their anatomic repair. However, those older patients with LV dysfunction achieved mLV/mRV pressure in less time (436 vs 730 days) than those without dysfunction. This may imply that the LV was generating the same pressure at the time of the double switch as in those with ventricular defect physiology (increased end-diastolic volume, reduced afterload) but fared better than in the trained group. This may illustrate that the PA band is a confounder, whereas the morphology (congenitally corrected transposition of the great arteries with small restrictive ventricular septal defect or no ventricular septal defect) is an independent predictor, especially when the authors state in

Table 1 that 20% of the nontrained group had good mLV versus 60% in the trained group ($P = .04$, statistically significant). Furthermore, in support of this is the well-known association of worsening survival in the older infant with transposition of the great arteries and an intact ventricular septum after the arterial switch compared with transposition of the great arteries and a ventricular septal defect.⁴ The additional benefit of ventricle volume load via a systemic circulation to PA shunt to the pressure load PA band afterload has been described and could be of benefit in corrected transposition.⁵ Finally, the ventricular activation time period (QRS) has been a strong marker of ventricular function.⁶ The authors did not state its respective duration for the 2 groups before and after the double switch. I would appreciate the authors' response based on their vast experience when managing this unique group of patients.

Jeffrey Shuhaiber, MD

Loyola University Stritch School of
Medicine
Maywood, Ill

References

1. Quinn DW, McGuirk SP, Metha C, Nightingale P, de Giovanni JV, Dhillon R, et al. The morphologic left ventricle that requires training by means of pulmonary artery banding before the double-switch procedure for congenitally corrected transposition of the great arteries is at risk of late dysfunction. *J Thorac Cardiovasc Surg.* 2008;135:1137-44.
2. Chiu IS, Wu SJ, Chen SJ, Wang JK, Wu MH, Lue HC. Sequential diagnosis of coronary arterial anatomy in congenitally corrected transposition of the great arteries. *Ann Thorac Surg.* 2003;75:422-9.
3. Angeli E, Raisky O, Bonnet D, Sidi D, Vouhé PR. Late reoperations after neonatal arterial switch operation for transposition of the great arteries. *Eur J Cardiothorac Surg.* 2008;34:32-6. Epub 2008 May 8.
4. Prêtre R, Tamisier D, Bonhoeffer P, Mauriat P, Pouard P, Sidi D, et al. Results of the arterial switch operation in neonates with transposed great arteries. *Lancet.* 2001;357:1826-30.
5. Ilbawi MN, Idriss FS, DeLeon SY, Muster AJ, Gidding SS, Duffy CE, et al. Preparation of the left ventricle for anatomical correction in patients with simple transposition of the great arteries. Surgical guidelines. *J Thorac Cardiovasc Surg.* 1987;94:87-94.
6. Bautista-Hernandez V, Marx GR, Gauvreau K, Mayer JE Jr, Cecchin F, del Nido PJ. Determinants of left ventricular dysfunction after anatomic repair of congenitally corrected transposition of the great arteries. *Ann Thorac Surg.* 2006;82:2059-65.

doi:10.1016/j.jtcvs.2008.07.076

CONTEGRA AND EXTRACARDIAC FONTAN

To the Editor:

We read with great interest the letter to the editor by Sersar and Jamjoom¹ in the *Journal* of November 2008, describing the use of the Contegra (Medtronic Inc, Minneapolis, MN) bovine jugular vein as a possible alternative extracardiac conduit in total cavopulmonary connection (TCPC).

We want to emphasize our bad experience with this particular conduit in patients undergoing the Fontan procedure, which was published in the *Journal* 7 years ago.²

The only 3 patients in whom the Contegra conduit was used for TCPC all experienced serious complications of conduit thrombosis; 2 of them died and 1 survived, but with major neurologic injury. The Contegra implantation was carried out at different institutions by different surgeons.

Evidence has meanwhile accumulated that thrombogenicity is indeed an inherent characteristic of the Contegra bovine jugular venous conduit. Thrombus formation has been shown to occur even in high-flow right ventricular to pulmonary artery conditions.^{3,4}

A fatal case of conduit thrombosis also occurred at the authors' institution. This was published by Baslaim⁵ in 2008 (1 of 18 patients undergoing operation between 2002 and 2005). An "operative technical cause" is suggested by the author as a possible cause.

We are looking forward to the structured clinical and echocardiographic follow-up in their patients. For the present, we continue to discourage the use of the Contegra bovine jugular vein as an extracardiac conduit in TCPC.

Paul Schoof, MD, PhD

Tjalling Waterbolk, MD

Divisions of Cardiothoracic Surgery
University Medical Center Nijmegen
and Groningen
The Netherlands

References

1. Sersar SI, Jamjoom AA. Contegra and extracardiac Fontan. *J Thorac Surg.* 2008;136:1393.

2. Schoof PH, Koch AD, Hazekamp MG, Waterbolk TW, Ebels T, Dion RA. Bovine jugular vein thrombosis in the Fontan circulation. *J Thorac Cardiovasc Surg.* 2002;124:1038-40.
3. Tiete AR, Sachweh S, Roemer U, Kozlik-Fredmann R, Reuchart B, Daebritz SH. Right ventricular outflow tract reconstruction with the Contegra bovine jugular vein conduit: a word of caution. *Ann Thorac Surg.* 2004;77:2151-6.
4. Shebani SO, McGuirk S, Baghai M, Stickley J, DeGiovanni JV, Bu'Lock F, et al. Right ventricular outflow tract reconstruction using Contegra valved conduit: natural history and conduit performance under pressure. *Eur J Cardiothorac Surg.* 2006;29:397-405.
5. Baslaim G. Bovine valved xenograft (Contegra) conduit in the extracardiac Fontan procedure: the preliminary experience. *J Card Surg.* 2008;23:146-9.

doi:10.1016/j.jtcvs.2009.02.051

LATE RUPTURE OF EXPANDED POLYTETRAFLUORO-ETHYLENE NEOCHORDAE USED FOR MITRAL VALVE REPAIR

To the Editor:

In the February issue of the *Journal*, Farivar and colleagues¹ described a case of late rupture of expanded polytetrafluoroethylene (ePTFE) neochordae after mitral valve repair. In the "Discussion," they state that there was only 1 previous case reported in the literature of ruptured neochordae, presented by Buttany and colleagues,² suggesting their case as the second reported. This is not correct. In 2007, the *Journal of Heart Valve Disease* published our report of 2 cases of acute mitral regurgitation caused by ruptured ePTFE neochordae.³ Our patients presented with symptoms of acute heart failure caused by sudden rupture of the chordae, which was different from the rare clinical manifestations, hemolysis, and hematuria experienced by the patient reported by Farivar and colleagues.

We first used ePTFE routinely to correct ruptured or elongated chordae tendineae during the late 1980s and have since implanted it in more than 500 patients without a single known case of ePTFE failure necessitating reoperation, until these 2 late events occurred. After a thorough literature

search regarding the possible mechanisms behind this unusual occurrence, we analyzed our histopathologic findings. We found that there was likely pannus formation and collagen infiltration through the porous structure that could account for progressive disruption before calcification ensued, which could be postulated as a form of repair/scarring.

In addition to the degenerative causes that could be responsible for the late rupture, we believe that the use of forceps and other metal instruments in the manipulation of the ePTFE could interrupt its electronegative charge and weaken the material. We currently avoid any kind of metallic manipulation.

Gonçalo Coutinho, MD

Lina Carvalho, MD, PhD

Manuel J. Antunes, MD, PhD, DSc

Center of Cardiothoracic Surgery

University Hospital

Coimbra, Portugal

References

1. Farivar RS, Sherman SK, Cohn LH. Late rupture of polytetrafluoroethylene neochordae after mitral valve repair. *J Thorac Cardiovasc Surg.* 2009;137:504-6.
2. Buttany J, Collins MJ, David TE. Ruptured synthetic expanded polytetrafluoroethylene chordae tendineae. *Cardiovasc Pathol.* 2004;13:182-4.
3. Coutinho GF, Carvalho L, Antunes MJ. Acute mitral regurgitation due to ruptured ePTFE neo-chordae. *J Heart Valve Dis.* 2007;16:278-81.

doi:10.1016/j.jtcvs.2009.02.050

BIVENTRICULAR PACING AFTER CARDIAC SURGERY

To the Editor:

We read with interest the article by Evonich and colleagues¹ in the October issue of the *Journal*. They performed a randomized study of pacing modes ("normal care," synchronous atrial-right ventricular pacing, and synchronous atrial-biventricular pacing) in 40 patients with impaired left ventricular function after cardiac surgery. They included both a crossover study of acute hemodynamic variables

early after operation (in 29 patients) and an assessment of clinical end points, such as intensive therapy unit and hospital stay and mortality. In contrast with experience in chronic heart failure, they showed no advantage to biventricular pacing in any area, although the trial was underpowered for the clinical end points.

They suggest this was the first randomized study in surgical patients but did not cite our 2005 publication.² We performed a similar randomized crossover study, with hemodynamic assessment of different pacing modes in a similar number of patients (25) early after surgery. We demonstrated a clear advantage to atrial-left ventricular pacing and regard this modality as an important addition to the postoperative care of the compromised patient.

Closer analysis of the results in the article by Evonich and colleagues¹ suggests that there is a serious error or that they too found an advantage but did not recognize it. In the second paragraph of the section titled "Hemodynamic Testing," they state a statistically significant difference in stroke volume, cardiac output, and left ventricular stroke work index between AAI and atrial-right ventricular pacing. But, in Table 2, the values they refer to in the text are in the column under atrial-biventricular pacing. If Table 2 is to be believed (and there is inconsistency in nomenclature, does AAI equate to "usual care"?), these investigators *did* see an advantage to biventricular pacing.

We believe biventricular pacing is of advantage to the postsurgical patient. Even if we are mistaken in our interpretation of this article, surgeons are left with 2 series of patients with contrasting results, perhaps because both have small numbers. This calls for a larger definitive study.

Michael Flynn, FRCS

John H. Dark, FRCS

Janet M. McComb, FRCP, MD

University of Newcastle upon Tyne

United Kingdom