Orthotopic vascularized right lung transplantation in the mouse

Wenjun Li, MD,^a Seiichiro Sugimoto, MD, PhD,^a Jiaming Lai, MD,^a G. Alexander Patterson, MD,^a Andrew E. Gelman, PhD,^{a,b} Alexander S. Krupnick, MD,^a and Daniel Kreisel, MD, PhD^{a,b}

Objectives: Orthotopic left lung transplantation in the mouse, as recently developed by our laboratory, represents a physiologic model for studies in transplantation biology. However, because of the limited overall respiratory contribution of the murine left lung, left lung transplant recipients remain healthy despite immune-mediated graft necrosis. We sought to develop a lung transplantation model in which animal survival depends on graft function.

Methods: Orthotopic vascularized right lung transplantations using cuff techniques were performed in syngeneic and allogeneic strain combinations. Grafts were assessed histologically or functionally by measuring arterial blood gases from 7 to 28 days after transplantation. In a parallel set of experiments, syngeneic and immunosuppressed allogeneic hosts underwent a left pneumonectomy 2 weeks after right lung transplantation, with assessment of graft function 1 week later.

Results: We performed 40 right lung transplantations, with a survival rate of 87.5%. Syngeneic grafts remain free of inflammation as far as 28 days after transplantation. On day 7, arterial oxygen levels in syngeneic recipients (481 ± 90 mm Hg) are equivalent to those in naive mice (503 ± 59 mm Hg) after left hilar occlusion. Alternatively, allogeneic grafts develop histologic evidence of acute rejection, and arterial oxygen levels are significantly decreased after left hilar clamping (53.3 ± 10.3 mm Hg). Both syngeneic and immunosuppressed allogeneic right lung recipients tolerate a left pneumonectomy.

Conclusions: Right lung transplantation followed by left pneumonectomy represents the first survival model of vascularized lung transplantation in the mouse and will therefore allow for the design of novel studies in experimental lung transplantation. (*J Thorac Cardiovasc Surg* 2010;139:1637-43)

Physiologic animal models have been instrumental for the advancement of clinical lung transplantation. To this end, technical refinements and immunosuppressive studies in canine models of pulmonary autotransplantation or allotransplantation have paved the way for the first successful human lung transplantations.¹⁻³ Introduction of cuff techniques has allowed for the widespread use of orthotopic lung transplantation in the rat.⁴ Undoubtedly, important insights into lung transplantation biology have been gained by using the rat model. However, the paucity of transgenic and knockout strains in the rat limits the use of this model for the design of mechanistic studies. Extending microsurgical techniques established in the rat model, our laboratory was the first to describe a method for orthotopic vascularized left lung transplantation in the mouse.⁵ Using this model, we

have reported important differences between requirements for allograft rejection in the lung compared with other organs.⁶ Notably, there exist marked similarities between observations in murine and human lung allografts, pointing out the potential translational value of the murine model.^{5,7}

Other murine transplantation models have clearly defined indicators of graft rejection, such as cessation of heart beat in the case of vascularized cardiac transplantation, animal death in the case of kidney transplantation with removal of the recipient's native kidneys, or sloughing and necrosis of the graft in the case of skin transplantation.⁸⁻¹⁰ One limitation of the orthotopic left lung transplantation model in the mouse is that recipient animals survive and appear healthy even after alloimmune-mediated necrosis of their allografts.¹¹ Therefore graft assessment relies on histologic examination of the transplanted left lung. We sought to develop a murine lung transplantation model in which survival depends on the function of the transplanted lung. Here we took advantage of the fact that the right murine lung is markedly larger than the left lung. The right murine lung, which is composed of 4 lobes, accounts for approximately 70% of the total lung volume.¹² Here, for the first time, we report the development of a technique for orthotopic vascularized right lung transplantation in the mouse. Syngeneic recipients have fully functional grafts and are able to survive a left pneumonectomy, thereby providing the first description of a survival model for experimental lung transplantation in the mouse.

From the Departments of Surgery^a and Pathology and Immunology,^b Washington University, St Louis, Mo.

Dr Kreisel is supported by a grant jointly sponsored by the National Heart, Lung, and Blood Institute and the Thoracic Surgery Foundation for Research and Education (1K08HL083983) and by a grant from the National Heart, Lung, and Blood Institute (1R01HL094601).

Disclosures: None.

Received for publication July 28, 2009; revisions received Oct 15, 2009; accepted for publication Nov 12, 2009; available ahead of print March 10, 2010.

Address for reprints: Daniel Kreisel, MD, PhD, Departments of Surgery and Pathology and Immunology, Campus Box 8234, 660 South Euclid Ave, Washington University in St Louis, St Louis, MO 63110-1013 (E-mail: kreiseld@wudosis.wustl.edu). 0022-5223/\$36.00

Copyright © 2010 by The American Association for Thoracic Surgery
doi:10.1016/j.jtcvs.2009.11.019

MATERIALS AND METHODS

Animals

Male inbred C57BL/6 (H-2b; B6) and BALB/c (H-2d) mice were purchased from Jackson Laboratories (Bar Harbor, Me). Animals weighing 22 to 28 grams were used as donors, and mice weighing 26 to 30 grams were used as recipients. Syngeneic transplantations were performed in the B6 → B6 strain, and allogeneic transplantations were performed in the B6 → BALB/c and BALB/c → B6 strain combinations. Some allogeneic transplants received treatment with anti-CD40 ligand antibody (MR1; 250 μ g administered intraperitoneally, day 0) and cytotoxic T-lymphocyte-associated antigen-4-immunoglobulin (200 μ g administered intraperitoneally, day 2). All animal procedures were approved by the Animal Studies Committee at Washington University School of Medicine. Animals received humane care in compliance with the "Guide for the care and use of laboratory animals" prepared by the National Academy of Sciences and published by the National Institutes of Health and the "Principles of laboratory animal care" formulated by the National Society for Medical Research.

Donor Procedure

Donor mice are anesthetized with an intraperitoneal injection of ketamine (50 mg/kg) and xylazine (10 mg/kg), intubated with a 20-gauge angiocatheter through a tracheostomy, and then attached to a rodent ventilator. After injection of 100 units of heparin into the penile vein, the chest is exposed through a median laparosternotomy. The right atrium, left atrium, and inferior vena cava are incised to vent the heart, and subsequently, 3 mL of cold low-potassium dextran glucose solution is injected into the main pulmonary artery to flush the lungs. The trachea is ligated after the lungs are inflated to end-tidal volume with room air, and the heart–lung block is excised. The right lung is then prepared for the placement of cuffs. A 22-gauge angiocatheter is used for the pulmonary artery, a 20-gauge angiocatheter is used for pulmonary veins, and an 18-gauge angiocatheter is used for the right mainstem bronchus.

Because of anatomic differences between the right and the left lungs in the mouse, we were forced to modify our previously described technique for the donor portion of the left lung transplantation.⁵ The pulmonary vein is located anterior to the pulmonary artery, and the right mainstem bronchus courses posterior to the pulmonary artery. The anatomic arrangement of these hilar structures necessitates a careful dissection to avoid damage. Importantly, although the left lung is made up of a single lobe, the right lung is composed of 4 lobes, namely the apical lobe, the azygous lobe, the cardiac lobe, and the diaphragmatic lobe (Figure 1, A). The pulmonary veins draining the lobes of the right lung converge in close proximity to the left atrium, resulting in a relatively short right pulmonary vein. Therefore the length of the cuff for the right pulmonary vein (0.3 mm excluding the extension handle) needs to be shorter than for left lung transplantation (0.6 mm; Figure 1, B). The length of the cuff for the pulmonary artery is 0.6 mm. After introduction of these structures through the cuffs and evert their ends, the cuffs are secured with a circumferential 10–0 nylon suture. Of note, as previously described for the left lung transplantation procedure, we cuff the bronchus at the time of implantation. The length of the bronchial cuff is 0.6 mm. The donor procedure takes approximately 40 minutes. The donor lung is stored in low-potassium dextran glucose solution at 4 °C until implantation.

Recipient Procedure

Recipient mice are anesthetized with an intraperitoneal injection of ketamine (50 mg/kg) and xylazine (10 mg/kg), intubated orotracheally with a 20-gauge intravenous catheter, and then placed in a left lateral decubitus position. A right thoracotomy is performed, a portion of the third rib is excised to enhance exposure, and a chest retractor is placed. Dissection is initiated in the superior portion of the pulmonary hilum, where the right mainstem bronchus is gently freed from the pulmonary artery. Of note, the right mainstem bronchus is markedly shorter than the left mainstem

FIGURE 1. A, Right lung anatomy: 1, apical lobe; 2, azygous lobe; 3, cardiac lobe; 4, diaphragmatic lobe. B, Donor pulmonary vein (PV) after placement of the cuff. C, Implantation of donor right lung is initiated by insertion of the cuffed donor pulmonary artery (PA) into the recipient PA. Only the inferior branch is used for later insertion of the cuffed donor PV. Br, Bronchus.

bronchus. Therefore during dissection of the right mainstem bronchus, special attention needs to be directed toward avoiding injury to the left bronchus. The right bronchus is occluded close to the carina with a slipknot (8–0 silk ligature), leading to a collapse of the right lung, which facilitates the subsequent dissection. To further improve exposure, we then excise the diaphragmatic lobe and attach a clamp to the remainder of the right lung for lateral retraction. The pulmonary artery is dissected free from the pulmonary vein and then occluded with a slipknot (10–0 silk ligature). The superior branch of the pulmonary vein is divided between two 10–0 silk ties. We then occlude the inferior branch of the pulmonary vein with a slipknot (8–0 silk ligature) close to its confluence with the superior branch. The recipient pulmonary artery is incised, and the cuffed donor pulmonary artery is inserted and secured as previously described for the left lung transplantation procedure (Figure 1, C).¹³ The donor lung is rotated medially to expose the recipient right mainstem bronchus. We then incise the recipient bronchus, insert the cuffed donor bronchus, and secure it with a circumferential ligature (10–0 nylon). Only the inferior branch of the recipient pulmonary vein is used for insertion of the cuffed donor pulmonary vein. The pulmonary venous anastomosis is also secured with a circumferential ligature (10–0 nylon). The recipient right lung is excised, and the slipknots on the pulmonary veins, pulmonary artery, and bronchus are released to reperfuse the lung. The recipient procedure takes approximately 70 to 80 minutes. The cold ischemic period lasts approximately 50 minutes, and the implantation of the donor lung, during which the graft is covered with an ice cold sponge, takes approximately 15 minutes.

Left Pneumonectomy

Two weeks after right lung transplantation, mice are placed in a right lateral decubitus position and a left thoracotomy is performed. The chest is entered through the third intercostal space. The left hilar structures are mass ligated with a 6–0 silk tie, and the left lung is excised. A piece of Gelfoam (Pfizer, New York, NY) is placed into the left chest cavity before closure to stabilize the mediastinum. Postoperatively, mice are maintained on 100% oxygen for 30 minutes.

Histology

At the time of death, lung grafts were harvested, inflation fixed in formaldehyde, embedded in paraffin, sectioned, and stained with hematoxylin and eosin.

Functional Graft Assessment

After a 4-minute period of occlusion of the left pulmonary hilum, arterial blood was drawn from the left ventricle while mice were ventilated with a fraction of inspired oxygen of 1.0. Blood gases were measured with an iSTAT Portable Clinical Analyzer (iMale STAT Corp, East Windsor, NJ).

Statistical Analysis

Statistical analysis was performed with the Student's *t* test. Oxygen levels are expressed as the mean $\pm$ standard error of the mean.

RESULTS

Macroscopic and Microscopic Appearance of Syngeneic and Allogeneic Grafts

A total of 40 right lung transplantations were performed, with a survival rate of 87.5%. Five animals died intraoperatively of bleeding ($n = 3$) or in the immediate postoperative period because of pneumothoraces ($n = 2$). Syngeneic B6 grafts were ventilated at 7 days ($n = 10$) and appeared grossly normal up to 28 days ($n = 16$) after transplantation (Figure 2). On histologic examination, these grafts had no

evidence of inflammation and were comparable with lungs not undergoing surgical intervention. On the other hand, allogeneic BALB/c $\rightarrow$ B6 right lung grafts were not ventilated at 7 days after transplantation and had histologic evidence of acute rejection with perivascular cuffing and peribronchiolar mononuclear infiltrates ($n = 3$, Figure 3). B6 $\rightarrow$ BALB/c ($n = 3$) right lung grafts were also acutely rejected 7 days after transplantation (data not shown).

Functional Assessment of Right Lung Grafts

To assess the grafts functionally, we compared arterial oxygen levels in naive B6 mice and syngeneic B6 $\rightarrow$ B6 right lung transplant recipients 7 days after engraftment after occlusion of the left pulmonary hilum. We found no significant differences between oxygen levels in these 2 groups after either 4 minutes (PaO_2 of 503 ± 59 mm Hg in naive B6 mice [$n = 4$] and PaO_2 of 481 ± 90 mm Hg in B6 $\rightarrow$ B6 right lung transplant recipients [$n = 4$], $P > .1$) or 10 minutes of left hilar occlusion (PaO_2 of 399.2 ± 16.9 mm Hg in naive B6 mice [$n = 3$] and PaO_2 of 362.5 ± 6.9 mm Hg in B6 $\rightarrow$ B6 right lung recipients [$n = 3$], $P > .1$), indicating that the syngeneic right lung transplants are functional (Figure 4). However, oxygen levels were significantly lower in allogeneic BALB/c $\rightarrow$ B6 right lung recipients compared with those in both naive B6 mice and syngeneic right lung recipients after 4 minutes of left hilar clamping (53.3 ± 10.3 mm Hg, $n = 3$; $P < .01$ compared with syngeneic transplant recipients and naive mice).

Right Lung Transplantation Followed by Left Pneumonectomy

We measured arterial oxygen levels in syngeneic B6 right lung transplant recipients 7 days after a left pneumonectomy. There were no statistically significant differences between these mice and naive B6 mice 7 days after a left pneumonectomy (PaO_2 of 534 ± 61 mm Hg in naive B6 mice after left pneumonectomy [$n = 3$] and PaO_2 of 461 ± 67 mm Hg in B6 $\rightarrow$ B6 right lung transplant recipients after left pneumonectomy [$n = 3$], $P = 0.23$; Figure 5). We have previously reported that in left lung transplants perioperative blockade of CD40/CD40 ligand and B7/CD28 costimulatory pathways prevents graft necrosis, and grafts stay ventilated for time periods exceeding 100 days.¹¹ BALB/c $\rightarrow$ B6 right lung transplant recipients that received this regimen tolerated a left pneumonectomy. Their arterial oxygen levels 7 days after left pneumonectomy were 373.1 ± 16.3 mm Hg ($n = 3$; $P > .1$ vs syngeneic B6 right lung transplant recipients after left pneumonectomy and $P < .05$ vs naive B6 mice after left pneumonectomy).

DISCUSSION

Outcomes after lung transplantation are markedly worse than those after transplantation of all other organs. Development of new experimental models in the mouse will be

FIGURE 2. Gross appearance and histology (hematoxylin and eosin stain, original magnification 100 $\times$) of a right B6 lung not undergoing surgical intervention (A and B), as well as a right B6 lung 7 (C and D; n = 10) and 28 (E and F; n = 3) days after transplantation into a B6 recipient. Tx, Right lung graft.

critical for the design of novel investigations in lung transplant biology. To this end, we and others have recently reported microsurgical techniques for orthotopic left lung transplantation in the mouse.^{5,13,14} Heterotopic murine tracheal transplantation has been widely used as a model for lung transplantation.¹⁵ However, this model is not likely to mirror immunologic and nonimmunologic events that lead to failure of vascularized human lung grafts. Unlike human

lung transplants, tracheal transplants are not vascularized and are not exposed to the external environment. Airway epithelial cells in tracheal allografts undergo apoptosis, leading to fibroproliferation and occlusion of the lumen. In stark contrast, we have recently reported that airway epithelial cells remain intact, maintain differentiation markers, and also upregulate antiapoptotic proteins in nonimmunosuppressed vascularized allografts that have undergone

FIGURE 3. Gross appearance (A) and histology (hematoxylin and eosin stain, original magnification 100 $\times$; B) 7 days after transplantation of a right BALB/c lung into a B6 recipient. Results are representative of 3 independent experiments.

extensive vascular necrosis.¹¹ These observations suggest that airway epithelial cells might be able to protect themselves and that insults in addition to the alloimmune response might be necessary to initiate changes in airway epithelial cells that eventually lead to the fibrotic changes seen in bronchiolitis obliterans.

We have observed several similarities between vascularized murine lung transplants and human lung transplants. Histologic changes of acute rejection in murine lungs are analogous to those observed in human lungs.⁵ Moreover, compared with orthotopic lung transplantation models in the rat, important advantages of the murine model are the wider availability of transgenic and knockout strains, as well as reagents for cellular and molecular analysis. The murine model has already allowed us to gain some new insight into requirements for acute lung rejection. Unlike the case for other organs, CD8⁺ T cells outnumber CD4⁺ T cells in both murine and human lungs that undergo acute rejection.^{7,16} Notably, CD4⁺ T cells are not necessary for the activation of alloreactive CD8⁺ T cells and for the acute rejection of murine lung grafts.⁷ Clinical observations in combined heart–lung transplantations in the 1980s have revealed that lung rejection is initiated at earlier time points than cardiac rejection.¹⁷ A recent study by our group has provided a potential mechanism for this observation.⁶ Although secondary lymphoid organs are critically important for the rejection of other organs, including the heart and skin, alloreactive T cells can be primed locally within lung grafts.¹⁸ In fact, we have observed clustering of recipient T cells around graft-resident dendritic cells at early time points after engraftment, indicating that the lung allograft represents a suitable environment for the activation of alloreactive T cells.

In this study we have modified techniques that we have previously described for transplantation of the left lung.^{5,13} Because of anatomic differences, we found right lung

transplantation to be technically more challenging, which also resulted in longer operative times for both the donor and recipient procedures. Therefore ischemic times are longer for right compared with left lung transplantations when both donor and recipient operations are performed consecutively by the same surgeon. Specifically, the right pulmonary vein is markedly shorter than the left pulmonary vein. We found it technically impossible to insert the cuffed donor pulmonary vein into the main right pulmonary vein of the recipient animal or perform separate inferior and superior pulmonary venous anastomoses, as has been previously described in the rat.¹⁹ Therefore we opted to ligate the superior pulmonary venous branch and use the inferior branch for insertion of the donor vessel. We did not experience pulmonary venous outflow obstruction with this technique.

Bilateral lung transplantation in mice would most closely mirror bilateral sequential lung transplantation in human subjects, which is currently the most commonly used technique worldwide. There are no published reports of successful bilateral lung transplantation in the rat, and in our hands attempts at bilateral sequential lung transplantation in the mouse have resulted in intraoperative animal death. Although further developments of microsurgical techniques might eventually result in successful bilateral lung transplantation in rodents, we had to rely on the removal of the native left lung to limit the recipient's lung function to the transplanted graft.

Unlike previous descriptions for left pneumonectomies in naive mice, we found it critical to place a piece of Gelfoam into the left chest cavity of right lung recipients at the time of the left pneumonectomy to prevent mediastinal shifting, which in our hands led to the death of the animal in the immediate postoperative period.²⁰ Similar to a previous report of right lung transplantation followed by left pneumonectomy in the rat, we performed the left pneumonectomy after a 2-week recovery period after the transplantation of the

FIGURE 4. A, There were no statistically significant differences in arterial PaO₂ levels between wild-type (WT) B6 mice (n = 3) and B6 recipients of syngeneic right B6 lungs 7 days after transplantation (n = 3) after 10 minutes of left hilar occlusion. B, PaO₂ levels were significantly lower in allogeneic BALB/c → B6 right lung transplant recipients (n = 3) than in either syngeneic B6 → B6 right lung transplant recipients (n = 4) or naive B6 mice (n = 4) after a 4-minute period of left hilar occlusion (*P < .01). After 4 minutes of left hilar occlusion, the pH was 7.37 ± 0.02 for naive B6 mice, 7.29 ± 0.02 for B6 → B6 transplant recipients, and 7.19 ± 0.02 for BALB/c → B6 transplant recipients. The Pco₂ was 33.3 ± 2.1 mm Hg for naive B6 mice, 46.9 ± 3.9 for B6 → B6 transplant recipients, and 46.7 ± 2.6 for BALB/c → B6 transplant recipients. The base deficit was 3.3 ± 0.7 for naive B6 mice, 3.6 ± 1.4 for B6 → B6 transplant recipients, and 8.7 ± 0.8 for BALB/c → B6 transplant recipients.

right lung.¹⁹ Attempts at left pneumonectomies less than 2 weeks after right lung transplantation resulted in death of the animals, which might be due to transiently impaired graft function caused by reperfusion injury.²¹ A better understanding of molecular mechanisms contributing to ischemia/reperfusion-related graft dysfunction might allow right lung recipients to tolerate a left pneumonectomy at earlier time points.

To our knowledge, there exist no reports of successful right pneumonectomies in the mouse, which is likely related to an inability of the left lung to compensate for the sudden loss of 70% of overall lung function. To this end, left lung transplant recipients did not survive a right pneumonectomy, even after prolonged recovery periods. Allogeneic right lung

FIGURE 5. There were no statistically significant differences in PaO₂ levels between wild-type B6 mice (n = 3) and B6 recipients of syngeneic right B6 lung transplants 7 days after left pneumonectomy (n = 3, P > .1). Lt PNx, Left pneumonectomy; Syn Rt Tx + Lt PNx, right syngeneic lung transplantation plus left pneumonectomy. Paco₂ levels were 35.2 ± 2.7 and 28 ± 6 mm Hg for the Lt PNx and Syn Rt Tx + Lt PNx groups, respectively. pH was 7.38 ± 0.02 and 7.35 ± 0.02 for the Lt PNx and Syn Rt Tx + Lt PNx groups, respectively. Base deficits were 2.9 ± 1.8 and 8.5 ± 2.6 for the Lt PNx and Syn Rt Tx + Lt PNx groups, respectively. PaO₂ levels in MR1- and cytotoxic T lymphocyte-associated antigen-immunoglobulin-treated BALB/c → B6 right lung recipients after left pneumonectomy (n = 3; Allo Rt Tx + Lt PNx) were 373.1 ± 16.3 mm Hg (P > .1 vs the Syn Rt Tx + Lt PNx group and P < .05 vs the Lt PNx group). Pco₂ levels were 28.3 ± 5.9, pH was 7.36 ± 0.03, and the base deficit was 8.3 ± 1.6.

recipients undergoing acute graft rejection were alive at 7 days after transplantation but displayed respiratory distress. It is possible that a gradual loss of right graft function in these nonimmunosuppressed allogeneic recipients might allow for the development of some compensatory mechanisms by the native left lung.²²

It is important to point out that compensatory growth of the right lung has been well described after left pneumonectomies in mice. This process involves expansion of multiple cell populations, such as endothelial and epithelial cells, and results in complete restoration of the original lung volume.²³ Compensatory lung growth is associated with the rapid upregulation of multiple genes and transcription factors.²⁴ Therefore processes regulating compensatory growth of the graft can influence right lung graft function after a left pneumonectomy and should be taken into consideration when designing studies and interpreting results with this model. Similarly, compensatory lung growth might affect graft function after lobar transplantation in the pediatric population.²⁵

In conclusion, this is the first description of vascularized aerated transplantation of the right lung in the mouse. Histopathologic changes observed in syngeneic and allogeneic grafts are virtually identical to those that we have reported after transplantation of the left lung.⁵ An important advantage of right lung transplantation is that right lung recipients

tolerate a left pneumonectomy where graft failure can result in animal death. Therefore because we have demonstrated that right allograft recipients that are treated with costimulatory blockade can tolerate a left pneumonectomy, this model might be valuable for long-term studies evaluating graft survival in immunosuppressed hosts. Thus this model represents an important addition to the armamentarium in experimental lung transplantation.

We thank Arlene Ligor for the medical illustrations.

References

- Hardy JD, Eraslan S, Dalton ML Jr, Alican F, Turner MD. Re-implantation and homotransplantation of the lung: laboratory studies and clinical potential. *Ann Surg.* 1963;157:707-18.
- Morgan E, Lima O, Goldberg M, Ferdman A, Luk SK, Cooper JD. Successful revascularization of totally ischemic bronchial autografts with omental pedicle flaps in dogs. *J Thorac Cardiovasc Surg.* 1982;84:204-10.
- Toronto Lung Transplant Group. Unilateral lung transplantation for pulmonary fibrosis. *N Engl J Med.* 1986;314:1140-5.
- Mizuta T, Kawaguchi A, Nakahara K, Kawashima Y. Simplified rat lung transplantation using a cuff technique. *J Thorac Cardiovasc Surg.* 1989;97:578-81.
- Okazaki M, Krupnick AS, Kornfeld CG, Lai JM, Ritter JH, Richardson SB, et al. A mouse model of orthotopic vascularized aerated lung transplantation. *Am J Transplant.* 2007;7:1672-9.
- Gelman AE, Li W, Richardson SB, Zinselmeyer BH, Lai J, Okazaki M, et al. Cutting edge: acute lung allograft rejection is independent of secondary lymphoid organs. *J Immunol.* 2009;182:3969-73.
- Gelman AE, Okazaki M, Lai J, Kornfeld CG, Kreisel FH, Richardson SB, et al. CD4+ T lymphocytes are not necessary for the acute rejection of vascularized mouse lung transplants. *J Immunol.* 2008;180:4754-62.
- Corry RJ, Winn HJ, Russell PS. Primarily vascularized allografts of hearts in mice. The role of H-2D, H-2 K, and non-H-2 antigens in rejection. *Transplantation.* 1973;16:343-50.
- Han WR, Murray-Segal LJ, Mottram PL. Modified technique for kidney transplantation in mice. *Microsurgery.* 1999;19:272-4.
- Boisgerault F, Liu Y, Anosova N, Dana R, Benichou G. Differential roles of direct and indirect allorecognition pathways in the rejection of skin and corneal transplants. *Transplantation.* 2009;87:16-23.
- Okazaki M, Gelman AE, Tietjens JR, Ibricevic A, Kornfeld CG, Huang HJ, et al. Maintenance of airway epithelium in acutely rejected orthotopic vascularized mouse lung transplants. *Am J Respir Cell Mol Biol.* 2007;37:625-30.
- Voswinckel R, Motejl V, Fehrenbach A, Wegmann M, Mehling T, Fehrenbach H, et al. Characterisation of post-pneumonectomy lung growth in adult mice. *Eur Respir J.* 2004;24:524-32.
- Krupnick AS, Lin X, Li W, Okazaki M, Lai J, Sugimoto S, et al. Orthotopic mouse lung transplantation as experimental methodology to study transplant and tumor biology. *Nat Protoc.* 2009;4:86-93.
- Jungraithmayr WM, Korom S, Hillinger S, Weder W. A mouse model of orthotopic, single-lung transplantation. *J Thorac Cardiovasc Surg.* 2009;137:486-91.
- Hertz MI, Jessurun J, King MB, Savik SK, Murray JJ. Reproduction of the obliterative bronchiolitis lesion after heterotopic transplantation of mouse airways. *Am J Pathol.* 1993;142:1945-51.
- Donnenberg VS, Burckart GJ, Zeevi A, Griffith BP, Iacono A, McCurry KR, et al. P-glycoprotein activity is decreased in CD4+ but not CD8+ lung allograft-infiltrating T cells during acute cellular rejection. *Transplantation.* 2004;77:1699-706.
- McGregor CG, Baldwin JC, Jamieson SW, Billingham ME, Yousem SA, Burke CM, et al. Isolated pulmonary rejection after combined heart-lung transplantation. *J Thorac Cardiovasc Surg.* 1985;90:623-6.
- Lakkis FG, Arakelov A, Konieczny BT, Inoue Y. Immunologic "ignorance" of vascularized organ transplants in the absence of secondary lymphoid tissue. *Nat Med.* 2000;6:686-8.
- Kawaguchi AT, Mizuta T, Shirai M, Ishibashi-Ueda H, Machida H, Kawashima Y. Right lung transplantation followed by left pneumonectomy in the rat. *Eur J Cardiothorac Surg.* 1996;10:1011-4.
- Sakurai MK, Greene AK, Wilson J, Fauza D, Puder M. Pneumonectomy in the mouse: technique and perioperative management. *J Invest Surg.* 2005;18:201-5.
- Marck KW, Prop J, Wildevuur CR. Lung transplantation in the rat. III. Functional studies in iso- and allografts. *J Surg Res.* 1983;35:149-58.
- Sakurai MK, Lee S, Arsenault DA, Nose V, Wilson JM, Heymach JV, et al. Vascular endothelial growth factor accelerates compensatory lung growth after unilateral pneumonectomy. *Am J Physiol Lung Cell Mol Physiol.* 2007;292:L742-7.
- Fernandez LG, Le Cras TD, Ruiz M, Glover DK, Kron IL, Laubach VE. Differential vascular growth in postpneumonectomy compensatory lung growth. *J Thorac Cardiovasc Surg.* 2007;133:309-16.
- Li D, Fernandez LG, Dodd-o J, Langer J, Wang D, Laubach VE. Upregulation of hypoxia-induced mitogenic factor in compensatory lung growth after pneumonectomy. *Am J Respir Cell Mol Biol.* 2005;32:185-91.
- Binns OA, DeLima NF, Buchanan SA, Lopes MB, Cope JT, Marek CA, et al. Mature pulmonary lobar transplants grow in an immature environment. *J Thorac Cardiovasc Surg.* 1997;114:186-94.