Christophe Jayle, MD
Jamil Hajj-Chahine, MD
Pierre Corbi, MD, PhD
Department of Cardio-thoracic
Surgery
Centre Hospitalier Universitaire de
Poitiers
Université de Poitiers
Poitiers, France

References

1. Augoustides J, Plappert T, Bavaria J. Aortic decision-making in the Loeys-Dietz syndrome: aortic root aneurysm and a normal-caliber ascending aorta and aortic arch. *J Thorac Cardiovasc Surg.* 2009; 138:502-3.
2. Loeys BL, Schwarze U, Holm T, Callewaert BL, Thomas GH, Pannu H, et al. Aneurysm syndromes caused by mutations in the TGF-beta receptor. *N Engl J Med.* 2006;355:788-98.
3. Melenovsky V, Adamira M, Kautznerova D, Voska L, Weichet J, Loeys B, et al. Aortic dissection in a young man with Loeys-Dietz syndrome. *J Thorac Cardiovasc Surg.* 2008;135:1174-5.

doi:10.1016/j.jtcvs.2009.11.066

Reply to the Editor:

We thank Drs Jayle, Hajj-Chahine, and Corbi for their report concerning an aortic root aneurysm in Loeys-Dietz syndrome (LDS) and their subsequent questions related to our clinical approach to this challenging aortopathy.¹ We congratulate this group on the successful surgical management of an aortic root aneurysm associated with LDS.

Their first question is whether to leave an elephant trunk during arch replacement in LDS. Although an elephant trunk is conceptually a possibility in this situation, it is not our standard approach at the University

of Pennsylvania because typically the descending aorta has a normal caliber until it dissects later in the natural history of this disease.²

Their second question relates to timing of surgical intervention for the aortic root in LDS. At the University of Pennsylvania, we recommend aortic root replacement in adults with LDS when the aortic root diameter is 40 to 45 mm. We typically also like to intervene before the development of aortic insufficiency so that, in the absence of aortic valve fenestrations, the native aortic valve can be preserved for optimal hemodynamics and long-term freedom from anticoagulation. In the setting of an aortic root replacement in LDS, we also favor hemiarch replacement under deep hypothermic circulatory arrest, not only to remove all the crossclamped aorta (which is likely at even higher risk for dissection) but also to protect further against future dissection of the distal ascending aorta and aortic arch.^{1,2-4}

Their third question concerned the ongoing surgical management of these patients with this aggressive aortopathy. At the University of Pennsylvania we recommend meticulous monitoring of the thoracic aorta for life in all patients with LDS. This clinical surveillance also includes regular thoracic aortic imaging to detect aneurysm and/or dissection to allow early surgical intervention to prevent death related to the aorta in this syndrome.

In summary, our clinical experience with LDS at the University of Pennsyl-

vania is that the surgical management of the thoracic aorta should be characterized by regular clinical examination, aortic imaging, and an aggressive surgical approach. The aortopathy associated with LDS merits meticulous management, given its life-threatening natural history.

John G.T. Augoustides, MD,
FASE, FAHA^a
Joseph E. Bavaria, MD^b
^aAssociate Professor
Cardiothoracic Section
Anesthesiology and Critical Care
^bProfessor and Vice-Chair
Director of the Thoracic Aortic Center
Division of Cardiothoracic Surgery
Department of Surgery
University of Pennsylvania School of
Medicine
Philadelphia, Pa

References

1. Augoustides J, Plappert T, Bavaria JE. Aortic decision-making in the Loeys-Dietz syndrome: aortic root aneurysm and a normal-caliber ascending aorta and aortic arch. *J Thorac Cardiovasc Surg.* 2009; 138:502-3.
2. Lee RS, Fazel S, Schwarze U, Fleischmann D, Berry GJ, Liang D, et al. Rapid aneurysmal degeneration of a Stanford type B aortic dissection in a patient with Loeys-Dietz syndrome. *J Thorac Cardiovasc Surg.* 2007;134:242-3.
3. Melenovsky V, Adamira M, Kautznerova D, Voska L, Weichet J, Loeys B, et al. Aortic dissection in a young man with Loeys-Dietz syndrome. *J Thorac Cardiovasc Surg.* 2008;135:1174-5.
4. Williams JA, Loeys BL, Nwakanma LU, Dietz HC, Spevak PJ, Patel ND, et al. Early surgical experience with Loeys-Dietz: a new syndrome of aggressive thoracic aortic aneurysm disease. *Ann Thorac Surg.* 2007;83:S757-63.

doi:10.1016/j.jtcvs.2009.12.020

Notice of Correction

Re: CCR3- and CXCR4-mediated interactions regulate migration of CD341 human bone marrow progenitors to ischemic myocardium and subsequent tissue repair. *J Thorac Cardiovasc Surg.* 2008;136:1044-53.

In the above-noted article, the spelling of Dr Sondermeijer's last name was incorrect. The corrected author list is printed below.

N. Bonaros, MD, H. Sondermeijer, MD, M. Schuster, PhD, R. Rauf, MD, S. F. Wang, MD, T. Seki, PhD, D. Skerrett, MD, S. Itescu, MD, and A. A. Kocher, MD