References

1. Benedetto U, Angeloni E, Luciani R, Refice S, Stefanelli M, Comito C, et al. Acute kidney injury after coronary artery bypass grafting: does rhabdomyolysis play a role? *J Thorac Cardiovasc Surg.* 2010;140:464-70.
2. Borders-Hemphill V. Concurrent use of statins and amiodarone. *Consult Pharm.* 2009;24:372-9.
3. Grundy SM, Cleeman JJ, Merz CN, Brewer HB Jr, Clark LT, Hunninghake DB, et al. Implications of recent clinical trials for the National Cholesterol Education Program Adult Treatment Panel III guidelines. *Circulation.* 2004;110:227-39.

doi:10.1016/j.jtcvs.2010.10.016

FUNCTIONAL MITRAL STENOSIS AFTER SURGICAL ANNULOPLASTY FOR ISCHEMIC MITRAL REGURGITATION: IMPORTANCE OF SUBVALVULAR TETHERING IN THE MECHANISM AND DYNAMIC DETERIORATION DURING EXERTION

To the Editor:

Kubota and associates¹ have presented detailed information of the interactions between mitral valve anatomy and function in the presence of mitral insufficiency caused by ischemic ventricular mechanism. They emphasize restriction of flow through the tethered mitral valve at leaflet edge level, which is aggravated by the presence of annuloplasty. The possibility of further aggravation by exercise is well discussed. I find that observations made by me and my colleague, Dr Edward Yellin,²⁻⁴ between 1959 and 2009 provide support for the excellent work described by Kubota and associates.

The normal valve transmits normal flow with a minimal gradient with less than full possible opening. When the normal diastolic annular expansion is fixed by annuloplasty, there is always a small gradient and a reduction of the mitral area to less than the annular area. However, as Kubota and associates found, the echo-derived mitral areas are much less than the geometric area of the annuloplasty rings used. They attribute this "functional mitral

stenosis" to leaflet tethering at the free edge with a reduced separation of the central anterior and posterior leaflet tips. A paper presented at the July 2009 meeting of the Society for Heart Valve Disease in Berlin (Frater, Papillary Displacement: Effect on Chordae and Leaflets) supports and confirms this opinion.

This was an in vitro anatomic observation of the effects of papillary displacement on the anterior and posterior leaflets in 6 pig and 3 human hearts with fixed reducing annuloplasties. In normal diastole the papillary muscles move apically, posteriorly, and apart.⁵ These are also the papillary displacements seen in myopathic ventricular enlargement. Apical displacement by itself puts the second-order chordae under tension, leaving the free edge first-order chordae loose, and allows the leaflet to retain flexibility. Separation of the papillary muscles, by contrast, puts both the second-order and the free edge first-order chordae under tension. Tautness of the free-edge chordae reduces all flexibility of the major parts of the anterior leaflet and the central scallop, rendering them resistant to opening by transvalvular flow. Whereas the normal separation of papillary muscles is 1 cm in systole and 2 cm in diastole, in a ventricle with a 65-mm diastolic dimension, the papillary separation may be 4 cm. The anterior leaflet is a taut sheet pointing posteriorly and the stiffened posterior scallops are parallel with the ventricular wall. The findings of this in vitro study provide clear confirmation of the mechanism of mitral stenosis at free edge level postulated by Kubota and associates.

Robert W. M. Frater, MD
Montefiore Medical Center
Department of Cardiothoracic
Surgery
Bronx, NY

References

1. Kubota K, Otsuji Y, Ueno T, Koriyama C, Levine RA, Sakata R, et al. Functional mitral stenosis after surgical annuloplasty for ischemic mitral regur-

gitation: importance of subvalvular tethering in the mechanism and dynamic deterioration during exertion. *J Thorac Cardiovasc Surg.* 2010;140:617-23.

2. Frater RWM. Mitral valve anatomy and prosthetic valve design. *Proc Staff Meet Mayo Clin.* 1961;36:582-92.
3. Yellin EL, Peskin C, Yoran C, Koenigsberg M, Matsumoto M, Laniado S, et al. Mechanism of mitral valve motion during diastole. *Am J Physiol.* 1980;241(Heart Circ. Physiol 10):H389-400.
4. Yellin EL, Nikolic S, Frater RWM. Left ventricular filling dynamics and diastolic function. *Prog Cardiovasc Dis.* 1990;32:247-71.
5. Babburi H, Oommen R, Brofferio A, Ilcail A, Frater RWM, Shirani J. Functional anatomy of the normal mitral apparatus: a transthoracic two-dimensional echocardiographic study. *J Heart Valve Dis.* 2003;12:180-5.

doi:10.1016/j.jtcvs.2010.10.017

Reply to the Editor:

We appreciate the comments of Dr Robert Frater on our recent manuscript¹ and are honored to reply to them. He has been a pioneer in recognizing that mitral leaflet diastolic opening, not only systolic closure, is controlled by the chordal apparatus.^{2,3} He and his colleagues observed only limited influence from changes in mitral inflow volume on mitral leaflet opening excursion.³ They further observed that papillary muscle rupture clearly increases mitral leaflet opening, indicating limited leaflet opening by chordal attachments.³ Therefore, they demonstrated flow-independent opening of the mitral leaflets as well as diastolic restriction or tethering of mitral leaflet opening by chordal attachments and tension, subsequently confirmed in patients by our group.⁴

In dilated ventricles, papillary muscle displacement causes greater chordal tension and restricts leaflet mobility in both systole and diastole. It leads to mitral regurgitation in systole and reduced leaflet opening in diastole.^{4,5} Therefore, subvalvular tethering by papillary muscle displacement is the central pathophysiology restricting mitral leaflet mobility and function. Of note is that ventricular dilatation alone causes significant mitral regurgitation whereas resultant mitral stenosis is