colleagues.¹ They introduced a case series with the use of innominate cannulation during repair of Stanford type A aortic dissection with ascending aorta and hemiarch replacement combined with stent-graft elephant trunk. We congratulate them for their excellent clinical results and also would like to discuss a few questions regarding cannulation during antegrade cerebral perfusion (ACP).

The combined use of ACP with hypothermic circulatory arrest (HCA) is widely accepted as a routine means of protecting the brain during complex aortic arch reconstruction; however, the risks of brain complications related to HCA and extensive ACP times also remain a substantial burden in aortic arch construction. Moreover, technical issues relating to the use of ACP during HCA, such as cannulation sites and types for ACP, unilateral perfusion or bilateral perfusion, and the flow rate and blood pressure used to selectively perfuse the brain, are still uncertain. ACP has become recognized as an important mode for arch replacement and can be performed in several different ways: unilateral cerebral perfusion through the right axillary artery or the right brachial artery, or the innominate artery. It is well documented that unilateral cerebral perfusion through the right axillary artery by direct cannulation has been successfully used in a majority of heart centers.² In a previous letter published elsewhere,³ we discussed the benefits and risks with Xu, whose group used a side vascular graft to perfuse the innominate artery in conjunction with the use of HCA during aortic arch surgery. Shi and colleagues¹ also summarized benefits with cannulation at the innominate artery site relative to right axillary artery cannulation: (1) No additional incision is needed. (2) The same technique and same size of cannula are used as in ascending aortic cannulation. (3) Cerebral perfusion pressure can be monitored continuously with a right arterial line.

Relative to Xu's method, cannulation at the innominate artery has more limitations. First, cannulation at the innominate artery avoids the additional incision, but you may have another cannula in your surgical field. Although the innominate artery is larger than the right axillary artery, it is shorter; moreover, you have to choose a cannula with a curved tip, because the tip of the cannula should be adjusted in direction at least twice during the procedure. Adjustment of the direction of the cannula may cause damage to the artery.⁴ A cannula with a curved tip generates greater pressure drops and may cause more blood damage. Moreover, as we mentioned in our previous letter,³ we still think that there is a potential risk when the innominate artery is used to perfuse the brain. When the innominate artery is used, blood flow distribution may depend on the resistance of the right axillary artery and the left common carotid artery. Our concern is that if the resistance of the right axillary artery decreases or increases, this may result in malperfusion to the brain. Although this has not been widely recognized in the study of ACP flow during HCA, both overperfusion and underperfusion have been shown to be detrimental to the brain. In our previous research, when the ACP (unilateral cerebral perfusion through the right axillary artery by direct cannulation) flow was lower than 10 mL/kg, transcranial Doppler ultrasonography could not detect the signal of blood flow in the middle cerebral artery.⁵ We think that no matter which way is used during aortic arch surgery, it is most important that we know whether the brain is at a suitable ratio of blood flow and oxygen consumption. To achieve this purpose, the best way to monitor the brain status is by combining transcranial Doppler ultrasonography with near-infrared spectroscopy during this procedure, not relying on pressure alone.

Bingyang Ji, MD
Xiaohua Wang, MD
Jinping Liu, MD
Cun Long, MD

Department of Cardiopulmonary
Bypass
Chinese Academy of Medical Science
Peking Union Medical College
Fuwai Hospital and Cardiovascular
Institute
Beijing, People's Republic of China

References

1. Shi E, Gu T, Yu L, Xiu Z, Zhang Z, Wang C, et al. Repair of Stanford type A aortic dissection with ascending aorta and hemiarch replacement combined with stent-graft elephant trunk technique by using innominate cannulation. *J Thorac Cardiovasc Surg*. 30 March 2011 [Epub ahead of print].
2. Di Eusano M, Schepens MA, Morshuis WJ, Dossche KM, Di Bartolomeo R, Pacini D, et al. Brain protection using antegrade selective cerebral perfusion: a multicenter study. *Ann Thorac Surg*. 2003;76:1181-9.
3. Ji B, Liu J, Chang Q, Yu C, Long C. Benefits and risks with using cannulation at the innominate artery with a side graft during arch surgery. *Ann Thorac Surg*. 2011;91:1649-50.
4. Undar A, Ji B, Rider A, Lukic B, Kunselman AR, Weiss WJ, et al. Comparison of four different pediatric 10F aortic cannulae during pulsatile versus non-pulsatile perfusion in a simulated neonatal model of cardiopulmonary bypass. *ASAIO J*. 2007;53:778-84.
5. Ji B, Duan X, Miao Na, Liu R, Yu K, Yang J, et al. Neuromonitoring combine TCD with NIRS during total aorta arch replacement procedure. *Chinese J ECC*. 2011;9:71-7.

doi:10.1016/j.jtcvs.2011.07.064

Reply to the Editor:

We thank Ji and colleagues for their insightful comments on the technique of innominate artery cannulation during surgery to correct Stanford type A aortic dissection as described in our recently published article.¹

Although one more cannula is in the surgical field when the innominate artery cannulation technique is used, we do not think that this cannula will add any difficulty to the operation. Gentle adjustment of the direction of the cannula by an experienced surgeon will do no damage to the artery. In Ji and colleagues' opinion, a cannula with a curved tip may generate higher pressure drop and cause more blood damage. In contrast, we think that the

diameter of the cannula is the key factor in generating the pressure drop, rather than the shape of the tip. In addition, relative to axillary artery cannulation, a larger size of cannula can be used for innominate artery cannulation, which may decrease blood damage, especially during periods of high flow rate.

The main comment of Ji and colleagues is that during antegrade cerebral perfusion (ACP), when the innominate artery is used as the cannulation site, blood flow distribution may depend on the resistance of 2 arteries, the right axillary artery and the right common carotid artery. Thus any decrease or increase in the resistance of the right axillary artery could result in malperfusion of the brain. Compared with direct axillary artery cannulation technique, the perfusion pressure for the common carotid artery can be monitored by direct right radical artery cannulation. Therefore, ACP can be adjusted by flow rate combined with perfusion pressure. Usually a flow rate of 8 to 10 mL · kg⁻¹ · min⁻¹ is used for ACP, while the perfusion pressure may be maintained between 40 and 50 mm Hg. When the perfusion pressure is too low, the flow rate can be increased to avoid underperfusion to the brain. When the perfusion pressure is too high, such as higher than 70 mm Hg, the flow rate can be decreased. We therefore believe that this technique may be better than ACP with a fixed flow rate. Our clinical results have also confirmed that ACP with innominate artery cannulation is an acceptable brain protective method during arch surgery.¹ We also do agree that transcranial Doppler ultrasonography combined with near-infrared spectroscopy is a more accurate means of monitoring brain perfusion state.

Enyi Shi, MD, PhD
Tianxiang Gu, MD, PhD
Department of Cardiac Surgery
First Affiliated Hospital
China Medical University
Shenyang, China

Reference

1. Shi E, Gu T, Yu L, Xiu Z, Zhang Z, Wang C, et al. Repair of Stanford type A aortic dissection with ascending aorta and hemiarch replacement combined with stent-graft elephant trunk technique by using innominate cannulation. *J Thorac Cardiovasc Surg*. 30 Mar 2011 [Epub ahead of print].

doi:10.1016/j.jtcvs.2011.08.029

DABIGATRAN ETEXILATE IN MECHANICAL VALVE THROMBOPROPHYLAXIS

To the Editor:

I read with interest the recently published study of McKellar and colleagues regarding the effectiveness of dabigatran etexilate for thromboprophylaxis of mechanical heart valves.¹ In this animal study, McKellar and colleagues¹ used a model of heterotopic aortic valve replacement in which, as described previously, the valve within the modified valved conduit remains open continuously throughout the cardiac cycle.² Surprisingly, this very important information regarding the valve function after this extra-anatomic valved conduit placement is missing from their recent study as a possible major limitation.¹

Historically, successful aortic valve replacement is considered the implantation of a mechanical or tissue valve in the aortic position with the prosthetic valve leaflets remaining open only during systole and closed during diastole. I believe that this historic definition of successful aortic valve replacement has never been challenged and certainly has never changed, so I have the following questions: (1) Did the authors make a modification in their model leading to a periodic, synchronous to cardiac cycle opening and closing of the mechanical valve? (2) If no modification to their previously published model has been made, how can they translate their findings into clinical aortic valve replacement practice in light of the extra-anatomic position of the mechanical prosthetic valve and the fact that this valve function is completely different from that considered clinically successful?

Mihalis E. Argiriou, MD
Evangelismos Hospital
Athens, Greece

References

1. McKellar SH, Abel S, Camp CL, Suri RM, Erath MH, Schaff HV. Effectiveness of dabigatran etexilate for thromboprophylaxis of mechanical heart valves. *J Thorac Cardiovasc Surg*. 2011; 141:1410-6.
2. McKellar SH, Thompson JL, Schaff HV. A model of heterotopic aortic valve replacement for studying thromboembolism prophylaxis in mechanical valve prostheses. *J Surg Res*. 2007; 141:1-6.

doi:10.1016/j.jtcvs.2011.05.036

Reply to the Editor:

We appreciate Argiriou's comments. The heterotopic porcine model described in our study is useful for screening antithrombotic therapy for mechanical valves.¹⁻³ The procedure is less costly in terms of time and equipment than heterotopic valve replacement, and mortality among experimental animals is relatively low. Rapid, robust thrombus formation is important in the swine model because of somatic growth and the considerable expense of housing animals for long periods. It is true that the tilting discs of the bileaflet valves do not open and close normally; however, we actually consider this an advantage of the model in that lack of leaflet washing might actually be expected to increase thrombogenicity of the valve. Indeed, we previously compared thrombus formation on mechanical valve prostheses in the heterotopic aortic and orthotopic mitral positions, and at 30 days, thrombus formation was greater on the heterotopically placed valves.⁴ Thus any therapies that reduce or eliminate thrombus formation in this heterotopic model should be effective in an orthotopic model.

Stephen H. McKellar, MD, MSc
Hartzell V. Schaff, MD
Division of Cardiovascular Surgery
Mayo Clinic College of Medicine
Rochester, Minn