subsequent reduction of renal blood flow observed during hypothermia are probably unrelated to increased levels of circulating catecholamines or increased renal sympathetic nerve activity. Other metabolic process taking place in response to lowering of temperature have been proposed.⁴

Overall, the clinical evidence regarding the potential benefits of hypothermia during pediatric heart surgery are not convincing. In fact, there is increasing retrospective evidence⁵ showing several benefits of normothermic cardiopulmonary bypass. Our own study has provided the first randomized trial evidence of the impact of normothermic cardiopulmonary bypass on renal function, contributing to the accumulating evidence of this technique in a low-risk group of children undergoing heart surgery.

Much more work is needed if we are to understand whether normothermia is safe in patients at higher risk, such as neonates and patients with complex congenital heart disease, who are more likely to be affected by cardiopulmonary bypass-induced postoperative renal dysfunction.

Massimo Caputo, MD^{a,b}

Chris A. Rogers, PhD^b

^aBristol Royal Hospital for Children
Bristol, UK

^bBristol Heart Institute
University of Bristol
Bristol, UK

References

1. Tveita T, Johansen K, Lien AH, Myklebust R, Lindal S. Morphological changes in tubular cells from in situ kidneys following experimental hypothermia and rewarming. *APMIS*. 2005;113:13-20.
2. Tveita T, Ytrehus K, Skandfer M, Oian P, Helset E, Myhre ES, et al. Changes in blood flow distribution and capillary function after deep hypothermia in rat. *Can J Physiol Pharmacol*. 1996;74:376-81.
3. Delin NA, Kjartansson KB, Pollock L, Schenk WG Jr. Redistribution of regional blood flow in hypothermia. *J Thorac Cardiovasc Surg*. 1965;49:511-6.
4. Broman M, Källskog O, Kopp UC, Wolgast M. Influence of the sympathetic nervous system on renal function during hypothermia. *Acta Physiol Scand*. 1998;163:241-9.

5. Durandy Y, Hulin S. Intermittent warm blood cardioplegia in the surgical treatment of congenital heart disease: clinical experience with 1400 cases. *J Thorac Cardiovasc Surg*. 2007;133:241-6.

doi:10.1016/j.jtcvs.2011.12.013

RESPIRATORY COMPLICATIONS IN PATIENTS WITH HETEROTAXY SYNDROME

To the Editor:

We read the interesting paper by Swisher and associates¹ on the respiratory complications after cardiac surgery in patients with heterotaxy syndrome. The authors compared the postoperative course of children with heterotaxy syndrome with that of patients with cardiac defects and similar surgical complexities following the Risk Adjustment in Congenital Heart Surgery.^{1,2}

The patients with heterotaxy syndrome showed higher surgical mortality and increased postoperative respiratory complications including postoperative hospital stay, longer mechanical ventilation, need for tracheostomies, and extracorporeal membrane oxygenation support.¹ Other very recent articles underlined specific problems and specific surgical solutions in children with heart disease and laterality defects.^{3,4} All these observations confirm the notion that the presence of genetic syndromes may significantly affect the perioperative management of patients with congenital heart defect.⁵

As described for patients with trisomy 21, deletion 22q11, Turner syndrome, and other genetic syndromes,⁵ the knowledge of specific cardiovascular and extracardiac defects deserves particular attention in the perioperative period in terms of specific diagnostic, operative, and intensive care protocols.

Heterotaxy syndrome characterized by complex cardiovascular malformations and extracardiac defects deserves specific cardiac and extracardiac preoperative assessment to recognize risk factors and to reduce the

operative mortality and postoperative complications.^{1,3,4}

Vanessa Martucci, MD

Dario Marino, MS

Paolo Versacci, MD

Department of Pediatrics

Sapienza University of Rome

Viale Regina Elena

Rome, Italy

References

1. Swisher M, Jonas R, Tian X, Lee ES, Lo CW, Leatherbury L. Increased postoperative and respiratory complications in patients with congenital heart disease associated with heterotaxy. *J Thorac Cardiovasc Surg*. 2011;141:637-44.
2. Jenkins KJ, Gauvreau K, Newburger JW, Spray TL, Moller JH, Iezzoni LI. Consensus-based method for risk adjustment for surgery for congenital heart disease. *J Thorac Cardiovasc Surg*. 2002;123:110-8.
3. McElhinney DB, Marx GR, Marshall AC, Mayer JE, Del Nido PJ. Cavopulmonary pathway modification in patients with heterotaxy and newly diagnosed or persistent pulmonary arteriovenous malformations after a modified Fontan operation. *J Thorac Cardiovasc Surg*. 2011;141:1362-70.
4. O'Connor MJ, Ravishanker C, Ballweg JA, Gillespie MJ, Gaynor JW, Tabbatt S, et al. Early systemic-to-pulmonary artery shunt intervention in neonates with congenital heart disease. *J Thorac Cardiovasc Surg*. 2011;142:106-12.
5. Formigari R, Michielon G, Digilio MC, Piacentini G, Carotti A, Giardini A, et al. Genetic syndromes and congenital heart defects: how is surgical management affected? *Eur J Cardiothorac Surg*. 2009;35:606-14.

doi:10.1016/j.jtcvs.2011.10.086

Reply to the Editor:

We thank Dr Martucci and her colleagues for their supportive letter, which emphasizes the general point that the presence of genetic syndromes may significantly affect the perioperative management of patients with a congenital heart defect. Specifically for patients with heterotaxy, we agree that these complex patients require "cardiac and extracardiac preoperative assessment to recognize risk factors and to reduce the operative mortality and postoperative complications." Extrapolating beyond the phenotype-genotype-prognosis paradigm of their cited reference,¹ we suggest adding "therapy" to the paradigm.

Patients in our study with heterotaxy and congenital heart disease (CHD) showed high postoperative morbidity/mortality, with some having respiratory complications.² Although this respiratory phenotype is often attributed to the CHD, airway mucus clearance and left-right patterning, abnormal in heterotaxy syndrome, both require motile ciliary function. Thus airway ciliary dysfunction (CD) similar to that of primary ciliary dyskinesia (PCD) may have relevance for increased respiratory complications seen in patients with heterotaxy. To better characterize the phenotype of a broad spectrum of patients with heterotaxy, 43 patients with CHD were assessed for airway CD. We used videomicroscopy to look for abnormal ciliary motion in nasal tissue and nasal nitric oxide noninvasive testing (typically low with PCD) to assess for CD. A remarkable 42% (18/43) of patients exhibited CD, with one third in the severe PCD range. Patients greater than 6 years old with CD showed increased respiratory disease similar to that seen in PCD.³ Our studies suggest patients with CHD who have heterotaxy have substantial risk of CD and respiratory disease that complicates their prognosis.

The phenotype-genotype-prognosis correlation is also important in that further research may allow future patients and families to correlate gene mutations with phenotype and prognosis. Now families face the very high recurrence risk ratio of 79.1 for heterotaxia (95% confidence interval, 32.9-190).⁴ Further study of our patients with heterotaxy who have CD through next-generation sequencing of all 14 known PCD genes in 13 heterotaxy patients with CD identified 10 novel coding variants in 6 PCD genes. Recently, some CD gene mutations have become commercially available. The paradigm of phenotype-genotype-prognosis is beneficial to promote translational research to the patient, family, and medical care team. Our research suggests that patients with CHD and heterotaxy will

be at higher perioperative risk if they have CD.³ These findings indicate that patients with CHD and heterotaxy may benefit from preoperative screening for CD. Recognition of specific surgical risks factors in genetic syndromes is leading to specific diagnostic protocols that improve phenotyping or predictive genotyping.¹ To add "therapy" to the paradigm, further studies are needed to evaluate whether perioperative protocols improving airway ciliary motility and enhancing mucus clearance may reduce respiratory complications and improve postoperative outcome for patients with CHD who have heterotaxy and CD.

Linda Leatherbury, MD^a

Cecilia W. Lo, PhD^b

Richard Jonas, MD^a

^aChildren's National Heart Institute
Children's National Medical Center
Washington, DC

^bDepartment of Developmental
Biology

University of Pittsburgh School of
Medicine
Pittsburgh, Pa

References

1. Formigari R, Michielon G, Digilio MC, Piacentini G, Carotti A, Giardini A, et al. Genetic syndromes and congenital heart defects: how is surgical management affected? *Eur J Cardiothorac Surg*. 2009;35:606-14.
2. Swisher M, Jonas R, Tian X, Lee ES, Lo CW, Leatherbury L. Increased postoperative and respiratory complications in patients with congenital heart disease associated with heterotaxy. *J Thorac Cardiovasc Surg*. 2011;141:637-44.
3. Nakhleh N, Swisher M, Francis R, Giese R, Chatterjee B, Connelly P, et al. Low nasal nitric oxide and ciliary dysmotility in patients with congenital heart disease and heterotaxy. *Am J Respir Crit Care Med*. 2009;179:A1223.
4. Oyen N, Poulsen G, Boyd HA, Wohlfahrt J, Jensen PKA, Melbye M. Recurrence of congenital heart defects in families. *Circulation*. 2009;120:295-301.

doi:10.1016/j.jtcvs.2011.11.006

LATE EROSION OF AN AMPLATZER SEPTAL OCCLUDER DEVICE

To the Editor:

We read with interest the article "Late erosion of an Amplatzer septal

occluder device 6 years after placement" by Taggart, Dearani, and Hagler¹ in the *Journal of Thoracic and Cardiovascular Surgery*.

It is assumed that cardiac perforation is a rare complication following transcatheter device closure of an atrial septal defect (ASD) or patent foramen ovale (PFO). We, too, had a patient who had similar complications of perforation 9 months after closure of an ASD with a BioSTAR septal occluder (NMT Medical, Boston, Mass).² Another patient came to us 3 years after ASD closure with a Solysafe device (Swissimplant AG, Solothurn, Switzerland) with complications of device fracture and embolization into the pulmonary artery. Our patients warranted emergency surgical exploration for device explantation, repair of the erosion, along with pericardial patch closure of the residual septal defect. Although innovative absorbable closure devices that gradually biodegrade after implantation have theoretically decreased the long-term complications, it must be borne in mind that these devices are not totally bioabsorbed. The nonabsorbable components, such as the metal struts, may create long-term complications by eroding into the adjacent tissues.

This case clearly illustrates that inasmuch as tissue erosion, and other complications such as strut fracture, may occur at any time starting from 72 hours to 6 years after implantation, echocardiographic follow-up is needed, not just in the early stages, but also in the long-term. We also believe that clearer guidelines and recommendations may help prevent the complications associated with septal occluder devices.

Although cardiac perforation may be "rare," it is a potentially "serious" complication after transcatheter device closure procedures. These patients arrived at the hospital in time to benefit from immediate diagnosis and emergency surgical treatment of their complications. The question we need to ask ourselves and also