bypass and will pull on the patch and perhaps on the AV valves, possibly making them leak. There is usually some atrial septum available, but if not, I would put the hole and snare as far lateral as possible, which should make the pull on the rest of the repair negligible.

John E. Foker, MD, PhD
on behalf of all authors

Division of Cardiothoracic Surgery
University of Minnesota
Minneapolis, Minn

References

1. Foker JE, Berry JM, Steinberger J. Ventricular growth stimulation to achieve two-ventricle repair in unbalanced common atrioventricular canal. *Prog Pediatr Cardiol.* 1999;10:173-86.
2. Foker JE, Berry JM, Harvey BA, Pyles LA. Mitral and tricuspid valve repair and growth in unbalanced atrial ventricular canal defects. *J Thorac Cardiovasc Surg.* 2012;143:S29-32.

<http://dx.doi.org/10.1016/j.jtcvs.2012.06.033>

WILL LOCAL USE OF ANTIBIOTICS BE ESSENTIAL?

To the Editor:

We appreciate the study by Schimmer and colleagues¹ on gentamicin-collagen sponge use for heart surgery. They demonstrated a reduction of wound complication after surgery by gentamicin-collagen sponge use.

We agree with Schimmer and colleagues¹ and strongly recommend applying antibiotics directly to the surgical site. Our previous study was on the prevention of surgical site infection by antibiotic spraying.² We used not only gentamicin but also cefazolin. In our study, the incidence of deep surgical site infection was 0.50% in 2612 patients. We chose gentamicin and cefazolin because they cover the most likely contaminants, such as *Staphylococcus aureus* and *Pseudomonas aeruginosa*. In the study by Schimmer and colleagues,¹ *S aureus* cases were low despite the use of only gentamicin.

Local use of antibiotics is not accepted by guidelines for prevention of

surgical site infection³ because there is no strong evidence. The references of the guidelines did not report better results than Schimmer and colleagues.¹

In reality, many surgeons apply antibiotics by local spraying or subcutaneous injection. We think local use of antibiotics is an essential method for postoperative reduction of wound complications.

Hiroshi Osawa, MD, PhD
Kouan Orii, MD, PhD

Division of Cardiothoracic Surgery
Shimada General Hospital
Chiba, Japan

References

1. Schimmer C, Özkur M, Sinha B, Hain J, Gorski A, Hager B, et al. Gentamicin-collagen sponge reduces sternal wound complications after heart surgery: a controlled, prospectively randomized, double-blind study. *J Thorac Cardiovasc Surg.* 2012;143:194-200.
2. Yoshii S, Hosaka S, Suzuki S, Takahashi W, Okuwaki H, Osawa H, et al. Prevention of surgical site infection by antibiotic spraying in the operative field during cardiac surgery. *Jpn J Thorac Surg.* 2001;49:279-81.
3. Guideline for prevention of surgical site infection, 1999. *Infect Control Hosp Epidemiol.* 1999;20:247-78.

<http://dx.doi.org/10.1016/j.jtcvs.2012.05.079>

Reply to the Editor:

We appreciate the kind comments¹ on our publication “Gentamicin-Collagen Sponge Reduces Sternal Wound Complications After Heart Surgery: A Controlled, Prospectively Randomized, Double-Blind Study”² regarding our results of the routine prophylactic retrosternal use of a gentamicin-collagen sponge in patients undergoing cardiac surgery. Our answers to the 3 points included in the letter to the Editor are as follows:

1. The most likely contaminants of deep sternal wound infections (mediastinitis) in our study were coagulase-negative staphylococci (68.4%), gram-negative bacteria (10.5%), *Propionibacterium acnes* (10.5%), and *Staphylococcus aureus* (5.3%).² This microbial

spectrum is similar to that in the Swedish study by Friberg and colleagues,³ in which coagulase-negative staphylococci formed the largest group of pathogens. The number of *S aureus* cases accounted for 2.4% (1 case) in the presented study, 21.7% in a Swedish study, and 37.3% in a US study.⁴ In our study and the Swedish study, there were no patients with methicillin-resistant *S aureus*. By contrast, methicillin-resistant *S aureus* infection was identified in 10.2% of the microbial isolates (27.3% of the *S aureus* cases) in the US study. The extent to which the dissimilar microbial spectra and possible antibiotic resistance influence the results cannot be clarified.

2. The guidelines for prevention of surgical site infections investigating intraoperative gentamicin implant before wound closure involve 2 randomized controlled trials.^{3,5,6} Figure 1 shows that the 2 studies were combined in a meta-analysis, and that a statistically significant difference was found favoring the gentamicin implant (odds ratio, 0.49; 95% confidence interval, 0.34-0.68; $I^2 = 0\%$).⁶

Engelmann and colleagues⁷ summarized these 2 studies in 2007 by observing that a slight reduction in infection was seen, but the population was too small to draw a definitive conclusion. Although the use of topical antibiotics is controversial, and they are not used by most cardiac surgeons, the existing studies demonstrate a reduction in the wound infection rate. More study is warranted before topical antibiotics can be recommended as a standard prophylaxis.⁷ Our study is a randomized controlled trial and therefore fulfils this demand and is a contribution toward more evidence in the prevention of surgical site infection.

FIGURE 1. Meta-analysis of 2 trials comparing the effect in cardiac surgery on surgical site infection incidence of inserting a gentamicin implant before wound closure versus no insertion of a gentamicin implant. *CI*, Confidence interval; *OR*, odds ratio.

3. We absolutely agree with Osawa and Orii¹ that in reality many surgeons use antibiotics locally, and that antibiotic use plays an essential part in many different methods to reduce sternal wound infection.

Christoph Schimmer, MD
 Rainer Leyh, MD, PhD
 Department of Cardiothoracic and
 Thoracic Vascular Surgery
 University of Würzburg
 University Hospital Würzburg
 Würzburg, Germany

References

- Osawa H, Orii K. Will local use of antibiotics be essential? *J Thorac Cardiovasc Surg.* 2012;144:742.
- Schimmer C, Oezkur M, Shinha B, Hain J, Groski A, Hager B, et al. Gentamicin-collagen sponge reduces sternal wound complications after heart surgery: a controlled, prospectively randomized, double-blind study. *J Thorac Cardiovasc Surg.* 2012;143:194-200.
- Friberg Ö, Svedjeholm R, Söderquist B, Granfeldt H, Vikerfors T, Källman J. Local gentamicin reduces sternal wound infections after cardiac surgery: a randomized controlled trial. *Ann Thorac Surg.* 2005;79:153-62.
- Bennett-Guerrero E, Ferguson TB, Lin M, Garg J, Mark DB, Scavo VA, et al. Effect of an implantable gentamicin-collagen sponge on sternal wound infections following cardiac surgery. *JAMA.* 2010;304:755-62.
- Clinical Guideline. *Surgical Site Infection. Prevention and Treatment of Surgical Site Infection.* Commissioned by the National Institute for Health and Clinical Excellence. RCOG Press; 2008:74.
- Eklund A, Valtonen M, Werkkala K. Prophylaxis of sternal wound infections with gentamicin-collagen implant: randomized controlled study in cardiac surgery. *J Hosp Infect.* 2005;59:108-12.
- Engelmann R, Shahian D, Shemin R, Guy TS, Bratzler D, Edwards F, et al. The Society of Thoracic Surgeons practice guideline series: antibiotic prophylaxis in cardiac surgery, part II:

antibiotic choice. *Ann Thorac Surg.* 2007;83:1569-76.

<http://dx.doi.org/10.1016/j.jtcvs.2012.06.017>

INTERLEUKIN-6 AND -10 AS MASTER PREDICTIVE MEDIATORS OF THE POSTCARDIOPULMONARY BYPASS INFLAMMATORY RESPONSE

To the Editor:

We read with great interest the well-documented work of Jouan and colleagues¹ on gene polymorphism and cytokine plasma levels as predictive factors of complications after cardiopulmonary bypass (CPB). Previous studies highlighted potential roles of cytokines (eg, interleukin [IL]-6 and -10) on post-CPB complications^{2,3} but without mechanistic explanations of these elevated productions. The interesting and well-documented study of Jouan and colleagues¹ now gives this mechanistic explanation. Genetic testing of IL6-572G>C and IL10-592C>A might thus be a tool for identifying patients at the highest risk of poor tolerance to the post-CPB inflammatory response. Another main interest of this study is to bring some doubts concerning the role (if any) of other pro- or anti-inflammatory compounds released during CPB, such as angiogenic growth factors and lipidic mediators, on post-CPB adverse effects.^{4,5} It is clear that IL-6 and IL-10 now can be considered as the master predictive control mediators of the post-CPB inflammatory response.

Yves Denizot, PhD^a
 Nathalie Nathan, MD, PhD^b
^aUniversité de Limoges
 Limoges, France
^bService d'Anesthésie Réanimation
 Chirurgicale
 CHU Dupuytren
 Limoges, France

References

- Jouan J, Golmard L, Benhamouda N, Durrleman N, Golmard JL, Ceccaldi C, et al. Gene polymorphisms and cytokine plasma levels as predictive factors of complications after cardiopulmonary bypass. *J Thorac Cardiovasc Surg.* 2012;144:467-73.
- Nathan N, Preux PM, Feiss P, Denizot Y. Plasma interleukin 4, 10 and 13 concentrations and complications after coronary artery bypass grafting surgery. *J Cardiothorac Vasc Surg.* 2000;14:156-60.
- Denizot Y, Lorgeot V, Cornu E, Nathan N. Plasma leukemia inhibitory factor, interleukin-6 and soluble interleukin-6 receptor levels during cardiopulmonary bypass with extracorporeal circulation. *Cytokine.* 1998;10:303-6.
- Denizot Y, Guglielmi L, Cornu E, Nathan N. Alterations in plasma angiogenic growth factor concentrations after coronary artery bypass graft surgery: relationships with post-operative complications. *Cytokine.* 2003;24:7-12.
- Nathan N, Denizot Y, Cornu E, Jauberteau MO, Chauvreau C, Feiss P. Cytokine and lipid mediator blood concentrations after coronary artery surgery. *Anesth Analg.* 1997;85:1240-6.

<http://dx.doi.org/10.1016/j.jtcvs.2012.05.080>

Reply to the Editor:

We appreciate the comments of Denizot and Nathan on our article “Gene Polymorphisms and Cytokine Plasma Levels as Predictive Factors of Complications After Cardiopulmonary Bypass.”¹ However, even if we