Risk factors for nonocclusive mesenteric ischemia after elective cardiac surgery

Heinrich Volker Groesdonk, MD,^{a,b} Matthias Klingele, MD,^c Sandra Schlempp, MS,^a Hagen Bomberg, MD,^a Wolfram Schmied, Dipl Psych,^a Peter Minko, MD,^d and Hans-Joachim Schäfers, MD^a

Objective: Nonocclusive mesenteric ischemia (NOMI) may occur after cardiopulmonary bypass. It is crucial to early identify patients who are at risk of developing this complication. The aim of this prospective study was to evaluate perioperative risk factors in a large cohort of patients undergoing elective cardiac surgery.

Methods: From January 1, 2010, to March 31, 2011, all patients scheduled for elective cardiac surgery were screened for participation in this trial. If NOMI was suspected, arterial angiography was performed. NOMI and non-NOMI patients were compared with respect to all variables assessed in this study. Additionally, odds ratios were calculated. Linear discriminant analyses as well as logistic regression analyses were performed to develop a model that identifies patients at risk for developing NOMI.

Results: Eight hundred sixty-five patients were included in the study, of whom 78 developed NOMI. Among preoperative parameters, renal insufficiency, diuretic therapy, and age >70 years showed the highest odds ratios for postoperative NOMI. The highest odds ratios for development of NOMI were observed with postoperative variables. In particular, the need for intra-aortic balloon pump support and serum lactate concentrations >5 mmol/L proved to be serious risk factors. Using a linear discriminant analysis with 7 variables, 92.3% of patients were correctly classified (sensitivity 76.9%, specificity 93.8%).

Conclusions: A high index of suspicion for NOMI in patients with the above-mentioned risk factors may decrease the diagnostic and therapeutic delay. To identify at-risk patients the developed risk equation is a useful tool with a high specificity. (*J Thorac Cardiovasc Surg* 2013;145:1603-10)

Non-occlusive mesenteric ischemia (NOMI) is a rare but serious complication after cardiac surgery. NOMI was first described by Ende in 1958¹ as summarizing all forms of mesenteric ischemia without occlusion of the mesenteric arteries. NOMI has been generally defined as “intestinal gangrene in the presence of a patent arterial tree.”² Although incidence rates around 1% have been reported, mortality is up to 90%.^{3,4}

The exact pathophysiology of NOMI is currently not fully understood; it is assumed that the key mechanism is an extreme reduction or maldistribution of splanchnic blood flow. Intestinal ischemia results in compromised integrity of the mucosal layer with bacterial translocation, bacteremia, and the development of multiorgan failure.³

The clinical signs of NOMI—oliguria, increased serum lactate levels, decreased oxygenation, or hypotension—are unspecific. The same is true for abdominal pain, which in addition may be suppressed by analgesation.^{5,6}

Angiography is currently the only means of establishing the diagnosis; in addition it allows for intra-arterial application of vasodilators as the specific treatment. Angiography, however, is an invasive procedure that cannot be applied too liberally. In addition, it must be instituted early enough to be able to correct the vascular pathology. To minimize morbidity and mortality due to NOMI, it is thus crucial to identify patients who are at risk of developing this complication.⁷

A variety of risk factors have been proposed on the basis of retrospective investigations of heterogeneous patient cohorts. Among others, advanced age, renal failure, and different forms of shock as well as the use of an intra-aortic balloon pump (IABP) have been found to be associated with NOMI.^{3,4,7} At this time there is no prospective identification of risk factors for NOMI.

Therefore, aim of this prospective study was to identify perioperative risk factors for NOMI in a large cohort of patients undergoing elective cardiac surgery.

METHODS

The study was designed as a prospective cohort study and approved by the local ethics committee (Landesärztekammer des Saarlandes; ID: 199/09). From January 1, 2010, to March 31, 2011, all patients scheduled for

From the Department of Thoracic and Cardiovascular Surgery,^a Department of Anesthesiology, Intensive Care Medicine and Pain Medicine,^b Department of Medicine, Division of Nephrology and Hypertension,^c and Department of Diagnostic and Interventional Radiology,^d Saarland University Medical Center, Homburg/Saar, Germany.

Disclosures: Authors have nothing to disclose with regard to commercial support. Received for publication May 7, 2012; revisions received Oct 6, 2012; accepted for publication Nov 6, 2012; available ahead of print Dec 10, 2012.

Address for reprints: Hans-Joachim Schäfers, MD, Department of Thoracic and Cardiovascular Surgery, University Medical Center, University of Saarland, Kirrbergerstraße 66421, Homburg/Saar, Germany (E-mail: h-j.schaefers@uks.eu). 0022-5223/\$36.00

Copyright © 2013 by The American Association for Thoracic Surgery
http://dx.doi.org/10.1016/j.jtcvs.2012.11.022

Abbreviations and Acronyms

CABG = coronary artery bypass graft
IABP = intra-aortic balloon pump
NOMI = nonocclusive mesenteric ischemia

cardiac surgery at our institution were screened for participation in this trial. The inclusion criterion was elective cardiac surgery with cardiopulmonary bypass. Exclusion criteria were age <18 years, refusal to participate, planned off-pump surgery, urgent cardiac surgery, hemodynamic instability with emergency indication for cardiac surgery, or insufficient knowledge of the German language. Written informed consent was obtained from all patients meeting the above-mentioned criteria. The patients' demographic and perioperative data were entered in a computerized data bank in addition to the medical record chart (see [Appendix 1](#)).

Radiographic Analysis

If NOMI was suspected arterial angiography was performed. The decision to perform angiography was based on the presence of at least 2 of 4 possible clinical indicators: new onset of oliguria (urine output < 0.5 mL/kg/hour for at least 6 hours) or anuria, abdominal distention with decreased or absent bowel sounds, serum lactate levels >5.0 mmol/L or metabolic acidosis (base excess <−5 mmol/L). In accordance to the definition of cardiogenic shock,⁸ angiography was only performed in patients with a systolic blood pressure >90 mm Hg, and a cardiac index >1.8 L/minute/m².

The examinations were performed on an undercouch tube/overcouch detector angiography unit (Axiom Artis, Siemens Medical Solutions, Forchheim, Germany) with a maximum field of view of 40 cm.

In all cases the common femoral artery was accessed and a 6F sheath was inserted. The superior mesenteric artery was intubated with a 4F catheter (Cordis, Johnson & Johnson Medical Products GmbH, Vienna, Austria) with a cobra tip configuration and a digital subtraction angiography was obtained by automatic contrast medium injection (contrast medium: Imeron 300, Bracco, Milan, Italy).

All images were assessed by an experienced radiologist and an intensivist on a consensus basis. In accordance with our previously published scoring system,⁹ NOMI was diagnosed whenever the total score was more than 1 ([Table 1](#)).

Statistics

Data analysis was performed using SPSS Statistics version 19 (IBM, Ehningen, Germany). As a first step, patients with and without NOMI as confirmed by angiography were compared with respect to all variables assessed in this study (see [Appendix 1](#)). For categorical variables, χ^2 tests were performed. For continuous variables the differences in manifestation (expressed as mean $\pm$ standard deviation) between NOMI and non-NOMI patients were compared with Student *t* tests (Welch's *t* tests in case of inhomogeneous variance). As a second step, odds ratios were calculated for all categorical dichotomized variables (with 95% confidence intervals) to describe the risk of developing NOMI.

Previous risk analyses only pointed toward differences between NOMI and non-NOMI patients regarding several variables. These analyses, however, did not exceed the description of the differences. Therefore, in a third step, linear discriminant analyses as well as logistic regression analyses were performed to develop a model that separates patients at high risk for developing NOMI from patients at low risk. To derive a medically sensitive predictive model, we stepwise excluded variables with low predictive values: After exclusion of the variable with the lowest predictive value, both linear discriminant and logistic regression analysis were calculated again to see how the coefficient loadings changed. Subsequently, the variable with the lowest predictive value by now was excluded, and analyses

were iterated until the final equation was found. In addition, we partially reincluded several variables to compare different possible predictive models. This was done when variables had been excluded due to statistical reasons, but were previously reported to influence NOMI.

RESULTS

During the study period, 1163 adult patients underwent elective cardiac surgery with extracorporeal circulation. Because 298 patients refused participation in the study the final study population consisted of 865 individuals (74%). Angiography was performed in 88 patients. Of these, 78 patients (9% of the total cohort) had the typical angiographic signs of NOMI in the absence of cardiogenic shock. Ten additional patients underwent angiography based on clinical suspicion; their findings were unremarkable. At the time of angiography systolic blood pressure was 120 ± 15 mm Hg, cardiac index was 2.7 ± 0.3 L/minute/m², and central venous saturation was $69.1\% \pm 1.2\%$. Relevant outcome data are shown in [Table 2](#).

Patients with NOMI differed in many respects from patients without NOMI ([Table 3](#)). Among preoperative parameters, renal insufficiency, diuretic therapy, and age >70 years showed the highest odds ratios for postoperative NOMI.

Intraoperative risk factors with clearly increased odds ratios for postoperative NOMI were duration of operation >240 minutes and cardiopulmonary bypass time >100 minutes.

The highest odds ratios for development of NOMI were observed with postoperative variables. In particular, the need for IABP support, levosimendan therapy, transfusion of >1 unit packed red blood cells, and re-exploration for bleeding proved to be serious risk factors. In addition, serum lactate concentrations >5 mmol/L and norepinephrine support >0.1 $\mu\text{g/kg/minute}$ were associated with a clearly increased risk to develop NOMI ([Table 4](#)).

Among the calculated models, the best classification regarding medical sensitivity and practice orientation was obtained by linear discriminant analysis. The final model included 7 variables (see below; for standardized canonical coefficients see [Appendix 2](#)). The discriminant function was highly significant (Wilks' Λ , 0.59; χ^2_7 , 449.41; $P < .001$). It was calculated with the following equation:

$$\begin{aligned} d = & 3.85 \times \text{postoperative IABP support (0 = no; 1 = yes)} \\ & + 1.91 \times \text{re-exploration for bleeding (0 = no; 1 = yes)} \\ & + 1.86 \times \text{postoperative need for more than 1 unit PRBC} \\ & \quad (0 = \text{no; } 1 = \text{yes}) \\ & + 1.52 \times \text{postoperative serum lactate level } >5 \text{ mmol/l} \\ & \quad (0 = \text{no; } 1 = \text{yes}) \end{aligned}$$

TABLE 1. Specific nonocclusive mesenteric ischemia (NOMI) score system⁹

	Criteria/score	Description
Vessel morphology	0: None	SMA trunk, branches, and mesenteric arcades normal
	1: Mild	SMA trunk normal, several SMA branches and/or mesenteric arcades slightly constricted
		SMA trunk slightly constricted, normal SMA branches and mesenteric arcades
		SMA trunk, several SMA branches and/or mesenteric arcades slightly constricted
	2: Moderate	SMA trunk, multiple SMA branches, and/or mesenteric arcades constricted
Contrast medium reflux into the aorta	0: None	No contrast medium reflux into the aorta
	1: Mild	Some contrast medium reflux into the aorta
	2: Severe	Severe contrast medium reflux with complete aortogram
Contrasting of the parenchyma	0: Complete	Homogeneous contrasting of the intestine
	1: Some deficiency	Inhomogeneous contrasting of the intestine
	2: Complete deficiency	No contrasting of the intestine
Distension of the intestine	0: None	Up to 3 air containing, distended bowel loops
	1: Mild	>3 air containing, distended bowel loops
	2: Severe	Ubiquitary distended bowel loops
Contrasting time of portal vein	0: Normal	<8 sec
	1: Slightly delayed	9-12 sec
	2: Delayed	>13 sec
Total score	No	0
	Mild (Grade 1)	1-5
	Moderate (Grade 2)	6-7
	Severe (Grade 3)	8-11

Summary of a specific NOMI score system used for image analysis with respect to vessel morphology, contrast medium reflux into the aorta, contrast enhancement and distension of the intestine, as well as the time of portal vein filling. The sum of each category results into a total score. SMA, Superior mesenteric artery.

+ 1.07 × postoperative levosimendan therapy
(0 = no; 1 = yes)

+ 0.77 × postoperative need for norepinephrine
>0.1 µg/kg/min
(0 = no; 1 = yes)

+ 0.36 × postoperative loss of sinus rhythm
(0 = no; 1 = yes)

− 0.32 (constant of the equation)

TABLE 2. Comparison of relevant outcome data

Outcome	Non-NOMI		P value
	NOMI (n = 78)	(n = 787)	
Mechanical ventilation (h)	166.5 ± 23.3	22.1 ± 2.0	<.001
Renal replacement therapy (%)	42 (53.4)	17 (2.2)	<.001
LOS ICU (d)	9.3 ± 1.1	1.8 ± 0.1	<.001
LOS IMC (d)	6.1 ± 0.7	2.2 ± 0.1	<.001
LOS hospital (d)	20.2 ± 1.5	11.8 ± 0.2	<.001
30-day mortality (%)	17 (22.2)	10 (1.2)	<.001

Data are expressed as n (%) or mean ± standard deviation. Boldface indicates statistically significant P values. NOMI, Nonocclusive mesenteric ischemia; LOS, length of stay; ICU, intensive care unit; IMC, intermediate care unit.

The discrimination value was 1.18; individuals scoring higher were suspected to experience NOMI. With our model we could account for 40.7% of the variance.

In total, 92.3% of patients were correctly classified as patients with or without NOMI (Table 5). The rate of correct prediction was 76.9% for patients with NOMI (sensitivity), and 93.8% for patients without NOMI (specificity).

A parallel logistic regression model with the same independent variables as used in the prior model could account for 50.1% of variance (Nagelkerke R^2 , 0.50; model χ^2 , 223.26; $P < .001$) and allowed for 93.9% correct predictions (Table 5). The regression model was inferior in terms of sensitivity (43.6%) but superior in specificity (98.9%) to the model derived from linear discriminant analyses. Importantly, 44 patients with NOMI had been mistakenly classified as asymptomatic using the regression equation, whereas the same applied for only 18 patients using the equation obtained through discriminant analyses.

DISCUSSION

A life-threatening complication of cardiac surgical procedures is intestinal ischemia.^{3,4} The pathomechanism of NOMI is not well understood. It is known that ileum and distal jejunum are most severely affected.⁷ It is as yet unclear if NOMI is simply the delayed result of

PM

TABLE 3. Summary of statistically significant differences between patients with and without nonocclusive mesenteric ischemia (NOMI)

	Risk factor	NOMI (n = 78)	Non-NOMI (n = 787)	P value
Demographics	Age (y)	72.0 ± 1.0	62.8 ± 0.5	<.001
	Height (cm)	169.0 ± 1.1	172.3 ± 0.3	.002
	EuroSCORE	8.6 ± 0.4	5.8 ± 0.1	<.001
Comorbidity	Atrial fibrillation (%)	20 (26)	115 (14)	.009
	Coronary artery disease (%)	48 (62)	325 (41)	<.001
	Pulmonary hypertension (%)	31 (40)	174 (22)	<.001
	Chronic obstructive pulmonary disease (%)	11 (14)	53 (7)	.02
	Renal insufficiency (%)	15 (19)	48 (6)	<.001
	Status past stroke (%)	7 (9)	28 (4)	.02
	Diuretic (%)	58 (74)	346 (44)	<.001
Medication	Phenprocoumon (%)	13 (17)	68 (9)	.02
	Statin (%)	34 (44)	232 (30)	.01
	Coronary artery bypass grafting operations	42 (54)	285 (36)	.002
Intraoperative	Operation time (min)	215.6 ± 8.4	166.7 ± 1.9	<.001
	Cardiopulmonary bypass time (min)	114.0 ± 6.4	82.8 ± 1.2	<.001
	Crossclamp time (min)	71.6 ± 4.3	54.4 ± 0.8	<.001
	Intra-aortic balloon pump support (%)	13 (17)	1 (0.1)	<.001
	Sinus rhythm (%)	14 (18)	365 (46)	<.001
	Blood loss (ml)	1116.1 ± 96.7	622.2 ± 17.5	<.001
	Re-exploration for bleeding (%)	18 (23)	13 (2)	<.001
	Packed red blood cells (n)	3.1 ± 0.2	1.7 ± 0.1	<.001
	Levosimendan therapy (%)	7 (9)	4 (0.5)	<.001
	Norepinephrine (μg/kg/min)	0.26 ± 0.03	0.06 ± 0.01	<.001
	Serum lactate >10 mmol/L	11 (14)	2 (0.3)	<.001
	Serum lactate >5 mmol/L	28 (36)	29 (4)	<.001
	Base excess <-5	39 (50)	124 (16)	<.001
	pH < 7.2	2 (3)	6 (0.8)	<.001
Postoperative	Urine output within first 24 h (mL)	2151 ± 163	2698 ± 59	.002
	Urine output within first 24 h (mL/kg/h)	1.13 ± 0.08	1.41 ± 0.03	.002

Data are expressed as n (%) or mean ± standard deviation. Boldface indicates statistically significant *P* values. EuroSCORE, European System for Cardiac Operative Risk Evaluation.

microcirculatory alterations initiated during cardiopulmonary bypass (CPB) or rather the manifestation of a separate disease. Several studies have also documented alterations in intestinal microcirculation during CPB.¹⁰⁻¹³ Impaired intestinal blood flow may result in breakdown of the intestinal barrier,¹⁴ which contributes to further aggravation of the physiologic alterations after CPB. Alternatively, it may be assumed that the changes in the microcirculation are a consequence of the vasopressor treatment that is necessary in many patients during and after operations involving CPB.¹⁵

NOMI is a difficult clinical problem due to the nonspecific nature of the symptoms and difficulty in diagnosis. Early identification of patients at highest risk will allow lowering the threshold for angiography as the essential diagnostic and therapeutic tool. Therefore, we initiated this prospective study to identify isolated risk factors and patients at risk for NOMI.

Previous studies have identified advanced age, severe heart failure, cardiogenic shock, chronic renal failure, and diuretic medication as important preoperative risk factors for the development of mesenteric ischemia.^{7,16} Among these, advanced age (>70 years), chronic renal failure,

and diuretic medication increases the odds ratio to develop NOMI in our study between 3- and 4-fold. These risk factors alone do not provide conclusive evidence, but may be indicative for the susceptibility of patients to develop NOMI.

The most prominent postoperative factors described in the literature are hypovolemic shock, cardiogenic shock, administration of α -adrenergic drugs, or the use of an IABP.^{7,16} In line with these findings, extended norepinephrine use, levosimendan therapy, or postoperative IABP support increases the odds ratio to develop NOMI in our study between 18- and 153-fold.

Vasopressor treatment is necessary in many patients during and after operations involving CPB.¹⁷⁻¹⁹ In this context, treatment with α -adrenergic catecholamines like norepinephrine may lead to an increased incidence of NOMI after CPB.¹² It may be assumed that these changes in the microcirculation are a consequence of a hypercontractile response of ileal microvessels after α -adrenoceptor stimulation.^{12,15,17} In addition, norepinephrine stimulates in a dose-dependent manner β receptors, which has been shown to increase intestinal oxygen consumption.²⁰

TABLE 4. Odds ratios of perioperative risk factors

		Odds ratio	95% CI	P value
Demographics	Age >70 y	2.82	1.74-4.60	<.001
	Height >170 cm	0.51	0.32-0.82	<.05
	EuroSCORE >7	2.33	1.65-4.01	<.001
	Atrial fibrillation	2.17	1.27-3.73	<.001
Comorbidity	Coronary artery disease	2.03	1.31-3.32	<.05
	Pulmonary hypertension	2.48	1.53-4.01	<.001
	Chronic obstructive pulmonary disease	2.46	1.22-4.95	<.001
	Renal insufficiency	3.95	2.12-7.37	<.001
Medication	Status past stroke	2.60	1.10-6.16	<.001
	Diuretics	3.79	2.24-6.43	<.001
	Phenprocoumon	2.12	1.11-4.03	<.05
	Statins	1.85	1.15-2.97	<.05
Intraoperative	Coronary artery bypass grafting operations	2.01	1.29-3.28	<.05
	Operation time >240 min	5.62	3.26-9.69	<.001
	Operation time >180 min	3.28	2.02-5.31	<.001
	Cardiopulmonary bypass time >100 min	3.83	2.38-6.17	<.001
Postoperative	Crossclamp time >60 min	2.19	1.37-3.49	<.05
	Intra-aortic balloon pump support	153.00	19.70-1188.08	<.001
	Loss of sinus rhythm	3.53	2.00-6.22	<.001
	Blood loss >1000 ml	7.22	4.29-12.14	<.001
	Re-exploration for bleeding	24.22	11.64-59.42	<.001
	Packed red blood cells >2	20.05	11.39-35.27	<.001
	Packed red blood cells >1	12.77	7.66-21.27	<.001
	Levosimendan therapy	18.78	5.37-65.71	<.001
	Norepinephrine >0.2 µg/kg/min	17.47	9.01-33.88	<.001
	Norepinephrine >0.1 µg/kg/min	12.14	7.18-20.52	<.001
	Serum lactate >10 mmol/L	139.09	17.81-1086.33	<.001
	Serum lactate >5 mmol/L	15.60	8.61-28.26	<.001

Data are expressed as odds ratio for each recorded categorical respectively dichotomized variable and 95% confidence interval (CI). Boldface indicates statistically significant *P* values. *EuroSCORE*, European System for Cardiac Operative Risk Evaluation.

Irrespective of the surgical procedure, patients after cardiac surgery are still vulnerable to low cardiac output syndrome or cardiogenic shock. Pharmacologic interventions are commonly limited to catecholamines, phosphodiesterase III inhibitors, and levosimendan, which has recently received more attention.²¹ Due to the cost of levosimendan it is usually restricted to refractory cardiogenic shock. Although levosimendan therapy is well tolerated, most of its adverse effects are due to its vasodilating effect.²¹ This side effect itself and/or the resulting need for increased vasopressor treatment may additionally aggravate intestinal

TABLE 5. Comparison between linear discriminant and logistic regression analysis to predict nonocclusive mesenteric ischemia (NOMI)

Clinical diagnosis	Predicted diagnosis			
	Linear discriminant analysis		Logistic regression analysis	
	non-NOMI	NOMI	non-NOMI	NOMI
non-NOMI	738 (93.8)	49 (6.2)	778 (98.9)	9 (1.1)
NOMI	18 (23.1)	60 (76.9)	44 (56.4)	34 (43.6)

Data are presented as n (%). Boldface indicates correct predictions. Linear discriminant analysis classified 92.3% of all patients correctly, whereas logistic regression analysis showed a correct prediction rate of 93.9% (n = 865).

hypoperfusion. Nevertheless, we do not think that levosimendan itself is a risk factor for the development of NOMI, but its need in the treatment of cardiogenic shock seems to be a clue for impaired intestinal perfusion.

IABP implantation after cardiac surgery is a commonly used form of circulatory support for patients with postoperative low cardiac output syndrome. Despite improving coronary perfusion and reducing left ventricular afterload, IABP use is known as risk factor for the development of lethal mesenteric ischemia.²² One explanation for this unwanted side effect is the malposition of the IABP balloon, resulting in compromised visceral blood flow. This hypothesis is supported by Rastan and colleagues²³ who found a malposition of the IABP balloon with consecutive visceral artery compromise in up to 97% of their study patients. After insertion of the IABP, the position was immediately confirmed by chest x-ray film. At angiography we only found 3 patients (23%) with a malposition of the IABP balloon, which was then corrected. Therefore, an incorrect-positioned IABP balloon may aggravate a state of intestinal hypoperfusion, but does not explain alone the >150-fold odds increase for the development of NOMI in our patients. Moreover, NOMI was confirmed after restoration of an adequate hemodynamic, underlining that NOMI is a distinct complication after cardiogenic shock. Therefore, despite an apparently re-established stable hemodynamic situation, NOMI should be kept in mind as a life-threatening complication.

Intestinal ischemia may lead to a systemic inflammatory response syndrome and sepsis thereby altering the coagulation system in both ways, hyper- and hypocoagulation.²⁴ Bleeding is additionally known to be a strong proinflammatory stimulus, which further aggravates systemic inflammatory response syndrome and sepsis.²⁴ A *circulus vitiosus* that ends up in distinct NOMI may be the result. This is supported by higher odds ratios to develop NOMI in patients with increased blood loss, need for >1 unit packed red blood cells, or need for chest re-exploration. In 15 of 18 patients, however, no surgical source of bleeding was found while re-exploring the chest, suggesting coagulopathy caused by NOMI.

Elevated serum lactate levels, metabolic acidosis, and increased serum creatinine kinase levels have been described

in previous studies to be indicative but not providing conclusive evidence for NOMI.⁴ We found significantly higher serum lactate levels in patients with NOMI compared with patients without. Furthermore, patients with serum lactate levels of >5 mmol/L had an increased odds ratio of 15.60 to show the angiographic signs of NOMI. This may be due to an increased intestinal lactate production as a result of anaerobic glycolysis. It is, however, impossible to discriminate between lactate production due to systemic or local hypoperfusion.

All these postoperative predictors have to be looked at in the context of preoperative morbidity and intraoperative events. Especially intraoperative physiology is difficult or impossible to define clearly; myocardial protection, suboptimal correction of cardiovascular pathology, and hemodynamic derangement cannot be quantified in clinical practice and may have effects that extend into postoperative treatment.

Despite the possibility to detect intestinal malperfusion by contrast multidetector row computed tomography,²⁵ catheter angiography remains the gold standard for the early diagnosis of NOMI. In addition to the direct visualization of the mesenteric blood flow, the inserted catheter allows selective mesenteric intra-arterial infusion of vasodilative drugs. Early recognition and correction of the vascular pathology is the only way to reduce the high morbidity and mortality of NOMI.⁷ Nevertheless, angiography is an invasive procedure that cannot be applied too liberally due to its potential side effects. To alleviate the decision process to perform angiography, we tried to identify patients who are at risk to develop NOMI by using a discriminant and regression approach, respectively. The final risk equation showed a good sensitivity and high specificity. Using this equation, we were able to exclude NOMI in >90% of cases. As a result the sensitivity was only 76%; this reflects our key intention to aim for a high specificity. In our mind the high mortality of untreated NOMI warrants a test that has the lowest possible false negative rate.

Linear discriminant analysis may be discussed controversially because this method is meant for the analysis of continuous independent variables. The use of supplemental categorical variables, in the form of dummy variables, is allowed. In our predictive model all continuous variables had to be eliminated because they showed no explanatory value. Because discriminant analysis is not designated for exclusively dichotomous data, our equation model may be problematic for statistical reasons. From a clinical perspective, the results obtained through linear discriminant analysis are superior to the results obtained through logistic regression analysis. In the first case, the predictive model had a false negative rate of 23.1% (n = 18 [2.1% of the total cohort]), whereas the predictive model derived from logistic regression analysis had a rate of 56.4% (n = 44 [5.1% of the total cohort]). Because false negative NOMI predictions

are associated with high mortality rates, type II error is much more important than sensitivity rate.

Another limitation of our study is that our control group (non-NOMI patients) did not undergo angiography. Therefore we cannot exclude that patients with clinical unapparent forms of NOMI were missed. The mortality in our control group was low (1.2%); in addition, none of these patients died within the first 10 postoperative days. These facts and the clinical course of the patients makes NOMI very unlikely.

Emergency operations are associated with a higher incidence of NOMI.²⁶ Due to the study design we excluded all urgent and emergency operations. Therefore, we cannot comment on whether or not these patients are at higher risk for the development of NOMI. Nevertheless, our risk equation consists of pre- and postoperative variables, which helps to easily identify risk patients independent of the operation indication.

CONCLUSIONS

Our study provides for the first time prospective observational data on risk factors for the development of NOMI in patients undergoing elective cardiac surgery. Summarizing our results, a high index of suspicion for NOMI in patients with the above-mentioned risk factors may decrease the diagnostic delay. This may lead to a decreased mortality due to earlier intervention. To better identify risk patients the developed risk equation is a useful tool with a high specificity and low type II error rate.

The authors thank Timo Speer for his support in database maintenance.

References

- Ende N. Infarction of the bowel in cardiac failure. *N Engl J Med*. 1958;258: 879-81.
- Haglund U, Hultén L, Ahren C, Lundgren O. Mucosal lesions in the human small intestine in shock. *Gut*. 1975;16:979-84.
- Klotz S, Vestring T, Rotker J, Schmidt C, Scheld HH, Schmid C. Diagnosis and treatment of nonocclusive mesenteric ischemia after open heart surgery. *Ann Thorac Surg*. 2001;72:1583-6.
- Abbond B, Daher R, Boujaoude J. Acute mesenteric ischemia after cardiopulmonary bypass surgery. *World J Gastroenterol*. 2008;14:5361-70.
- Howard TJ, Plaskon LA, Wiebke EA, Wilcox MG, Madura JA. Nonocclusive mesenteric ischemia remains a diagnostic dilemma. *Am J Surg*. 1996;171:405-8.
- Bassiouny HS. Nonocclusive mesenteric ischemia. *Surg Clin North Am*. 1997; 77:319-26.
- Trompeter M, Brazda T, Remy CT, Vestring T, Reimer P. Non-occlusive mesenteric ischemia: etiology, diagnosis, and interventional therapy. *Eur Radiol*. 2002; 12:1179-87.
- Van de Werf F, Bax J, Betriu A, Blomstrom-Lundqvist C, Crea F, Falk V, et al. Management of acute myocardial infarction in patients presenting with persistent ST-segment elevation: the Task Force on the Management of ST-Segment Elevation Acute Myocardial Infarction of the European Society of Cardiology. *Eur Heart J*. 2008;29:2909-45.
- Minko P, Groesdonk H, Stroeder J, Miodek J, Graeber S, Buckner A, et al. A scoring system for the assessment of angiographic findings in non-occlusive mesenteric ischemia (NOMI). *Rof*. 2012;184:805-9.
- Tao W, Zwischenberger JB, Nguyen TT, Vertrees RA, McDaniel LB, Nutt LK, et al. Gut mucosal ischemia during normothermic cardiopulmonary bypass

results from blood flow redistribution and increased oxygen demand. *J Thorac Cardiovasc Surg.* 1995;110:819-28.

11. Doguet F, Litzler PY, Tamion F, Richard V, Hellot MF, Thuillez C, et al. Changes in mesenteric vascular reactivity and inflammatory response after cardiopulmonary bypass in a rat model. *Ann Thorac Surg.* 2004;77:2130-7; author reply 2137.

12. O'Dwyer C, Woodson LC, Conroy BP, Lin CY, Deyo DJ, Uchida T, et al. Regional perfusion abnormalities with phenylephrine during normothermic bypass. *Ann Thorac Surg.* 1997;63:728-35.

13. Bombberg H, Bierbach B, Flache S, Wagner I, Glaser L, Groesdonk HV, et al. Endothelin and vasopressin influence splanchnic blood flow distribution during and after cardiopulmonary bypass. *J Thorac Cardiovasc Surg.* 2012 Apr 30 [Epub ahead of print].

14. Rocke DA, Gaffin SL, Wells MT, Koen Y, Brock Utine JG. Endotoxemia associated with cardiopulmonary bypass. *J Thorac Cardiovasc Surg.* 1987; 93:832-7.

15. Tofukuji M, Stahl GL, Metais C, Tomita M, Agah A, Bianchi C, et al. Mesenteric dysfunction after cardiopulmonary bypass: role of complement C5a. *Ann Thorac Surg.* 2000;69:799-807.

16. Lock G. Acute intestinal ischaemia. *Best Prac Res Clin Gastroenterol.* 2001;15: 83-98.

17. Khan TA, Bianchi C, Ruel M, Feng J, Sellke FW. Differential effects on the mesenteric microcirculatory response to vasopressin and phenylephrine after cardiopulmonary bypass. *J Thorac Cardiovasc Surg.* 2007;133:682-8.

18. Argenziano M, Chen JM, Choudhri AF, Cullinane S, Garfein E, Weinberg AD, et al. Management of vasodilatory shock after cardiac surgery: identification of predisposing factors and use of a novel pressor agent. *J Thorac Cardiovasc Surg.* 1998;116:973-80.

19. Morales DL, Gregg D, Helman DN, Williams MR, Naka Y, Landry DW, et al. Arginine vasopressin in the treatment of 50 patients with postcardiotomy vasodilatory shock. *Ann Thorac Surg.* 2000;69:102-6.

20. Salak N, Pajk W, Knotzer H, Hofstotter H, Schwarz B, Mayr A, et al. Effects of epinephrine on intestinal oxygen supply and mucosal tissue oxygen tension in pigs. *Crit Care Med.* 2001;29:367-73.

21. Mebazaa A, Pitsis AA, Rudiger A, Toller W, Longrois D, Ricksten SE, et al. Clinical review: practical recommendations on the management of perioperative heart failure in cardiac surgery. *Crit Care.* 2010;14:201.

22. Venkateswaran RV, Charman SC, Goddard M, Large SR. Lethal mesenteric ischaemia after cardiopulmonary bypass: a common complication? *Eur J Cardiothorac Surg.* 2002;22:534-8.

23. Rastan AJ, Tillmann E, Subramanian S, Lehmkuhl L, Funkat AK, Leontyev S, et al. Visceral arterial compromise during intra-aortic balloon counterpulsation therapy. *Circulation.* 2010;122(11 suppl):S92-9.

24. van der Poll T, de Boer JD, Levi M. The effect of inflammation on coagulation and vice versa. *Curr Opin Infect Dis.* 2011;24:273-8.

25. Mitsuyoshi A, Obama K, Shinkura N, Ito T, Zaima M. Survival in nonocclusive mesenteric ischemia: early diagnosis by multidetector row computed tomography and early treatment with continuous intravenous high-dose prostaglandin E(1). *Ann Surg.* 2007;246:229-35.

26. Allen KB, Salam AA, Lumsden AB. Acute mesenteric ischemia after cardiopulmonary bypass. *J Vasc Surg.* 1992;16:391-5; discussion 395-6.

APPENDIX 1. Summary of all documented data analyzed in this study

Data		Description
Demographics	Sex	
	Girls	Sex of each patient
	Boys	Sex of each patient
	Age	Age, documented in years (y)
	Weight	Body weight, documented in kilograms
	Size	Size, documented in centimeters
	EuroSCORE	EuroSCORE, documented in absolute value
	Ejection fraction	Ejection fraction, documented in percent (%)
	New York Heart Association functional class	New York Heart Association functional class, documented in absolute value
	Sinus rhythm	Documented sinus rhythm at the day of admission to the hospital
	Arterial hypertension	Systolic arterial pressure >140 mm Hg and/or diastolic arterial pressure >90 mm Hg
	Karnofsky Index	Karnofsky Index, documented in absolute value
	Smoker	Active smoker at the day of admission to the hospital
Comorbidity	Pack years	No. of pack years = packs smoked/d × years as a smoker
	Coronary artery disease	Confirmed by cardiac catheter examination
	Vitium	All kinds of valvular disease with indication for surgery
	Chronic pulmonary thromboembolism	Confirmed by angiography
	Pulmonary hypertension	Mean pulmonary artery pressure >25 mm Hg at rest or >30 mm Hg with exercise
	Chronic obstructive pulmonary disease	Global Initiative for Chronic Obstructive Lung Disease ≥2
	Renal insufficiency	Effective glomerular filtration rate <60 mL/min
	Hemodialysis	Terminal renal failure with the need for dialysis
Medication	Malignancy	All kinds of malignant disease
	Status past stroke	Status past stroke with or without residua
	Beta blocker	All kinds of beta blocker therapy for at least 4 wk before operation
	ACE inhibitor	All kinds of ACE inhibitor therapy for at least 4 wk before operation
	AT-1 blocker	All kinds of AT-1 blocker therapy for at least 4 wk before operation
	Calcium antagonist	All kinds of calcium antagonist therapy for at least 4 wk before operation
	Diuretic	All kinds of diuretic therapy for at least 4 wk before operation
	Acetylsalicylic acid	All kinds of acetylsalicylic acid therapy for at least 4 wk before operation
	Phenprocoumon	All kinds of phenprocoumon therapy for at least 4 wk before operation
	Amiodarone	All kinds of amiodarone therapy for at least 4 wk before operation
	Statin	All kinds of statin therapy for at least 4 wk before operation

(Continued)

APPENDIX 1. Continued

	Data	Description
Intraoperative	Procedure type	
	CABG operations	All kinds of coronary artery bypass surgery with cardiopulmonary bypass
	Valve operations	All kinds of valve surgery with cardiopulmonary bypass
	Pulmonary thromboendarterectomy	Pulmonary thromboendarterectomy
	Combination surgery	All kinds of combination surgery with cardiopulmonary bypass
	Redo CABG and/or valve	All kinds of redo surgery with cardiopulmonary bypass
	Operation time	Operation time, documented in minutes
	CPB time	Cardiopulmonary bypass time, documented in minutes
	Crossclamp time	Aortic crossclamp time, documented in minutes
Postoperative	HCA time	HCA time, documented in minutes
	IABP support	Postoperative need for IABP support
	Loss of sinus rhythm	Documented loss of sinus rhythm for at least 6 h
	Blood loss	Blood loss, documented in milliliters
	Re-exploration for bleeding	Need for re-exploration of the chest due to bleeding
	Packed red blood cells	No. of packed red blood cells
	Levosimendan therapy	Postoperative need for levosimendan therapy
	Norepinephrine therapy	Postoperative need for norepinephrine therapy, documented in micrograms/kilogram/minute
	Serum lactate	Postoperative serum lactate levels
	Base excess	Postoperative base excess values
	pH value	Postoperative pH values
	Urine output within first 24 h	Postoperative urine output, documented in milliliters
	Urine output within first 24 h (ml/kg/h)	Postoperative urine output, documented in milliliters/kilogram/hour
	Mechanical ventilation	Time of postoperative mechanical ventilation, documented in hours
Outcome	Renal replacement therapy	Need for renal replacement therapy due to new onset of acute kidney injury
	LOS ICU	LOS in the ICU, documented in days
	LOS IMC	LOS in the IMC unit, documented in days
	LOS hospital	LOS in the hospital, documented in days
	30-d mortality	30-d mortality including all causes of death

In total 52 isolated risk factors were included in our risk stratification calculations. *ACE*, Angiotensin-converting enzyme; *AT-1*, angiotensin-1; *CABG*, coronary artery bypass grafting; *CPB*, cardiopulmonary bypass; *HCA*, hypothermic circulatory arrest; *IABP*, intra-aortic balloon pump; *LOS*, length of stay; *ICU*, intensive care unit; *IMC*, intermediate care unit; *EuroSCORE*, European System for Cardiac Operative Risk Evaluation.

APPENDIX 2. Summary of stepwise discriminant analysis

Predictor	Standardized coefficient loadings
Postoperative IABP support	0.451
Re-exploration for bleeding	0.348
Postoperative need for more than 2 PRBC	0.456
Postoperative serum lactate level >5 mmol/L	0.351
Postoperative need for norepinephrine >0.1 µg/kg/min	0.221
Postoperative levosimendan therapy	0.117
Postoperative loss of sinus rhythm	0.179
Canonical correlation	0.638
Eigenvalue	0.687
Wilks' lambda	0.593
χ^2	449.41 (<i>df</i> = 7; <i>P</i> < .001)

IABP, Intra-aortic balloon pump; *PRBC*, packed red blood cells; *df*, degrees of freedom.