University of Sydney
Royal Prince Alfred Hospital
Sydney, Australia

References

1. Cao C, Manganas C, Horton M, Bannon P, Munkholm-Larsen S, Ang SC, et al. Angiographic outcomes of radial artery versus saphenous vein in coronary artery bypass graft surgery: a meta-analysis of randomized controlled trials. *J Thorac Cardiovasc Surg.* 2012 Aug 4. [Epub ahead of print].
2. Souza DS, Johansson B, Bojo L, Karlsson R, Geijer H, Filbey D, et al. Harvesting the saphenous vein with surrounding tissue for CABG provides long-term graft patency comparable to the left internal thoracic artery: results of a randomized longitudinal trial. *J Thorac Cardiovasc Surg.* 2006;132:373-8.
3. Lopes RD, Hafley GE, Allen KB, Ferguson TB, Peterson ED, Harrington RA, et al. Endoscopic versus open vein-graft harvesting in coronary-artery bypass surgery. *N Engl J Med.* 2009;361:235-44.
4. Davis Z, Jacobs HK, Zhang M, Thomas C, Castellanos Y. Endoscopic vein harvest for coronary artery bypass grafting: technique and outcomes. *J Thorac Cardiovasc Surg.* 1998;116:228-35.

<http://dx.doi.org/10.1016/j.jtcvs.2012.12.078>

CLINICAL IMPLICATIONS RELATED TO PREOPERATIVE DETECTION OF STAGE IA LUNG ADENOCARCINOMA To the Editor:

We read with interest the article by Tsutani and colleagues.¹ They aimed to determine clinical predictors of nodal involvement in stage IA lung adenocarcinoma and successfully identified that tumor size <0.8 cm and maximum standardized uptake value <1.5 are predictive for stage N0 status. We would like to discuss the following interesting items.

One of the most important indications for sublobar resection is the absence of nodal disease (ie, stage N0), which is usually preoperatively investigated by computed tomography (CT) and positron emission tomography (PET)–CT scan.² A more accurate evaluation of nodal status can be obtained by intraoperative nodal dissection with frozen section, as reported by the authors.¹ However, the intraoperative evaluation of every hilar and mediastinal nodal station appears to be too difficult to be routinely performed in

clinical practice. To avoid this, Tsutani and colleagues identified 2 factors statistically significant for preoperative prediction of N0 status. In particular, it is remarkable that these 2 parameters are easily achievable by the chest CT and PET–CT scan that are always performed when staging the disease of every patient with lung cancer.

We work daily with patients who should be excluded from surgery because their poor general health conditions or advanced age strongly advise against lobectomy. Tsutani and colleagues' findings should be particularly useful with these patients because they allow clinicians to preoperatively determine patients in which lobectomy can be avoided and sublobar resection with nodal sampling is enough. This increases the number of candidates for surgery.

Another item that we would like to underline is that the chance to preoperatively determine N0 status and therefore to plan a sublobar resection allows surgeons to understand the proper access to adopt. In particular, the possibility of performing a wedge resection with nodal sampling should increase the chance to perform video-assisted thoracic surgery,³ increasing once again the number of candidates for surgery.

One more interesting point is that tumors determined to be <0.8 cm or with maximum standardized uptake value <1.5 are characterized by a minor pathologic invasiveness and therefore are related to a better disease-free survival after sublobar resection.

Finally, we have 1 suggestion regarding the technique used in PET–CT scan for lesions affected by physiologic motion as pulmonary nodules. We suggest the use of respiratory gating⁴ because this method results in lesion volumes closer to those assessed by CT and improves measurements of tracer uptake.

Alessandro Baisi, MD^a
Matilde De Simone, MD, PhD^b
Ugo Cioffi, MD, PhD^b
^aThoracic Surgery Unit
Azienda Ospedaliera San Paolo

Milano
University of Milan
Milan, Italy

^bDepartment of Surgery
Fondazione IRCCS Ca' Granda
Ospedale Maggiore Policlinico
Milano
University of Milan
Milan, Italy

References

1. Tsutani Y, Miyata Y, Nakayama H, Okumura S, Adachi S, Yoshimura M, et al. Prediction of pathologic node-negative clinical stage IA lung adenocarcinoma for optimal candidates undergoing sublobar resection. *J Thorac Cardiovasc Surg.* 2012;144:1365-71.
2. Okada M, Koike T, Higashiyama M, Yamato Y, Kodama K, Tsubota N. Radical sublobar resection for small-sized non-small lung cancer: a multicenter study. *J Thorac Cardiovasc Surg.* 2006;132:769-75.
3. Sienel W, Dango S, Kirschbaum A, Cucuruz B, Horth W, Stremmel C, et al. Sublobar resections in stage IA non-small cell lung cancer: segmentectomies result in significantly better cancer-related survival than wedge resections. *Eur J Cardiothorac Surg.* 2008;33:728-34.
4. Werner MK, Parker JA, Kolodny GM, English JR, Palmer MR. Respiratory gating enhances imaging of pulmonary nodules and measurement of tracer uptake in FDG PET/CT. *Am J Roentgenol.* 2009;193:1640-5.

<http://dx.doi.org/10.1016/j.jtcvs.2012.12.082>

Reply to the Editor:

We are grateful for the letter by Baisi and colleagues regarding our study,¹ and we are delighted by their thoughtful insights into our results. We appreciate their viewpoint that 2 parameters (solid tumor size <0.8 cm on high-resolution computed tomography [HRCT] or a maximum standardized uptake volume [SUVmax] <1.5 on 18F-fluorodeoxyglucose positron emission tomography–computed tomography [PET–CT]) for predicting no nodal metastasis are easily achievable by the chest CT and PET–CT scan that are always performed in staging the disease of every patient with lung cancer.²

Generally, clinical physicians measure tumor size according to the TNM Classification of Malignant Tumors (TNM) by including the ground-glass opacity (GGO) components visualized on HRCT. We have found^{3,4} that

in addition to the SUVmax found on PET-CT,² the predictive values of the solid tumor size found on HRCT for pathologic high-grade malignancy and prognosis in patients with clinical stage IA lung adenocarcinoma are greater than those of whole tumor size, including solid and GGO components. We strongly recommend that solid tumor size be used to determine the T descriptor in the TNM classification of lung cancer and should be defined as the true tumor size in cases of lung adenocarcinoma with a GGO component visualized on HRCT.

The suggestion regarding a technique in PET-CT scan for lung lesions affected by physiologic motion is justified. The difference between the CT volume and the gated PET volume is significantly less than the difference between the CT volume and the ungated PET volume.⁵ Respiratory gating has the potential to enhance imaging of organs in motion, such as lungs, and although some additional time is needed for setup and acquisition, it can be incorporated practically and efficiently into clinical routine. Alternatively, breath-holding might be used during diagnostic PET-CT if true respiratory gating is not available.

Our belief, based on our data, is that the optimal surgical strategy for cT1N0M0 adenocarcinomas can be chosen using the solid tumor size found on HRCT and an SUVmax found on 18F-fluorodeoxyglucose PET-CT.

Morihito Okada, MD, PhD

Yasuhiro Tsutani, MD, PhD

Yoshihiro Miyata, MD, PhD

Department of Surgical Oncology

Hiroshima University

Hiroshima, Japan

References

1. Tsutani Y, Miyata Y, Nakayama H, Okumura S, Adachi S, Yoshimura M, et al. Prediction of pathologic node-negative clinical stage IA lung adenocarcinoma for optimal candidates undergoing sublobar resection. *J Thorac Cardiovasc Surg.* 2012;144:1365-71.
2. Okada M, Nakayama H, Okumura S, Daisaki H, Adachi S, Yoshimura M, et al. Multicenter analysis of high-resolution computed tomography and posi-

tron emission tomography/computed tomography findings to choose therapeutic strategies for clinical stage IA lung adenocarcinoma. *J Thorac Cardiovasc Surg.* 2011;141:1384-91.

3. Tsutani Y, Miyata Y, Nakayama H, Okumura S, Adachi S, Yoshimura M, et al. Prognostic significance of using solid versus whole tumor size on high-resolution computed tomography for predicting pathologic malignant grade of tumors in clinical stage IA lung adenocarcinoma: a multicenter study. *J Thorac Cardiovasc Surg.* 2012;143:607-12.
4. Tsutani Y, Miyata Y, Nakayama H, Okumura S, Adachi S, Yoshimura M, et al. Solid tumors versus mixed tumors with a ground-glass opacity component in patients with clinical stage IA lung adenocarcinoma: prognostic comparison using high-resolution computed tomography findings. *J Thorac Cardiovasc Surg.* December 14, 2012 [Epub ahead of print].
5. Werner MK, Parker JA, Kolodny GM, English JR, Palmer MR. Respiratory gating enhances imaging of pulmonary nodules and measurement of tracer uptake in FDG-PET/CT. *Am J Roentgenol.* 2009;193:1640-5.

<http://dx.doi.org/10.1016/j.jtcvs.2012.12.080>

NEW STANDARD IN LUNG CANCER STAGING? A WORD OF WARNING

To the Editor:

We read with interest the recent article by Yendamuri and Demmy¹ concerning the field of our long-standing scientific interest. We are glad that they share our fascination with the transcervical approach to the mediastinum; however, fascination with technique should not let us turn a blind eye to other, perhaps more important, aspects of surgery.

As the Yendamuri and Demmy¹ stated, transcervical extended mediastinal lymphadenectomy (TEMLA) is only used in a very few centers. We therefore draw the attention of the surgical community to some important issues missed by the Yendamuri and Demmy.¹

Discussing complications of TEMLA, Yendamuri and Demmy¹ only mentioned the recurrent laryngeal nerve palsy in 2.3% of patients. This statement may yield a false impression of safety. In fact, the rate of complications is high: in the original article on 83 patients,² it was 13.2% and apart

from recurrent nerve palsy included respiratory insufficiency, arrhythmia, pneumothorax, and deterioration in mental status. What is even more important, general fitness of 21% of patients deteriorated after TEMLA, to the degree of making them medically unfit for operation, although all were initially considered surgical candidates.²

The next article, analyzing results in 256 patients, showed a complication rate of 11.3% with a mortality as high as 1.2%. Again, more than 21% of the 176 node-negative patients became unfit for surgery because of deterioration of their general condition.³

In an updated series of 587 patients published 3 years later, the mortality was 0.9%.⁴

Complications of TEMLA may be severe. Apart from the recurrent nerve palsy mentioned by Yendamuri and Demmy,¹ they include perforation of the main bronchus, superior vena caval laceration, left phrenic nerve palsy, and injury to the innominate artery.

The least important drawback of the TEMLA is its operative time, which may be as long as 161 minutes.

The readers should be aware that these results were achieved in a very experienced center, by surgeons especially devoted to this technique. The complication rate in less experienced hands is likely to be even higher.

The first question to be answered is whether it is acceptable to use a staging procedure that is associated with mortality comparable to pulmonary lobectomy. The second, equally important, question is whether it is acceptable to use a staging procedure that makes 21% of patients unfit for curative-intent surgery. The obvious answer to both these questions is no.

A different consideration is the use of transcervical lymph node dissection as an element of curative-intent pulmonary resection. If extended mediastinal lymphadenectomy does influence survival, this approach would be interesting. At present, there are no published data in this regard. Also, data regarding