predict outcomes in infants undergoing cardiac surgery in a lesion-dependent fashion. *J Thorac Cardiovasc Surg*. Epub 2012 Aug 20.

2. Cantinotti M, Lorenzoni V, Storti S, Moschetti R, Murzi B, Marotta M, et al. Thyroid and brain natriuretic peptide response in children undergoing cardiac surgery for congenital heart disease. *Circ J*. 2012;77:188-97.
3. Lacour-Gayet F, Clarke D, Jacobs J, Comas J, Daebritz S, Daenen W, et al. The Aristotle score: a complexity-adjusted method to evaluate surgical results. *Eur J Cardiothorac Surg*. 2004;25:911-24.
4. Simsic JM, Cuadrado A, Kirshbom PM, Kanter KR. Risk adjustment for congenital heart surgery (RACHS): is it useful in a single-center series of newborns as a predictor of outcome in a high-risk population? *Congenit Heart Dis*. 2006;1:148-51.
5. Cantinotti M, Passino C, Storti S, Ripoli A, Zyw L, Clerico A. Clinical relevance of time course of BNP levels in neonates with congenital heart diseases. *Clin Chim Acta*. 2011;412:2300-4.

<http://dx.doi.org/10.1016/j.jtcvs.2012.12.086>

Reply to the Editor:

We appreciate the thoughtful comments that Cantinotti and colleagues expressed in their letter. Cantinotti and colleagues are to be commended for their important body of work, which has advanced the understanding of B-type natriuretic peptide (BNP) levels in pediatric patients with congenital heart disease.¹

We agree that levels of BNP are age dependent both in healthy neonates and in those with congenital heart disease. Also, as we indicated in the recent article under discussion,² the rapid changes in BNP levels after surgical intervention (within hours of the operation) imply that there are physiologic changes associated with the specific surgical interventions because of the particular underlying cardiac defects that determine the release of BNP. Indeed, neonates with univentricular physiology have very high preoperative BNP levels relative to those of age-matched healthy neonates and also relative to those with left-to-right shunts. In our study, there was also a trend toward higher BNP levels in neonates with univentricular physiology relative to those undergoing an arterial switch operation. We believe this is one reason that studies grouping neonates into single cohorts,^{3,4} without consideration

for the specific cardiac defects, have not found the postoperative decreases in BNP that both we² and Cantinotti and colleagues¹ found.

We therefore caution against relying on disease severity scoring systems that group diverse cardiac defects within particular categories or combine various diagnostic criteria, such as the Risk Adjustment for Congenital Heart Surgery 1 and Aristotle Basic Complexity scores, for the interpretation of perioperative BNP levels. Indeed, these scoring systems do not fully account for the variability in outcomes seen, demonstrating that other factors remain important.⁵ In attempting to understand better the physiologic release of a biomarker and its implications for prognostication and goal-directed therapy, the use of such scoring systems is therefore likely to fall short in accounting for the variability in levels of the biomarker. In fact, in Figure 1 of the letter of Cantinotti and colleagues, neonates in Aristotle Basic Complexity categories II and III can be seen to have more variability in BNP levels than those in category IV, probably because of the differences in cardiac lesions and the physiologic changes associated with surgery that inherently differ by category in these systems. A further example of the specificity we described in our study was shown in preliminary data from our previous work, where we found that *preoperative* BNP level in patients undergoing a total cavopulmonary connection (Fontan procedure) was the only significant predictor of adverse postoperative outcomes.⁶

There is clearly much to learn about the mechanisms involved in BNP release in patients with congenital heart disease and the utility of this biomarker in this population. We agree with Cantinotti and colleagues that further investigations into BNP changes during the perioperative period are warranted. On the basis of our combined data to date, we believe that fully understanding and interpreting perioperative BNP changes will require accounting for age, the cardiac lesion, and the

intervention that is undertaken. We thank both the Editor and Cantinotti and colleagues for this opportunity to highlight this important topic.

Rambod Amirmovin, MD^{a,b}

Peter Oishi, MD^{a,b}

Jeffrey R. Fineman, MD^{a,b}

Roberta L. Keller, MD^a

^aDepartment of Pediatrics

^bCardiovascular Research Institute

University of California

San Francisco, Calif

References

1. Cantinotti M, Passino C, Storti S, Ripoli A, Zyw L, Clerico A. Clinical relevance of time course of BNP levels in neonates with congenital heart diseases. *Clin Chim Acta*. 2011;412:2300-4.
2. Amirmovin R, Keller RL, Herrera C, Hsu JH, Datar S, Karl TR, et al. B-type natriuretic peptide levels predict outcomes in infants undergoing cardiac surgery in a lesion-dependent fashion. *J Thorac Cardiovasc Surg*. Epub 2012 Aug 20.
3. Niedner MF, Foley JL, Riffenburgh RH, Bichell DP, Peterson BM, Rodarte A. B-type natriuretic peptide: perioperative patterns in congenital heart disease. *Congenit Heart Dis*. 2010;5:243-55.
4. Hsu JH, Keller RL, Chikovani O, Cheng H, Hollander SA, Karl TR, et al. B-type natriuretic peptide levels predict outcome after neonatal cardiac surgery. *J Thorac Cardiovasc Surg*. 2007;134:939-45.
5. Al-Radi OO, Harrell FE Jr, Caldarone CA, McCrindle BW, Jacobs JP, Williams MG, et al. Case complexity scores in congenital heart surgery: a comparative study of the Aristotle Basic Complexity score and the Risk Adjustment in Congenital Heart Surgery (RACHS-1) system. *J Thorac Cardiovasc Surg*. 2007;133:865-75.
6. Hsu JH, Oishi PE, Keller RL, Chikovani O, Karl TR, Azakie A, et al. Perioperative B-type natriuretic peptide levels predict outcome after bidirectional cavopulmonary anastomosis and total cavopulmonary connection. *J Thorac Cardiovasc Surg*. 2008;135:746-53.

<http://dx.doi.org/10.1016/j.jtcvs.2013.01.002>

TRACHEAL REGENERATION: MYTH OR FACT?

To the Editor:

We read with great interest the article entitled, "Tracheal regeneration: Evidence of bone marrow mesenchymal stem cell involvement," by Seguin and coworkers.¹ We congratulate Seguin and coworkers¹ for their convincing demonstration of bone marrow-derived stem cell migration to the grafted area after tracheal replacement