improvement in the 15-year survival in women with radial artery (RA) grafting. We congratulate Dimitrova and coworkers on their well-designed analysis and the results in their publication. They concluded that women who received 2 arterial conduits, coronary artery bypass grafting (CABG) with left internal thoracic artery and RA, had a significant survival benefit up to 15 years compared with women who received a saphenous vein (SV) as a second conduit for myocardial revascularization.

Use of the RA as the second conduit of choice for CABG has several advantages: the length of the artery per se (~20 cm), its large diameter, its exposure to arterial blood pressure, the thickness of the wall, and its lower susceptibility to atherogenesis. Clinical studies have documented the superior long-term patency of the RA over SV grafts.²

It would be interesting to compare the survival of men undergoing CABG with the left internal thoracic artery and RA with those who received SV grafting. There may be a difference in survival between men and women using the RA as a second conduit.

Study results have shown enhanced physiologic characteristics related to endothelial production or vessel relaxation in response to nitric oxide in the RA compared with the SV.³ Joannides and colleagues⁴ reported the influence of gender difference on RA dimension in flow-dependent dilatation. Men have a higher diameter of the RA compared with women. Herman and coworkers⁵ have shown a deleterious effect of physiologic levels of androgen hormone in men and a decrease of androgen levels related to age as a benefit in coronary artery disease and endothelium function in developing the atherogenic process.

Majmudar and colleagues⁶ showed the effects of estrogen in increasing nitric oxide release in the arterial vasculature. Estrogen has its own cardioprotective effect, elevating high-density lipoprotein, lowering

low-density lipoprotein, reducing fibrinogen, maintaining the vasculature tone, inhibiting platelet aggregation and adhesion, and inhibiting the development of atherosclerotic plaques. This study was specifically targeted to forearm and arm arterial vascular resistance.

We can speculate that estrogen hormone in the premenopausal woman plays a positive role in the modulation of endothelium flow-mediated dilatation of the RA. On the other hand, men who receive RA conduits should have better results when they are aged more than 60 years because of diminishing androgen hormone and its deleterious effect on the arterial wall.⁵

These complex gender- and age-specific considerations should be taken into account in future studies of arterial conduits for revascularization.

Andrea Ruzza, MD, PhD

Lawrence S. C. Czer, MD

Division of Cardiothoracic Surgery

Cedars Sinai Medical Center

Los Angeles, Calif

References

1. Dimitrova KR, Hoffman DM, Geller CM, Ko W, Lucido DJ, Dincheva GR, et al. Radial artery grafting in women improves 15-year survival. *J Thorac Cardiovasc Surg*. 2012 [Epub ahead of print].
2. Goldman S, Sethi GK, Holman W, Thai H, McFalls E, Ward HB, et al. Radial artery grafts vs saphenous vein grafts in coronary artery bypass surgery: a randomized trial. *JAMA*. 2011;305:167-74.
3. Shapira OM, Xu A, Aldea GS, Vita JA, Shemin RJ, Keaney JF Jr. Enhanced nitric oxide-mediated vascular relaxation in radial artery compared with internal mammary artery or saphenous vein. *Circulation*. 1999;100(19 Suppl):II322-7.
4. Joannides R, Costentin A, Iacob M, Compagnon P, Lahary A, Thuillez C. Influence of vascular dimension on gender difference in flow-dependent dilatation of peripheral conduit arteries. *Am J Physiol Heart Circ Physiol*. 2002;282:H1262-9.
5. Herman SM, Robinson JT, McCredie RJ, Adams MR, Boyer MJ, Celermajer DS. Androgen deprivation is associated with enhanced endothelium-dependent dilatation in adult men. *Arterioscler Thromb Vasc Biol*. 1997;17:2004-9.
6. Majmudar NG, Robson SC, Ford GA. Effects of the menopause, gender, and estrogen replacement therapy on vascular nitric oxide activity. *J Clin Endocrinol Metab*. 2000;85:1577-83.

<http://dx.doi.org/10.1016/j.jtcvs.2013.01.025>

Reply to the Editor:

We thank the authors for their comments and interest in our study showing improved long-term survival after coronary artery bypass grafting (CABG) using the radial artery (RA) in women. We agree with the many important advantages of RA grafting, which they so clearly outline. We also found their novel suggestion that age-dependent levels of androgens and estrogens may affect patency of RA conduits and, thus, survival in men and women to be intriguing.

We are, thus, analyzing our clinical data to better define the relationship of age and sex on RA patency and survival after CABG using the RA as the second arterial graft. We again appreciate their interest in our work and their excellent suggestion for further investigation.

Kamellia R. Dimitrova, MD

Robert F. Tranbaugh, MD

Division of Cardiac Surgery

Beth Israel Medical Center

New York, NY

<http://dx.doi.org/10.1016/j.jtcvs.2013.03.028>

Reply to the Editor:

We thank Tavlasioglu and colleagues¹ for their insightful comments in the February issue of the *Journal*. Our recently published article revealed that a small, hyperkinetic heart can be a risk factor for systolic anterior motion after mitral valve repair.² Although the underlying mechanism of this finding has not been determined, Tavlasioglu and colleagues¹ proposed that the mechanism may involve a morphologic aspect. We agree with this idea, because a small heart can be associated with closer proximity of mitral coaptation to the left ventricular outflow tract, which in an experimental study was demonstrated to contribute to the development of systolic anterior motion.³ In addition, they proposed that the small, hyperkinetic heart can

be grouped as “narrow aorta–mitral angle and associated factors.” Their proposal is based on the assumption that aorta–mitral angle must be narrow in a heart with a low systolic diameter. We think that this assumption needs to be proved, however, and we cannot argue in favor or against it because of the lack of data concerning aorta–mitral angle in our study.

We would like to add another possibility with respect to the mechanism of our finding. We speculate that the mechanism also involves a functional aspect. A hyperkinetic heart can be associated with high flow velocity in left ventricular out-flow tract, which has also been

demonstrated in an experimental study to contribute to the development of systolic anterior motion.⁴ We therefore believe that the mechanism may be multifactorial and should not be explained solely from a morphologic point of view.

Susumu Manabe, MD^a

Hitoshi Kasegawa, MD^b

^aDepartment of Cardiovascular
Surgery

Teikyo University Hospital
Tokyo, Japan

^bDepartment of Cardiovascular
Surgery

Sakakibara Heart Institute
Tokyo, Japan

References

1. Tavlasoglu M, Durukan AB, Gurbuz HA. Is a “narrow aorta-mitral angle and associated factors” associated with development of systolic anterior motion? *J Thorac Cardiovasc Surg*. 2013; 145:617.
2. Manabe S, Kasegawa H, Fukui T, Tabata M, Shinozaki T, Shimokawa T, et al. Influence of left ventricular function on development of systolic anterior motion after mitral valve repair. *J Thorac Cardiovasc Surg*. Epub 2012 Aug 10.
3. He S, Hopmeyer J, Lefebvre XP, Schwammenthal E, Yoganathan AP, Levine RA. Importance of leaflet elongation in causing systolic anterior motion of the mitral valve. *J Heart Valve Dis*. 1997;6: 149-59.
4. Lefebvre XP, He S, Levine RA, Yoganathan AP. Systolic anterior motion of the mitral valve in hypertrophic cardiomyopathy: an in vitro pulsatile flow study. *J Heart Valve Dis*. 1995;4:422-38.

<http://dx.doi.org/10.1016/j.jtcvs.2013.01.025>