strategies other than pharmacologic treatment, or the use of other drugs of the same class with predominantly analgesic effects not affecting brain receptors (eg, morphine derivatives) may induce the surgeon to refrain from administering a life-threatening drug cocktail that may be responsible for triggering or worsening delirium.

Francesco Pollari, MD
Giuseppe Santarpino, MD

Theodor Fischlein, MD
Department of Cardiac Surgery
Klinikum Nürnberg
Nuremberg, Germany

References

1. Arenson BG, MacDonald LA, Grocott HP, Hiebert BM, Arora RC. Effect of intensive care unit environment on in-hospital delirium after cardiac surgery. *J Thorac Cardiovasc Surg.* 2013;146:172-8.
2. Inouye SK. Delirium in older persons. *N Engl J Med.* 2006;354:1157-65. Erratum in: *N Engl J Med.* 2006;354:1655.
3. Santarpino G, Fasol R, Sirch J, Ackermann B, Pfeiffer S, Fischlein T. Impact of bispectral index monitoring on postoperative delirium in patients undergoing aortic surgery. *HSR Proc Intensive Care Cardiovasc Anesth.* 2011;3:47-58.
4. Martin BJ, Buth KJ, Arora RC, Baskett RJ. Delirium: a cause for concern beyond the immediate postoperative period. *Ann Thorac Surg.* 2012;93:1114-20.
5. Zeller A, Crestani F, Camenisch I, Iwasato T, Itoharu S, Fritschy JM, et al. Cortical glutamatergic neurons mediate the motor sedative action of diazepam. *Mol Pharmacol.* 2008;73:282-91.

<http://dx.doi.org/10.1016/j.jtcvs.2013.09.079>

Reply to the Editor:

We are appreciative and in agreement with the point of view that Pollari and colleagues have expressed regarding our study outlining some of the risk factors for postoperative delirium after cardiac surgery. One of the messages inherent in our observational study is that certain potentially modifiable factors can, if appropriately commented on the study by Santarpino and colleagues examining the effect of cerebral monitoring as a method of identifying patients who might have a greater incidence of delirium. Although it is exciting that the use of

a cerebral monitor might represent a clinically available biomarker, the data supporting the use of processed electroencephalographic monitoring (eg, the bispectral index and near-infrared spectroscopy cerebral oximetry) are quite sparse. Importantly, their report appears to have confused near-infrared spectroscopy and processed electroencephalographic monitoring. Thus, although we are appreciative of their interest in our report, their points regarding near-infrared spectroscopy would be much better supported by a study that had not used a processed electroencephalography (bispectral index) to estimate the cerebral saturation. Finally, we do agree with the authors that a focus on preventative measures is a necessary direction of future research in addressing delirium in the postoperative cardiac surgery patient.

Benjamin G. Arenson, BSc^a
Lindsey A. MacDonald, BSc^b
Hilary P. Grocott, MD, FRCPC^a
Brett M. Hiebert, MSc^c
Rakesh C. Arora, MD, PhD, FRCSC^d

^aDepartment of Anesthesia and
Perioperative Medicine
^bUniversity of Manitoba Faculty of
Medicine

Winnipeg, Manitoba, Canada
^cCardiac Sciences Program
Winnipeg Regional Health Authority
Winnipeg, Manitoba, Canada

^dDepartments of Anesthesia and
Perioperative Medicine and Surgery
University of Manitoba Faculty of
Medicine
Winnipeg, Manitoba, Canada

<http://dx.doi.org/10.1016/j.jtcvs.2013.11.059>

THE EFFECTS OF BICUSPID AORTA

To the Editor:

Congratulations to Katayama and colleagues¹ for this study showing that tricuspid valves are superior in

the aortic position. The diameter of the aorta increases 10% in systole relative to diastole. Tricuspid aortic leaflets do not resist this increase, but bicuspid aortic leaflets do resist because of their position. Bicuspid aortic leaflets thus undergo trauma during systole. In addition, tricuspid aortic leaflets get closer to the aortic wall during systole than do bicuspid ones. This causes bicuspid aortic leaflets to be exposed to more friction and shear stress. Twisting of the ascending aorta laterally and dorsally and localization of the coaptation line of bicuspid leaflets, however, were ignored in this simulation. The differences in the characteristics will influence the flow patterns. Configuration of the aortic valve and ascending aorta should therefore be considered in the simulation.

The noncoronary wall has the highest dp/dt in the ascending aorta. Therefore, in people with normal (tricuspid) aortic valves, the longest free edge is that of the noncoronary leaflet. Ninety-five percent of aortic dissections start from the wall of the noncoronary cusp. One of the most important factors affecting the surgical success of repair of bicuspid aorta is the localization of the commissure on the annulus. The localization of the commissures at the anteroposterior position on the annulus would increase surgical success. In this position, the noncoronary wall of the aorta would be less influenced by the dp/dt ratio. We think that the anatomic positions of the commissures on the annulus should be taken into consideration in this study.

Habib Cakir, MD
Mert Kestelli
Ismail Yurekli, MD
Koksai Donmez, MD
Department of Cardiovascular
Surgery

Katip Celebi University Izmir Atatürk
Training and Research Hospital
Izmir, Turkey