- (seventh) edition of the TNM classification of malignant tumours. *J Thorac Oncol*. 2007;2:706-14.
16. Rusch VW, Asamura H, Watanabe H, Giroux DJ, Rami-Porta R, Goldstraw P. Members of IASLC Staging Committee. The IASLC lung cancer staging project: a proposal for a new international lymph node map in the forthcoming seventh edition of the TNM classification for lung cancer. *J Thorac Oncol*. 2009;4:568-77.
 17. Oki M, Saka H, Kitagawa C, Kogure Y, Murata N, Ichihara S, et al. Prospective study of endobronchial ultrasound-guided transbronchial needle aspiration of lymph nodes versus transbronchial lung biopsy of lung tissue for diagnosis of sarcoidosis. *J Thorac Cardiovasc Surg*. 2012;143:1324-9.
 18. Herth FJ, Eberhardt R, Vilmann P, Krasnik M, Ernst A. Real-time endobronchial ultrasound guided transbronchial needle aspiration for sampling mediastinal lymph nodes. *Thorax*. 2006;61:795-8.
 19. Oki M, Saka H, Kitagawa C, Kogure Y, Murata N, Adachi T, et al. Transesophageal bronchoscopic ultrasound-guided fine needle aspiration for diagnosis of sarcoidosis. *Respiration*. 2013;85:137-43.
 20. Nakajima T, Yasufuku K, Iyoda A, Yoshida S, Suzuki M, Sekine Y, et al. The evaluation of lymph node metastasis by endobronchial ultrasound-guided transbronchial needle aspiration: crucial for selection of surgical candidates with metastatic lung tumors. *J Thorac Cardiovasc Surg*. 2007;134:1485-90.
 21. Zielinski M, Szlubowski A, Kołodziej M, Orzechowski S, Laczynska E, Pankowski J, et al. Comparison of endobronchial ultrasound and/or endoesophageal ultrasound with transcervical extended mediastinal lymphadenectomy for staging and restaging of non-small-cell lung cancer. *J Thorac Oncol*. 2013;8:630-6.
 22. Tournoy KG, Keller SM, Annema JT. Mediastinal staging of lung cancer: novel concepts. *Lancet Oncol*. 2012;13:e221-9.
 23. Annema JT, van Meerbeeck JP, Rintoul RC, Dooms C, Deschepper E, Dekkers OM, et al. Mediastinoscopy vs endosonography for mediastinal nodal staging of lung cancer: a randomized trial. *JAMA*. 2010;304:2245-52.
 24. Zhang R, Ying K, Shi L, Zhang L, Zhou L. Combined endobronchial and endoscopic ultrasound-guided fine needle aspiration for mediastinal lymph node staging of lung cancer: a meta-analysis. *Eur J Cancer*. 2013;49:1860-7.
 25. Medford AR, Agrawal S. Single bronchoscope combined endoscopic-endobronchial ultrasound-guided fine-needle aspiration for tuberculous mediastinal nodes. *Chest*. 2010;138:1274.
 26. Hwangbo B, Lee HS, Lee GK, Lim KY, Lee SH, Kim HY, et al. Transesophageal needle aspiration using a convex probe ultrasonic bronchoscope. *Respirology*. 2009;14:843-9.
 27. Oki M, Saka H, Kitagawa C, Sato S. Bronchoscopic transesophageal ultrasound-guided needle aspiration: an alternative to the conventional transesophageal ultrasound-guided needle aspiration technique. *J Thorac Cardiovasc Surg*. 2010;139:1659-61.
 28. Oki M, Saka H, Kitagawa C. Transesophageal bronchoscopic ultrasound-guided fine-needle aspiration for diagnosis of peripheral lung cancer. *Ann Thorac Surg*. 2011;91:1613-6.
 29. Sharples LD, Jackson C, Wheaton E, Griffith G, Annema JT, Dooms C, et al. Clinical effectiveness and cost-effectiveness of endobronchial and endoscopic ultrasound relative to surgical staging in potentially resectable lung cancer: results from the ASTER randomised controlled trial. *Health Technol Assess*. 2012;16:1-75, iii-iv.
 30. Lee HS, Lee GK, Lee HS, Kim MS, Lee JM, Kim HY, et al. Real-time endobronchial ultrasound-guided transbronchial needle aspiration in mediastinal staging of non-small cell lung cancer: how many aspirations per target lymph node station? *Chest*. 2008;134:368-74.
 31. Block MI. Endobronchial ultrasound for lung cancer staging: how many stations should be sampled? *Ann Thorac Surg*. 2010;89:1582-7.
 32. Uemura S, Yasuda I, Kato T, Doi S, Kawaguchi J, Yamauchi T, et al. Preoperative routine evaluation of bilateral adrenal glands by endoscopic ultrasound and fine-needle aspiration in patients with potentially resectable lung cancer. *Endoscopy*. 2013;45:195-201.
 33. Liberman M, Duranceau A, Grunenwald E, Thiffault V, Khereba M, Ferraro P. Initial experience with a new technique of endoscopic and ultrasonographic access for biopsy of para-aortic (station 6) mediastinal lymph nodes without traversing the aorta. *J Thorac Cardiovasc Surg*. 2012;144:787-92.
 34. Lee BE, Kletsman E, Rutledge JR, Korst RJ. Utility of endobronchial ultrasound-guided mediastinal lymph node biopsy in patients with non-small cell lung cancer. *J Thorac Cardiovasc Surg*. 2012;143:585-90.

EDITORIAL COMMENTARY

Pathologic staging of the mediastinum: When and how?

Jacob A. Klapper, MD, and Chadrick E. Denlinger, MD

Pathologic staging of mediastinal lymph nodes before surgical resection is the accepted standard for patients with non-small cell lung cancer (NSCLC).¹ Mediastinoscopy, endobronchial ultrasound (EBUS), and endoscopic ultrasound are all acceptable means of obtaining tissue. Two prospective studies compared the sensitivity of EBUS with mediastinoscopy among patients with NSCLC

and concluded that the 2 modalities are equivalent.^{2,3} In the first study patients were randomized to either mediastinoscopy or EBUS followed by mediastinoscopy if EBUS was negative. The sensitivity of mediastinoscopy alone was 79% compared with 85% for EBUS. The addition of mediastinoscopy in patients previously evaluated by EBUS increased the sensitivity to 94%.² In the second study, by Yasufuku and colleagues,³ each patient was evaluated by both EBUS and mediastinoscopy and the sensitivities of the 2 were 81% and 79%, respectively.

The sensitivity of 73% for combined EBUS and endoscopic ultrasound presented by Oki and colleagues⁴ in this issue is congruent with prior surgical series where patients went on to resection if the mediastinum was

From the Division of Cardiothoracic Surgery, Medical University of South Carolina, Charleston, SC.

Disclosures: Authors have nothing to disclose with regard to commercial support. Address for reprints: Chadrick E. Denlinger, MD, 25 Courtenay Dr, ART Ste 7018, Charleston, SC 29425. (E-mail: denlinge@musc.edu).

J Thorac Cardiovasc Surg 2014;148:1177-8

0022-5223/\$36.00

Copyright © 2014 by The American Association for Thoracic Surgery

<http://dx.doi.org/10.1016/j.jtcvs.2014.09.001>

pathologically negative, but it was substantially less than a collective sensitivity rate of 89% presented in a recent review by Silvestri and colleagues.⁵ Although the reasons for the difference in sensitivity are unclear, 1 likely explanation is a difference in pretest probability. The mean lymph node size of 8.2 mm in the study by Oki and colleagues⁴ was similar to the reported size of 6.9 mm by Yasufuku and colleagues.³ Although positron emission tomography scan findings were not included in these results from Yasufuku and colleagues,⁴ the size indicates that these patients likely had no radiographic evidence of mediastinal involvement. In fact, patients with bulky mediastinal lymph nodes were excluded from the current study by Oki and colleagues.⁴ The greater sensitivity in previously published series may reflect the inclusion of patients with radiographically suspicious lymph nodes that were essentially confirmed with EBUS. This concept is further supported by the understanding that patients with a positive mediastinoscopy that were missed by EBUS frequently have micrometastatic disease.³

The disease burden as well as treatment plans for patients with stage IIIA NSCLC are each quite variable. Many institutions treat patients with involved mediastinal lymph nodes with either neoadjuvant chemotherapy with or without radiation followed by restaging and resection or with definitive chemotherapy and radiation. The inclusion of surgery for those patients who can be treated with a lobectomy was supported by the Intergroup 0139 prospective randomized study.⁶ Although the overall study did not indicate that surgical resection conveyed any survival advantage, these results were influenced by extremely high mortality rate among patients undergoing pneumonectomy. In an unplanned post hoc analysis that included approximately 60% of the original cohort, patients treated with a lobectomy demonstrated a significant survival advantage compared with patients treated with chemotherapy and radiation followed by more radiation.

A recent study of patients⁷ who underwent surgical resection and were incidentally found to have mediastinal lymph node involvement demonstrated 5-year survival rates

of 66.6% and 36.4% for those with single-station and multistation disease, respectively. These survival rates are substantially greater than expected for patients with stage IIIA disease.⁸ It remains unclear if this difference in survival relates primarily to a selection bias—where patients with minimal N2 disease and an excellent performance status were treated surgically and less suitable patients were treated with chemotherapy and radiation—or if there is an oncologic benefit of primary surgical resection in this patient population. Further investigations may address if preoperative pathologic staging of the mediastinum provides any benefit for patients who are radiographically negative or if these patients would be best treated with surgical resection and adjuvant chemotherapy for incidentally identified lymph node involvement.

References

1. National Comprehensive Cancer Network. Available at: http://www.nccn.org/professionals/physician_gls/pdf/nscl.pdf. Accessed August 29, 2014.
2. Annema JT, van Meerbeeck JP, Rintoul RC, Dooms C, Deschepper E, Dekkers OM, et al. Mediastinoscopy vs endosonography for mediastinal nodal staging of lung cancer: a randomized trial. *JAMA*. 2010;304:2245-52.
3. Yasufuku K, Pierre A, Darling G, de Perrot M, Waddell T, Johnston M, et al. A prospective controlled trial of endobronchial ultrasound-guided transbronchial needle aspiration compared with mediastinoscopy for mediastinal lymph node staging of lung cancer. *J Thorac Cardiovasc Surg*. 2011;142:1393-400.
4. Oki M, Saka H, Ando M, Kitagawa C, Kogure Y, Seki Y. Endoscopic ultrasound-guided fine needle aspiration and endobronchial ultrasound-guided transbronchial needle aspiration: are two better than one in mediastinal staging of non-small cell lung cancer? *J Thorac Cardiovasc Surg*. 2014;148:1169-77.
5. Silvestri GA, Gonzalez AV, Jantz MA, Margolis ML, Gould MK, Tanoue LT, et al. Methods for staging non-small cell lung cancer: diagnosis and management of lung cancer, 3rd ed: American College of Chest Physicians evidence-based clinical practice guidelines. *Chest*. 2013;143(5 Suppl):e211S-50S.
6. Albain KS, Swann RS, Rusch VW, Turrisi AT, Shepherd FA, Smith C, et al. Radiotherapy plus chemotherapy with or without surgical resection for stage III non-small cell lung cancer: a phase III randomized controlled trial. *Lancet*. 2009;374:379-86.
7. Cho HJ, Kim SR, Kim HR, Han JO, Kim YH, Kim DK, et al. Modern outcome and risk analysis of surgically resected occult N2 non-small cell lung cancer. *Ann Thorac Surg*. 2014;97:1920-5.
8. Goldstraw P, Crowley J, Chansky K, Giroux DJ, Groome PA, Rami-Porta R, et al. The IASCLC lung cancer staging project: Proposals for the revision of the TNM stage groupings in the forthcoming (seventh) edition of the TNM classification of malignant tumors. *J Thorac Oncol*. 2007;2:706-14.