13. Folliquet TA, Laborde F, Zannis K, Ghorayeb G, Haverich A, Shrestha M. Sutureless Perceval aortic valve replacement: results of two European centers. *Ann Thorac Surg*. 2012;93:1483-8.
14. Santarpino G, Pfeiffer S, Schmidt J, Concistrè G, Fischlein T. Sutureless aortic valve replacement: first-year single center experience. *Ann Thorac Surg*. 2012;94:504-9.
15. Shrestha M, Timm R, Höffeler K, Koigeldiyev N, Khaladj N, Hagl C, et al. Minimally invasive aortic valve replacement with self-anchoring Perceval valve. *J Heart Valve Dis*. 2013;22:230-5.
16. Glauber M, Miceli A, Bevilacqua S, Farneti PA. Minimally invasive aortic valve replacement via right anterior minithoracotomy: early outcomes and midterm follow-up. *J Thorac Cardiovasc Surg*. 2011;142:1577-9.
17. Santarpino G, Pfeiffer S, Concistrè G, Fischlein T. Perceval S aortic valve implantation in mini-invasive surgery: the simple sutureless solution. *Interact Cardiovasc Thorac Surg*. 2012;15:357-60.
18. Karimov JH, Santarelli F, Murzi M, Glauber M. A technique of an upper V-type ministernotomy in the second intercostal space. *Interact Cardiovasc Thorac Surg*. 2009;9:1021-2.
19. Miceli A, Murzi M, Gilmanov D, Fugà R, Ferrarini M, Solinas M, et al. Minimally invasive aortic valve replacement using right minithoracotomy is associated with better outcomes than ministernotomy. *J Thorac Cardiovasc Surg*. Epub 2013 Sep 12.
20. Gilmanov D, Farneti PA, Miceli A, Bevilacqua S, Glauber M. Perceval S sutureless aortic valve prosthesis implantation via a right anterior minithoracotomy. *Multimedia Manual Cardiothorac Surg*. Epub 2013 Jul 12; doi:10.1093/mmts/mmt012.
21. Santarpino G, Pfeiffer S, Concistrè G, Grossmann I, Hinzmann M, Fischlein T. The Perceval S aortic valve has the potential of shortening surgical time: does it also result in improved outcomes? *Ann Thorac Surg*. 2013;96:77-82.
22. Brennan JM, Edwards FH, Zhao Y, O'Brien SM, Douglas PS, Peterson ED. Developing Evidence to Inform Decision About Effectiveness—Aortic Valve Replacement (DECADE AVR) Research Team. Long-term survival after aortic valve replacement among high-risk elderly patients in the United States. Insight from The Society of Thoracic Surgeons adult cardiac surgery database, 1991 to 2007. *Circulation*. 2012;126:1621-9.
23. Miller DC, Blackstone EH, Mack MJ, Svensson LG, Kodali SK, Kapadia S, et al. Transcatheter (TAVR) versus surgical (AVR) aortic valve replacement: occurrence, hazard, risk factors, and consequences of neurological events in the PARTNER trial. *J Thorac Cardiovasc Surg*. 2012;143:832-43.
24. Neyt M, Van Brabant H, Devriese S, Van De Sande S. A cost-utility analysis of transcatheter aortic valve implantation in Belgium: focusing on a well-defined and identifiable population. *BMJ Open*. 2012;2:e001032.
25. D'Onofrio A, Mesina A, Lorusso R, Alfieri OR, Fusari M, Rubino P, et al. Sutureless aortic valve replacement as an alternative treatment for patients belonging to the "gray zone" between transcatheter aortic valve implantation and conventional surgery: a propensity-matched, multicenter analysis. *J Thorac Cardiovasc Surg*. 2012;144:1010-8.
26. Pradelli L, Zaniolo O. Perceval Sutureless valves in isolated and concomitant AVR procedures: an economic model shows overall decrease of costs for isolated or combined operations. *Farmeconomia. Health Economics and Therapeutic Pathways*. 2012;13:159-74.
27. Zoghbi WA, Chambers JB, Dusmenil JG, Foster E, Gottdiener JS, Grayburn PA, et al. Recommendations for evaluation of prosthetic valves with echocardiography and Doppler ultrasound: a report From the American Society of Echocardiography's Guidelines and Standards Committee and the Task Force on Prosthetic Valves, developed in conjunction with the American College of Cardiology Cardiovascular Imaging Committee, Cardiac Imaging Committee of the American Heart Association, the European Association of Echocardiography, a registered branch of the European Society of Cardiology, the Japanese Society of Echocardiography and the Canadian Society of Echocardiography, endorsed by the American College of Cardiology Foundation, American Heart Association, European Association of Echocardiography, a registered branch of the European Society of Cardiology, the Japanese Society of Echocardiography, and Canadian Society of Echocardiography. *J Am Soc Echocardiogr*. 2009;22:975-1014.

EDITORIAL COMMENTARY

Minimally invasive Perceval S implantation: Does this fill a sizable niche between sutured and transcatheter aortic valve replacement?

Michael O. Kayatta, MD, Hanna A. Jensen, MD, PhD, and Vinod H. Thourani, MD

See related article on pages 2838-43.

From the Structural Heart and Valve Center, Division of Cardiothoracic Surgery, Joseph B. Whitehead Department of Surgery, Emory University School of Medicine, Atlanta, Ga.

Disclosures: V.H.T. serves on the advisory boards of St Jude Medical, Edwards Lifesciences, Boston Scientific, and Abbot Medical and engages in research for Medtronic, Edwards Lifesciences, St Jude, and Sorin Medical. All other authors have nothing to disclose with regard to commercial support.

Received for publication Oct 2, 2014; accepted for publication Oct 2, 2014

Address for reprints: Vinod H. Thourani, MD, Emory University Hospital Midtown, 550 Peachtree St NE, 6th Floor, Medical Office Tower, Atlanta, GA 30308 (E-mail: vthoura@emory.edu).

J Thorac Cardiovasc Surg 2014;148:2843-4

0022-5223/\$36.00

Copyright © 2014 by The American Association for Thoracic Surgery

<http://dx.doi.org/10.1016/j.jtcvs.2014.10.004>

In this issue of the *Journal*, Miceli and colleagues¹ present a well-written article on their early experience with placement of the sutureless Perceval S (Sorin SpA, Milan, Italy) valve through minimally invasive surgical approaches of either partial sternotomy or right anterior thoracotomy. Of all patients coming to their centers for isolated aortic valve replacement (AVR), 40% (n = 281) were found eligible for these techniques. These patients were elderly (mean age, 76 years) but on average were a relatively healthy intermediate-risk population. In-hospital outcomes were excellent, with very low mortality and only 1.8% of patients having greater than trace paravalvular leak. This rate of leaks is extremely low, and has not been seen by other groups.^{2,3} Overall, the new postoperative pacemaker rate was only 4.2%. These results are to be commended and it is to be hoped will be reproducible

with wider adoption of this class of valves and increasing surgeon experience.

During the past decade, there has been an explosion of disruptive technologies in the management of patients requiring AVR. Broadly, management can be divided into surgical AVR requiring cardiopulmonary bypass and transcatheter AVR (TAVR) techniques avoiding the bypass circuitry. Traditionally, surgeons have used suture-based valves, the majority of which are stented, with stentless aortic valves less commonly used. The current series of Miceli and colleagues¹ reports on the newest sutureless valve technology for AVR.

Currently, minimally invasive AVR is being performed in a minority of heart centers. Compared with AVR through a full sternotomy, minimally invasive AVR generally has a slightly longer cardiopulmonary bypass time, which the use of the sutureless valve system could minimize by requiring significantly less suturing. It is hoped that sutureless valve therapies will lead to an increased adoption of minimally invasive techniques that may offer some benefits not only in cosmesis and patient satisfaction but also in improved respiratory mechanics.

The potential advantages of sutureless valves when compared with TAVR are the decreased paravalvular leak rates and decreased need for postoperative pacemakers. Miceli and colleagues¹ have noted that the sutureless valve technology is a viable alternative to high-risk TAVR; however, the median euroSCORE of their patients was only 8 (interquartile range 5-12). For comparison, in the landmark randomized, controlled Placement of AoRtic TraNscathetER Valves (PARTNER) trial⁴ the mean euroSCORE for the high-risk TAVR population was 29.3 ± 16.3 . Further studies are required in patient popula-

tions with a comparable risk profile before conclusions can be drawn about the optimal valve for the high-risk patients with aortic stenosis. It should also be noted that newer generations of TAVR are under development that may narrow the gap of paravalvular leak between TAVR and surgical AVR.⁵

Economic considerations may determine the widespread adoption of this highly innovative technology. Although the current prices are approximately half those of TAVR valves, they are approximately 2 to 3 times those of stented valves. A robust cost analysis will help determine the overall penetrance of sutureless valves.

For the surgical community, the results of this study represent yet another important advancement that expands our armamentarium to treat aortic valve disease. The full implications and the most appropriate target population must be determined in larger trials.

References

1. Miceli A, Santarpino G, Pfeiffer S, Murzi M, Gilmanov D, Concistré G, et al. Minimally invasive aortic valve replacement with Perceval S sutureless valve: early outcomes and one-year survival from two European centers. *J Thorac Cardiovasc Surg*. 2014;148:2838-43.
2. Folliguet TA, Laborde F, Zannis K, Ghorayeb G, Haverich A, Shrestha M. Sutureless Perceval aortic valve replacement: results of two European centers. *Ann Thorac Surg*. 2012;93:1483-8.
3. D'Onofrio A, Rizzoli G, Messina A, Alfieri O, Lorusso R, Salizzoni S, et al. Conventional surgery, sutureless valves, and transapical aortic valve replacement: what is the best option for patients with aortic valve stenosis? A multicenter, propensity-matched analysis. *J Thorac Cardiovasc Surg*. 2013;146:1065-70; discussion 1070-1.
4. Smith CR, Leon MB, Mack MJ, Miller DC, Moses JW, Svensson LG, et al., PARTNER Trial Investigators. Transcatheter versus surgical aortic-valve replacement in high-risk patients. *N Engl J Med*. 2011;364:2187-98.
5. Généreux P, Head SJ, Hahn R, Daneault B, Kodali S, Williams MR, et al. Paravalvular leak after transcatheter aortic valve replacement: the new Achilles' heel? A comprehensive review of the literature. *J Am Coll Cardiol*. 2013;61:1125-36.