CONCLUSIONS

Creation of a central shunt using an ePTFE tube graft as an initial palliative procedure for patients with pulmonary atresia with VSD and MAPCAs is highly effective in rehabilitating the diminutive PAs, and a multistage approach focusing on rehabilitation of the diminutive PAs has produced encouraging results and good PA growth.

References

1. Amark KM, Karamlou T, O'Carroll A, MacDonald C, Freedom RM, Yoo SJ, et al. Independent factors associated with mortality, reintervention, and achievement of complete repair in children with pulmonary atresia with ventricular septal defect. *J Am Coll Cardiol*. 2006;47:1448-56.
2. Bull K, Somerville J, Ty E, Spiegelhalter D. Presentation and attrition in complex pulmonary atresia. *J Am Coll Cardiol*. 1995;25:491-9.
3. Reddy VM, Liddicoat JR, Hanley FL. Midline one-stage complete unifocalization and repair of pulmonary atresia with ventricular septal defect and major aortopulmonary collaterals. *J Thorac Cardiovasc Surg*. 1995;109:832-44; discussion 44-5.
4. Reddy VM, McElhinney DB, Amin Z, Moore P, Parry AJ, Teitel DF, et al. Early and intermediate outcomes after repair of pulmonary atresia with ventricular septal defect and major aortopulmonary collateral arteries: experience with 85 patients. *Circulation*. 2000;101:1826-32.
5. Duncan BW, Mee RB, Prieto LR, Rosenthal GL, Mesia CI, Qureshi A, et al. Staged repair of tetralogy of Fallot with pulmonary atresia and major aortopulmonary collateral arteries. *J Thorac Cardiovasc Surg*. 2003;126:694-702.
6. Liava'a M, Brizard CP, Konstantinov IE, Robertson T, Cheung MM, Weintraub R, et al. Pulmonary atresia, ventricular septal defect, and major aortopulmonary collaterals: neonatal pulmonary artery rehabilitation without unifocalization. *Ann Thorac Surg*. 2012;93:185-91.
7. Watterson KG, Wilkinson JL, Karl TR, Mee RB. Very small pulmonary arteries: central end-to-side shunt. *Ann Thorac Surg*. 1991;52:1132-7.
8. Barozzi L, Brizard CP, Galati JC, Konstantinov IE, Bohuta L, d'Udekem Y. Side-to-side aorto-GoreTex central shunt warrants central shunt patency and pulmonary arteries growth. *Ann Thorac Surg*. 2012;92:1476-82.
9. Gates RN, Laks H, Johnson K. Side-to-side aorto-Gore-Tex central shunt. *Ann Thorac Surg*. 1998;65:515-6.
10. Grosse-Wortmann L, Yoo SJ, van Arsdell G, Chetan D, MacDonald C, Benson L, et al. Preoperative total pulmonary blood flow predicts right ventricular pressure in patients early after complete repair of tetralogy of Fallot and pulmonary atresia with major aortopulmonary collateral arteries. *J Thorac Cardiovasc Surg*. 2013;146:1185-90.
11. Honjo O, Al-Radi OO, MacDonald C, Tran KC, Sapra P, Davey LD, et al. The functional intraoperative pulmonary blood flow study is a more sensitive predictor than preoperative anatomy for right ventricular pressure and physiologic tolerance of ventricular septal defect closure after complete unifocalization in patients with pulmonary atresia, ventricular septal defect, and major aortopulmonary collaterals. *Circulation*. 2009;120:S46-52.
12. Song SW, Park HK, Park YH, Cho BK. Pulmonary atresia with ventricular septal defects and major aortopulmonary collateral arteries. *Circ J*. 2009;73:516-22.
13. Metras D, Chetaille P, Kreitmann B, Fraisse A, Ghez O, Riberi A. Pulmonary atresia with ventricular septal defect, extremely hypoplastic pulmonary arteries, major aorto-pulmonary collaterals. *Eur J Cardiothorac Surg*. 2001;20:590-6; discussion 6-7.
14. Zhang Y, Hua Z, Yang K, Zhang H, Yan J, Wang X, et al. Outcomes of the rehabilitative procedure for patients with pulmonary atresia, ventricular septal defect and hypoplastic pulmonary arteries beyond the infant period. *Eur J Cardiothorac Surg*. 2014;46:297-303; discussion 303.
15. Brizard CP, Liava'a M, d'Udekem Y. Pulmonary atresia, VSD and Mapcas: repair without unifocalization. *Semin Thorac Cardiovasc Surg Pediatr Card Surg Annu*. 2009;139-44.
16. d'Udekem Y, Alphonso N, Norgaard MA, Cochrane AD, Grigg LE, Wilkinson JL, et al. Pulmonary atresia with ventricular septal defects and major aortopulmonary collateral arteries: unifocalization brings no long-term benefits. *J Thorac Cardiovasc Surg*. 2005;130:1496-502.
17. Fouilloux V, Bonello B, Kammache I, Fraisse A, Mace L, Kreitmann B. Management of patients with pulmonary atresia, ventricular septal defect, hypoplastic pulmonary arteries and major aorto-pulmonary collaterals: Focus on the strategy of rehabilitation of the native pulmonary arteries. *Arch Cardiovasc Dis*. 2012;105:666-75.

EDITORIAL COMMENTARY

Shunt first: One way to do it

Robert D. B. Jaquiss, MD

See related article on pages 515-20.

The management of children with ventricular septal defect, pulmonary valve atresia, diminutive central pulmonary arteries, and multiple aorto-pulmonary collateral

arteries is complex, because of the wide variability in pulmonary blood flow. Nowhere is the cliché that “every patient is unique” more apt than with this condition. In addition to the intrinsic anatomic variability that bedevils surgeons caring for such patients, recommendations are widely divergent for optimal provision of pulmonary blood supply. As to whether amalgamation of multiple aorto-pulmonary collateral arteries with native pulmonary arteries (unifocalization) should be performed if native pulmonary arteries are present, a reader of the *Journal* will have learned, in 2005, that unifocalization “brings no long-term benefits,”¹ only to discover a few years later that “unifocalization . . . is essential to achieve excellent outcomes.”²

A possible advantage of employing unifocalization is the potential to achieve a single-stage complete repair.^{3,4}

From the Division of Cardiothoracic Surgery, Duke University School of Medicine, Durham, NC.

Disclosures: Author has nothing to disclose with regard to commercial support.

Received for publication Nov 9, 2014; accepted for publication Nov 10, 2014; available ahead of print Dec 5, 2014.

Address for reprints: Robert D. B. Jaquiss, MD, Division of Cardiothoracic Surgery, Duke University School of Medicine, DUMC 3474, Durham, NC 27710 (E-mail: robert.jaquiss@duke.edu).

J Thorac Cardiovasc Surg 2015;149:520-1
0022-5223/\$36.00

Copyright © 2015 by The American Association for Thoracic Surgery
<http://dx.doi.org/10.1016/j.jtcvs.2014.11.026>

By contrast, a multistage approach is obviously necessary if preliminary “rehabilitation” of the native pulmonary arteries is favored, either by direct association of the right ventricle to the main pulmonary artery,⁵ or by creation of a central shunt. The central shunt may be accomplished by direct anastomosis,⁶ although this technique has typically been accomplished more recently by a polytetrafluoroethylene interposition graft.¹

In the present report by Kim and colleagues,⁷ the authors have forsworn the potential efficiency of a single-stage approach for a strategy of maximizing the growth of the native pulmonary arteries by use of an initial central shunt. The shunts from aorta to the native main pulmonary artery were constructed in 1 of 2 forms, depending on the size of the main pulmonary artery. The resultant configurations are demonstrated by the superb figures in the article. The outcomes in this report are indeed excellent, with the vast majority of patients (13 of 15) achieving complete repair. No operative deaths occurred at any stage, although 2 patients died in late follow-up.

Although these are outstanding results, this approach is very resource intensive. The mean hospital stay after the central shunt procedure was nearly 6 weeks, which seems long for a group whose surgery did not require cardiopulmonary bypass. Furthermore, a large number of operations, 3.8 per patient, were necessary to achieve complete repair, which was not accomplished until 24 months after the initial shunt. Stated another way, these data (3.8 operations in 24 months) suggest that the typical patient faces cardiovascular surgery every 6 months, presumably with multiple, interspersed, cardiac catheterizations and computed tomography angiograms. The aggregate financial costs of this strategy, not to mention the aggregate dosages of anesthetic medication and ionizing radiation, each with potential significant attendant risk, are obviously substantial.

Although unifocalization was not initially planned for any of the patients in the present series, it was ultimately necessary for a number of them, as demonstrated in the flow diagram outlining the details of operative staging. In addition, as is obvious from that diagram, 4 patients required a second systemic-to-pulmonary shunt before complete repair. Thus, the touted advantage of the “symmetrical distribution of pulmonary blood flow through the MPA [main pulmonary artery] using a central shunt” was achieved in only 73% of patients. This result is remarkably similar to the experience reported by Liava’a and colleagues,⁸ wherein a second shunt was necessary in 4 of 20 patients whose initial procedure was a systemic-to-pulmonary shunt.

An alternative way to enhance native pulmonary artery growth is to initially associate the main pulmonary artery to the right ventricle, as advocated by Metras and colleagues.⁵ At the price of an initial operation requiring cardiopulmonary bypass, this approach permits antegrade catheter intervention, which is not possible using the central shunt approach in the present report. Kim and colleagues⁷ concede this but believe they can simply defer catheter interventions until after placement of a right-ventricle-to-pulmonary-artery conduit.

In assessing the merit of the approach advocated in the present report, it is important to define success carefully. If the metric is operative survival, this approach is indeed successful. If the yardstick is the rate of ultimately achieving complete repair with a closed ventricular septal defect, the approach is successful, albeit expensive in procedure number and resource utilization. However, if success is defined as achievement of optimal functional status accompanied by low right ventricular pressure, the jury is still out, as the authors cannot provide us with right ventricular pressure data, as they admirably concede in their discussion of limitations. We, and their patients, eagerly await further reports of ongoing follow-up to answer this crucial question. What can be said is that the approach advocated in this report is safe and effective; whether it is the best approach remains to be seen.

References

1. d’Udekem Y, Alphonso N, Nørgaard MA, Cochrane AD, Grigg LE, Wilkinson JL, et al. Pulmonary atresia with ventricular septal defects and major aortopulmonary collateral arteries: unifocalization brings no long-term benefits. *J Thorac Cardiovasc Surg.* 2005;130:1496-502.
2. Davies B, Mussa S, Davies P, Stickley J, Jones TJ, Barron DJ, et al. Unifocalization of major aortopulmonary collateral arteries in pulmonary atresia with ventricular septal defect is essential to achieve excellent outcomes irrespective of native pulmonary artery morphology. *J Thorac Cardiovasc Surg.* 2009;138:1269-75.
3. Reddy VM, Liddicoat JR, Hanley FL. Midline one-stage complete unifocalization and repair of pulmonary atresia with ventricular septal defect and major aortopulmonary collaterals. *J Thorac Cardiovasc Surg.* 1995;109:832-44; discussion 844-5.
4. Lofland GK. The management of pulmonary atresia, ventricular septal defect, and multiple aorta pulmonary collateral arteries by definitive single stage repair in early infancy. *Eur J Cardiothorac Surg.* 2000;18:480-6.
5. Metras D, Chetaille P, Kreitmann B, Fraisse A, Ghez O, Riberi A. Pulmonary atresia with ventricular septal defect, extremely hypoplastic pulmonary arteries, major aorto-pulmonary collaterals. *Eur J Cardiothorac Surg.* 2001;20:590-6; discussion 596-7.
6. Watterson KG, Wilkinson JL, Karl TR, Mee RB. Very small pulmonary arteries: central end-to-side shunt. *Ann Thorac Surg.* 1991;52:1132-7.
7. Kim H, Sung SC, Choi KH, Lee HD, Ban GH, Chang YH. The role of a central shunt using an expanded polytetrafluoroethylene tube graft for rehabilitation of diminutive pulmonary arteries in patients with pulmonary atresia with ventricular septal defect and major aortopulmonary collaterals. *J Thorac Cardiovasc Surg.* 2015;149:515-20.
8. Liava’a M, Brizard CP, Konstantinov IE, Robertson T, Cheung MM, Weintraub R, et al. Pulmonary atresia, ventricular septal defect, and major aortopulmonary collaterals: neonatal pulmonary artery rehabilitation without unifocalization. *Ann Thorac Surg.* 2012;93:185-91.