- coronary intervention: a propensity score analysis. *Eur J Cardiothorac Surg.* 2012;41:94-101.
7. Jeong DS, Lee YT, Chung SR, Jeong JH, Kim WS, Sung K, et al. Revascularization in left main coronary artery disease: comparison of off-pump coronary artery bypass grafting vs percutaneous coronary intervention. *Eur J Cardiothorac Surg.* 2013;44:718-24.
 8. Yi G, Joo HC, Youn YN, Hong S, Yoo KJ. Stent versus off-pump coronary bypass grafting in the second-generation drug-eluting stent era. *Ann Thorac Surg.* 2013; 96:535-41.
 9. Weiss AJ, Zhao S, Tian DH, Taggart DP, Yan TD. A meta-analysis comparing bilateral internal mammary artery with left internal mammary artery for coronary artery bypass grafting. *Ann Cardiothorac Surg.* 2013;2: 390-400.
 10. Grau JB, Ferrari G, Mak AW, Shaw RE, Brizzio ME, Mindich BP, et al. Propensity matched analysis of bilateral internal mammary artery versus single left internal mammary artery grafting at 17-year follow-up: validation of a contemporary surgical experience. *Eur J Cardiothorac Surg.* 2012;41:770-5.
 11. Schwann TA, Tranbaugh RF, Dimitrova KR, Engoren MC, Kabour A, Hoffman DM, et al. Time-varying survival benefit of radial artery versus vein grafting: a multiinstitutional analysis. *Ann Thorac Surg.* 2014;97:1328-34.
 12. Shi WY, Hayward PA, Fuller JA, Tatoulis J, Rosalion A, Newcomb AE, et al. Is the radial artery associated with improved survival in older patients undergoing coronary artery bypass grafting? An analysis of a multicentre experience. *Eur J Cardiothorac Surg.* February 15, 2015 [Epub ahead of print].
 13. Tranbaugh RF, Dimitrova KR, Friedmann P, Geller CM, Harris LJ, Stelzer P, et al. Radial artery conduits improve long-term survival after coronary artery bypass grafting. *Ann Thorac Surg.* 2010;90:1165-72.
 14. Kappetein AP, Dawkins KD, Mohr FW, Morice MC, Mack MJ, Russell ME, et al. Current percutaneous coronary intervention and coronary artery bypass grafting practices for three-vessel and left main coronary artery disease. Insights from the SYNTAX run-in phase. *Eur J Cardiothorac Surg.* 2006;29:486-91.
 15. Kobayashi J. Current status of coronary artery bypass grafting. *Gen Thorac Cardiovasc Surg.* 2008;56:260-7.
 16. Zamvar V, Williams D, Hall J, Payne N, Cann C, Young K, et al. Assessment of neurocognitive impairment after off-pump and on-pump techniques for coronary artery bypass graft surgery: prospective randomised controlled trial. *BMJ.* 2002; 325:1268.
 17. Diegeler A, Hirsch R, Schneider F, Schilling LO, Falk V, Rauch T, et al. Neuro-monitoring and neurocognitive outcome in off-pump versus conventional coronary bypass operation. *Ann Thorac Surg.* 2000;69:1162-6.
 18. Keeling WB, Williams ML, Slaughter MS, Zhao Y, Puskas JD. Off-pump and on-pump coronary revascularization in patients with low ejection fraction: a report from the society of thoracic surgeons national database. *Ann Thorac Surg.* 2013; 96:83-8.
 19. Moss E, Puskas JD, Thourani VH, Kilgo P, Chen EP, Leshnower BG, et al. Avoiding aortic clamping during coronary artery bypass grafting reduces postoperative stroke. *J Thorac Cardiovasc Surg.* 2015;149:175-80.
 20. Calafiore AM, Di Mauro M, Teodori G, Di Giammarco G, Cirmeni S, Contini M, et al. Impact of aortic manipulation on incidence of cerebrovascular accidents after surgical myocardial revascularization. *Ann Thorac Surg.* 2002; 73:1387-93.
 21. Kim KB, Kang CH, Chang WI, Lim C, Kim JH, Ham BM, et al. Off-pump coronary artery bypass with complete avoidance of aortic manipulation. *Ann Thorac Surg.* 2002;74:S1377-82.

<http://dx.doi.org/10.1016/j.jtcvs.2015.02.030>

OPCAB IS NOT FOR EVERY SURGEON AND NOT FOR EVERY PATIENT

Reply to the Editor:

Drs Yanagawa and Puskas have written a thorough, thoughtful, and articulate piece of effective advocacy in defense of off-pump coronary artery bypass (OPCAB), fortified by selective citation. I will not attempt to reply point by point.

I suspect we all agree that there are no perfect clinical trials. When meta-analyzers put the conclusions and imperfections of multiple trials in the salad spinner, out comes...salad. When differences between the modalities being compared are dramatic, none of that matters! The problem for OPCAB, now deep into its second decade of practice, is that convincingly demonstrated benefits have been illusory, and certain deficits (graft patency) have been stubbornly persistent.

When a procedure is called into question, a popular defense strategy used by surgeons is "not in my hands." I've used it myself! In the world of OPCAB, few surgeons bring as much legitimacy to that defense as Dr Puskas, whose randomized trials have been uniquely positive for OPCAB. But even Dr Puskas cannot be expected to perform every coronary artery bypass graft, and the super-expert is not a good solution to a public health problem the size of coronary disease.

A large part of me is disappointed by these developments. I have spent more than 15 years acquiring what I consider to be expertise in OPCAB. Certain technical benefits still have strong appeal. However, I agree completely with the authors' "not for every surgeon, not for every patient" precept, and I am dramatically expanding my application of the second "not."

Craig R. Smith, MD
Department of Surgery
Columbia University Medical Center of New York
Presbyterian Hospital
New York, NY

<http://dx.doi.org/10.1016/j.jtcvs.2015.03.005>

HARD ENDPOINTS FROM LARGE RANDOMIZED, CLINICAL TRIALS ARE MORE IMPORTANT THAN EARLY REPORTS OF SURROGATE ENDPOINTS FROM SMALL STUDIES

Reply to the Editor:

Thank you for the opportunity to comment on the letter by Yanagawa and Puskas, especially regarding our recent paper on graft patency in the Danish On-Pump versus Off-Pump Randomization Study (DOORS).¹

The conversion rate in DOORS was 12.8%,² similar to that in the American Randomized On-pump Off-pump Bypass (ROOBY) trial.³ Outside clinical trials, off-pump coronary artery bypass grafting (OPCAB) is often preferred to on-pump coronary artery bypass grafting (ONCAB) for patients who need relatively few grafts, tending to keep conversion rates low. Yanagawa and Puskas refer to a number of 2.2% as being the proportion of conversions in the Society of Thoracic Surgeons database, but the only reference

made in the letter concerning the Society of Thoracic Surgeons database is the one regarding Keeling and co-workers,⁴ who reported a conversion rate of 5.2%.

Yanagawa and Puskas write that “inexplicably, more perioperative bleeding was found in OPCAB patients than in ONCAB patients in DOORS,” suggesting that this was an indication of incomplete comfort level with OPCAB among some of the operating surgeons. This finding is different from those from earlier studies and—as described in the article—more likely caused by a higher degree of heparinization of the OPCAB patients in this study compared to earlier studies.⁵ This difference was not large enough to cause differences in the proportion of patients receiving perioperative blood transfusions²; however, we do not recommend that this higher degree of heparinization be implemented clinically. It is interesting that in DOORS the overall 30-day mortality was 1.7% in a population with a median euroSCORE value of 5.² This suggests that the included surgical teams were able to perform both types of operations with reasonable safety.

Regarding graft patency, the question of differences in patency of proximal anastomoses is raised. As explained in an earlier reply to Smith,⁶ these differences were mainly found in the proximal parts of jump grafts that were placed on the heart. It is thus not correct to assume that the problem was seen in sites not expected to differ between the operative techniques.

The propensity to attend follow-up angiography was unevenly distributed among groups, which caused us to perform an extensive analysis of possible confounders. To quote Smith, “The authors have thoroughly and thoughtfully undertaken such an analysis, to the satisfaction of the *Journal*’s fastidious statistical reviewers. They conclude that the propensity to undergo follow-up angiography, although not perfectly balanced, did not affect the overall results or conclusions.”⁷

We therefore also believe that the conclusions are valid.

Inferior graft patency after OPCAB compared with ONCAB in DOORS confirmed the findings of numerous earlier studies.^{8–11} Even in the Surgical Management of Arterial Revascularization Therapies (SMART) study conducted by Puskas and colleagues,¹² graft patency was 84.4% after ONCAB, versus 78.4% after OPCAB. This suggests a difference on the order of magnitude of that found in DOORS, even though the small number of patients undergoing follow-up angiography in the SMART study ($n = 57$) precluded any meaningful statistical comparison.

I would not argue with Yanagawa and Puskas with regard to the superiority of OPCAB to percutaneous coronary intervention. There is, of course, solid evidence that in most cases ONCAB also performs better than percutaneous coronary intervention.^{13,14} I also agree that evidence is accumulating in favor of arterial versus venous grafts,

although there is no reason to believe that this effect should be different for OPCAB than for ONCAB.

Yanagawa and Puskas refer to early studies that have shown superiority of OPCAB relative to ONCAB regarding such surrogate endpoints as lower levels of myocardial enzymes, reduced transfusion requirements, and better scores in neurocognitive tests. They could have added subclinical markers of renal damage,^{15–17} lung function,¹⁸ inflammatory markers,¹⁶ and gastrointestinal complications.¹⁹ None of these apparent differences, however, have survived the test of being able to improve the health of patients in the large randomized, controlled, clinical trials that have been performed during the last decade.²⁰

The lower graft patency after OPCAB that we confirmed in DOORS,¹ the higher risk of postoperative cardiac death found in the ROOBY trial,³ and the worse overall long-term survival in balanced cohorts shown by Takagi and co-workers²¹ are real causes of concern. Early findings from studies without the necessary statistical power from a period with the publication bias of successful pioneers should not distract us from this fact.

Kim Houliand, MD, PhD
Department of Vascular Surgery
Kolding Hospital
Kolding, Denmark

References

- Houliand K, Fenger-Grøn M, Holme SJ, Kjeldsen BJ, Madsen SN, Rasmussen BS, et al; DOORS Study Group. Graft patency after off-pump coronary artery bypass surgery is inferior even with identical heparinization protocols: results from the Danish On-pump Versus Off-pump Randomization Study (DOORS). *J Thorac Cardiovasc Surg*. 2014;148:1812–9.e2.
- Houliand K, Kjeldsen BJ, Madsen SN, Rasmussen BS, Holme SJ, Nielsen PH, et al; DOORS Study Group. On-pump versus off-pump coronary artery bypass surgery in elderly patients. *Circulation*. 2012;125:2431–9.
- Shroyer AL, Grover FL, Hattler B, Collins JF, McDonald GO, Kozora E, et al; Veterans Affairs Randomized On/Off Bypass (ROOBY) Study Group. On-pump versus off-pump coronary-artery bypass surgery. *N Engl J Med*. 2009;361:1827–37.
- Keeling WB, Williams ML, Slaughter MS, Zhao Y, Puskas JD. Off-pump and on-pump coronary revascularization in patients with low ejection fraction: a report from the Society of Thoracic Surgeons National database. *Ann Thorac Surg*. 2013;96:83–8; discussion 88–9.
- Houliand K, Kjeldsen BJ, Madsen S, Rasmussen BS, Holme S, Schmidt TA, et al; DOORS Study Group. The impact of avoiding cardiopulmonary by-pass during coronary artery bypass surgery in elderly patients. The Danish On-pump Off-pump Randomisation Study (DOORS). *Trials*. 2009;10:47–55.
- Houliand K, Kjeldsen BJ. Definition of proximal anastomosis in jump grafts in the DOORS study. *J Thorac Cardiovasc Surg*. 2015;149:943–4.
- Smith CR. #SLOPCAB: Is the party over? *J Thorac Cardiovasc Surg*. 2014;148:1788–9.
- Khan NE, De Souza A, Mister R, Flather M, Clague J, Davies S, et al. A randomized comparison of off-pump and on-pump multivessel coronary-artery bypass surgery. *N Engl J Med*. 2004;35:21–8.
- Lingaas PS, Hol PK, Lundblad R, Rein KA, Mathisen L, Smith HJ, et al. Clinical and radiologic outcome of off-pump coronary surgery at 12 months follow-up: a prospective randomized trial. *Ann Thorac Surg*. 2006;81:2089–95.
- Widimsky P, Straka Z, Stros P, Jirasek K, Dvorak J, Votava J, et al. One-year coronary bypass graft patency: a randomized comparison between off-pump and on-pump surgery angiographic results of the PRAGUE-4 trial. *Circulation*. 2004;110:3418–23.

11. Hattler B, Messenger JC, Shroyer AL, Collins JF, Haugen SJ, Garcia JA, et al; Veterans Affairs Randomized On/Off Bypass (ROOBY) Study Group. Off-Pump coronary artery bypass surgery is associated with worse arterial and saphenous vein graft patency and less effective revascularization: results from the Veterans Affairs Randomized On/Off Bypass (ROOBY) trial. *Circulation*. 2012; 125:2827-35.
12. Puskas JD, Williams WH, O'Donnell R, Patterson RE, Sigman SR, Smith AS, et al. Off-pump and on-pump coronary artery bypass grafting are associated with similar graft patency, myocardial ischemia, and freedom from reintervention: long-term follow-up of a randomized trial. *Ann Thorac Surg*. 2011;91:1836-42; discussion 1842-3.
13. Mohr FW, Morice MC, Kappetein AP, Feldman TE, Ståhle E, Colombo A, et al. Coronary artery bypass graft surgery versus percutaneous coronary intervention in patients with three-vessel disease and left main coronary disease: 5-year follow-up of the randomised, clinical SYNTAX trial. *Lancet*. 2013;381:629-38.
14. Farkouh ME, Domanski M, Sleeper LA, Siami FS, Dangas G, Mack M, et al; FREEDOM Trial Investigators. Strategies for multivessel revascularization in patients with diabetes. *N Engl J Med*. 2012;367:2375-84.
15. Lamy A, Devereaux PJ, Prabhakaran D, Taggart DP, Hu S, Paolasso E, et al; CORONARY Investigators. Off-pump or on-pump coronary-artery bypass grafting at 30 days. *N Engl J Med*. 2012;366:1489-97.
16. Raja SG, Berg GA. Impact of off-pump coronary artery bypass surgery on systemic inflammation: current best available evidence. *J Card Surg*. 2007;22: 445-55.
17. Ascione R, Lloyd CT, Underwood MJ, Gomes WJ, Angelini GD. On-pump versus off-pump coronary revascularization: evaluation of renal function. *Ann Thorac Surg*. 1999;68:493-8.
18. Güler M, Kirali K, Tokar ME, Bozbuğa N, Omeroğlu SN, Akinci E, et al. Different CABG methods in patients with chronic obstructive pulmonary disease. *Ann Thorac Surg*. 2001;71:152-7.
19. Raja SG, Haider Z, Ahmad M. Predictors of gastrointestinal complications after conventional and beating heart coronary surgery. *Surgeon*. 2003;1:221-8.
20. Houlind K. On-pump versus off-pump coronary artery bypass surgery: what is the status after ROOBY, DOORS, CORONARY and GOPCABE? *Future Cardiol*. 2013;9:569-79.
21. Takagi H, Unemoto T. All-Literature Investigation of Cardiovascular Evidence (ALICE) Group. Worse long-term survival after off-pump than on-pump coronary artery bypass grafting. *J Thorac Cardiovasc Surg*. 2014;148:1820-9.

<http://dx.doi.org/10.1016/j.jtcvs.2015.03.026>

SURGEON VERSUS DEVICE IN AORTIC VALVE REPLACEMENT To the Editor:

We believe that in-hospital outcomes after valve replacement are at least partially related to the skill of the surgeon, the choice of valve, and the complexities of perioperative management. From a technical perspective, mechanical valve outcomes are likely to be affected by sizing mismatch in larger patients¹; from a management perspective, mechanical valve outcomes are likely to be affected by the approach to anticoagulation, which is still not well defined.² Surgical and institutional expertise and experience thus still remain important factors affecting outcomes of mechanical versus bioprosthetic valves.

Our study was based on the national cohort and included all patients who received bioprosthetic and mechanical aortic valves. Data on mortality and

FIGURE 1. In-hospital mortalities for mechanical and bioprosthetic aortic valve replacements according hospital valve volume, with patients matched by propensity score within each hospital. Patients were selected from the Nationwide Inpatient Sample in the years 1998 through 2011. Error bars represent the 95% confidence interval.

complications, such as thromboembolism, were included for each treatment modality. We found increased in-hospital mortality associated with mechanical valves relative to bioprosthetic valves, even after adjustment for all known confounders using propensity scores. Further, when patients were propensity score matched within each center, the difference in mortality persisted.³

Hospitals with lower volumes were more likely to use mechanical valves, and it is possible that quality of care in these hospitals is an unmeasured variable. The accompanying editorial written by a respected expert focuses on this aspect of the evidence.⁴ The editorial highlights that surgeon expertise, volume, and anticoagulation protocol for mechanical and biologic valves are important and can serve as surrogate markers for quality of medical care and higher mortality in institutions with lower volumes, including the “failure to rescue” for complications. We believe, however, that persistence of findings after propensity score matching within each center is evidence that factors beyond the facilities’ quality of care levels are responsible for the higher mortality associated with mechanical valves (Figure 1). Even if we assume that there are no other factors, one can still argue that this can also be a device safety issue. When the device technology is complex and requires proper protocols for sizing, placement and management that are more likely to be well implemented in institutions with higher volumes, should we really separate the effect of the device safety from delivery of care? The device manufacturers and regulators need to be aware of the issues affecting implant performance under usual care, and we believe that the complexity of care and the device should be viewed as a package from a device safety and effectiveness perspective.⁵

We also recognize that the safety issues might be related to some specific mechanical valves and not to all devices. The absence of unique device identifiers makes it extremely difficult, if not impossible, to assess manufacturer-specific product performance in the real-world setting.⁶ The Society