Authors have nothing to disclose with regard to commercial support.

Domingos Ramos De Souza, MD, PhD^a

Michael R. Dashwood, PhD^b

Ninos Samano, MD, PhD^a

^aDepartment of Cardiothoracic and Vascular Surgery
Faculty of Medicine and Health
Örebro University
Örebro, Sweden

^bSurgical and Interventional Sciences
Royal Free Hospital Campus
University College Medical School
London, United Kingdom

References

1. Benedetto U, Angelini GD. Saphenous vein graft harvesting and patency: still an unanswered question. *J Thorac Cardiovasc Surg.* 2016;152:1462-3.
2. Angelini G. Surgical interventions for veins. CTSNet: SCTS Ionescu University program at the 2015 Annual Meeting of the Society for Cardiothoracic Surgery in Great Britain and Ireland. 2016. Available at: http://www.ctsnet.org/article/surgical-interventions-veins?utm_source=iContact&utm_medium=email&utm_campaign=CTSNet&utm_content=Pulse+11%2F15%2F16. Accessed November 9, 2016.
3. Roubos N, Rosenfeldt FL, Richards SM, Conyers RA, Davis BB. Improved preservation of saphenous vein grafts by the use of glycyl trinitrate-verapamil solution during harvesting. *Circulation.* 1995;92:31-6.
4. Wise ES, Brophy CM. The case for endothelial preservation via pressure-regulated distension in the preparation of autologous saphenous vein conduits in cardiac and peripheral bypass operations. *Front Surg.* 2016;3:54.
5. Shukla N, Wan S, Angelini GD, Jeremy JY. Low nanomolar thapsigargin inhibits the replication of vascular smooth muscle cells through reversible endoplasmic reticular stress. *Eur J Pharmacol.* 2013;714:210-7.
6. Samano N, Geijer H, Liden M, Fremes S, Bodin L, Souza D. The no-touch saphenous vein for coronary artery bypass grafting maintains a patency, after 16 years, comparable to the left internal thoracic artery: a randomized trial. *J Thorac Cardiovasc Surg.* 2015;150:880-8.
7. Dashwood MR, Savage K, Tsui JC, Dooley A, Shaw SG, Alfonso MSF, et al. Retaining perivascular tissue of human saphenous vein grafts protects against surgical and distension-induced damage and preserves endothelial nitric oxide synthase and nitric oxide synthase activity. *J Thorac Cardiovasc Surg.* 2009;138:334-40.
8. Souza DS, Dashwood MR, Tsui JC, Filbey D, Bodin L, Johansson B, et al. Improved patency in vein grafts harvested with surrounding tissue: results of a randomized study using three harvesting techniques. *Ann Thorac Surg.* 2002;73:1189-95.

<http://dx.doi.org/10.1016/j.jtcvs.2017.02.010>

DISTENDED OR NOT DISTENDED? FAT OR NUDE SAPHENOUS VEIN? IN MEDIO STAT VIRTUS

Reply to the Editor:

In an editorial comment, Michael Jessen¹ argues that although most surgeons² are advocating the use of all arterial grafts for coronary artery bypass grafting, 70% of patients in the United States still

received at least 1 saphenous vein graft (SVG). In addition to the underuse of all arterial grafting, an SVG is considered a good option to graft the right coronary system, especially in case of target stenosis less than 80%³ and the bilateral internal thoracic artery is grafted to the left coronary system. Moreover, in elderly patients, the use of a single internal thoracic artery to the left anterior descending artery along with the SVG has been widely considered to achieve good outcomes.

Thus, along with Jessen,¹ we wonder why so few efforts have been made to improve SVG patency. In this scenario, the idea by Souza and coworkers^{4,5} to harvest the SVG with surrounding tissue looks appealing and promising. In fact, as brilliantly argued by the authors, with this harvesting technique there is no manipulation of the graft, preserving both adventitia and endothelium, and avoiding graft spasm. But if it is true that surrounding tissue acts as a buffer as demonstrated by Dashwood and colleagues,⁶ SVG distention, even at 300 mm Hg for 1 minute, can be dangerous for endothelial integrity.⁶ In fact, the same group evaluated the effect of graft distention on the luminal endothelium by CD31 staining. Even if conventional SVG harvesting with denudation and distention showed the highest degree of endothelium damage (4.66), veins with perivascular tissue (PVT) and distention (no-touch technique) showed CD31 staining (7.54) similar to veins without PVT and no distention (7.50), whereas veins with PVT without distention showed the lowest grade of damage (8.38). In the same study, this trend was confirmed by other analyses. Although PVT plays a protective role in the case of SVG distention, the latter is somewhat dangerous for vein endothelium and should be avoided, even in case of low pressure distention (300 mm Hg).

Finally, another issue to solve is where harvested SVG should be stored before grafting: saline solution, warm blood, or other solution. Some in vitro or animal studies are consistent in demonstrating the detrimental effects of saline on vascular endothelium and therefore graft patency, but there is no agreement to decree warm blood superior as a storage medium.⁷

We believe that the protective role of surrounding tissue in case of SVG harvesting is flawless and separated from the action of avoiding overdilation, but as in Latin, in medio stat virtus, that is, virtue stands in the middle. The best choice may be harvesting the SVG with the surrounding tissue to take advantage of its effect as a buffer, but connecting the SVG to the arterial cannula allows the vein to distend at the patient's own arterial pressure, as suggested by Angelini and coworkers.⁸

Michele Di Mauro, MD

Massimiliano Foschi, MD

Fabrizio Costantino Tancredi, MD

Gabriele Di Giammarco, MD

Department of Cardiac Surgery

SS Annunziata Hospital

Chieti, Italy

Authors have nothing to disclose with regard to commercial support.

References

1. Jessen ME. Efforts to improve bypass graft patency have not been "in vein." *J Thorac Cardiovasc Surg.* 2015;150:889-90.
2. Benedetto U, Caputo M, Zakkar M, Bryan A, Angelini GD. Are three arteries better than two? Impact of using the radial artery in addition to bilateral internal thoracic artery grafting on long-term survival. *J Thorac Cardiovasc Surg.* 2016;152:862-9.
3. Di Mauro M, Contini M, Iacò AL, Bivona A, Gagliardi M, Varone E, et al. Bilateral internal thoracic artery on the left side: a propensity score-matched study of impact of the third conduit on the right side. *J Thorac Cardiovasc Surg.* 2009;137:869-74.
4. Souza DSR, Dashwood MR, Samano N. Saphenous vein graft harvesting and patency: no-touch harvesting is the answer. *J Thorac Cardiovasc Surg.* 2017;154:1300-1.
5. Souza DS, Dashwood MR, Tsui JC, Filbey D, Bodin L, Johansson B, et al. Improved patency in vein grafts harvested with surrounding tissue: results of a randomized 102 study using three harvesting techniques. *Ann Thorac Surg.* 2002;73:1189-95.
6. Dashwood MR, Savage K, Tsui JC, Dooley A, Shaw SG, Alfonso MSF, et al. Retaining perivascular tissue of human saphenous vein grafts protects against surgical and distension-induced damage and preserves 97 endothelial nitric oxide synthase and nitric oxide synthase activity. *J Thorac Cardiovasc Surg.* 2009;138:334-40.
7. Tsakoka M, Montgomery-Taylor S, Tsakokc T. Storage of saphenous vein grafts prior to coronary artery bypass grafting: is autologous whole blood more effective than saline in preserving graft function? *Interact Cardiovasc Thorac Surg.* 2012;15:720-5.
8. Angelini GD, Breckenridge IM, Williams HM, Newby AC. A surgical preparative technique for human saphenous vein coronary artery bypass grafts which preserves medial and endothelial functional integrity. *J Thorac Cardiovasc Surg.* 1987;94:393-8.

<http://dx.doi.org/10.1016/j.jtcvs.2017.03.006>