Authors have nothing to disclose with regard to commercial support.

Authors have nothing to disclose with regard to commercial support.

with normalized left ventricular function should be assessed for donation.

Although we strongly agree with their comment, we also emphasize that young donors with or without cardiopulmonary resuscitation must be subjected to hormonal resuscitation and optimal hemodynamic management, as described in the past.⁴ Ejection fraction measured at the time of procurement, 4 days later in the experience of Galeone and colleagues,¹ remains the final deciding factor in deciding whether to use the donor heart.

Louis P. Perrault, MD, PhD
Michel Carrier, MD
Department of Cardiac Surgery
Montreal Heart Institute
Université de Montréal
Montreal, Quebec, Canada

References

1. Galeone A, Varnous S, Lebreton G, Barreda E, Hariri S, Pavie A, et al. Impact of cardiac arrest resuscitated donors on heart transplant recipients' outcome. *J Thorac Cardiovasc Surg.* 2017;153:622-30.
2. Perrault LP, Carrier M. Expanding the pool of cardiac donors: Is it really possible after cardiac arrest? *J Thorac Cardiovasc Surg.* 2017;153:631.
3. Trivedi JR, Gallo M, Slaughter MS. Use of resuscitated donor hearts for transplantation: same results either side of the Atlantic Ocean. *J Thorac Cardiovasc Surg.* 2017. In press.
4. Zaroff JG, Rosengard BR, Armstrong WF, Babcock WD, D'Alessandro A, Dec GW, et al. Consensus conference report: maximizing use of organs recovered from the cadaver donor: cardiac recommendations, March 28-29, 2001, Crystal City, Va. *Circulation.* 2002;106:836-41.

<http://dx.doi.org/10.1016/j.jtcvs.2017.04.004>

**OLD EUROPE
CAREFULLY LOOKS AT A
NEW HEART: CARDIAC
ARREST-RESUSCITATED
DONORS SHOULD NOT BE**

**TURNED DOWN FOR HEART TRANSPLANT
AT FIRST GLANCE**

Reply to the Editor:

Expansion of organ donor criteria is an essential strategy to implement the limited organ availability; reluctance exists in accepting organs for transplantation from donors who sustained a cardiac arrest because of the concern that warm ischemic injury could negatively influence graft and patient survival. We showed that history of cardiac arrest resuscitation in donors with a preserved left ventricular function at the time of organ procurement does not affect early and late outcomes of heart recipients.¹ We also found

a significantly better long-term survival in recipients receiving an organ from a cardiac arrest–resuscitated donor (CARD), probably because of the younger age of donors who sustained a cardiac arrest in our series, as well as the ischemic preconditioning effect of cardiac arrest. Donors with a history of cardiac arrest are considered as marginal donors and usually refused as heart donors. Our results demonstrated that donors with a history of cardiac arrest should not be excluded a priori from heart transplantation; however, they need to be carefully evaluated because not all CARs are suitable for heart transplant; in our experience, CARs with poor left ventricular function and higher inotropic support were obviously turned down for heart transplantation.

Previous reports similarly showed that a short period of warm ischemia due to cardiac arrest does not negatively affect recipients' outcome,^{2,3} and recently a case series has been published on 3 successful adult heart transplants from non–heart-beating donors.⁴

Expansion of donor criteria, including the use of CARs, use of non–heart-beating donors, and development of optimal preservation and perfusion techniques to minimize organ injury and improve organ recovery after cardiac arrest, is required to increase the number of organs available for heart transplantation and reduce the unacceptable discrepancy between the growing number of recipients waiting for transplantation and the ongoing shortage of organs.

Antonella Galeone, MD, PhD
Guillaume Lebreton, MD, PhD
Pascal Leprince, MD, PhD
Department of Thoracic and Cardiovascular Surgery
University Pierre et Marie Curie
Paris 6, Assistance Publique des Hôpitaux de Paris
Groupe Hospitalier Pitié-Salpêtrière
Paris, France

References

1. Galeone A, Varnous S, Lebreton G, Barreda E, Hariri S, Pavie A, et al. Impact of cardiac arrest resuscitated donors on heart transplant recipients' outcome. *J Thorac Cardiovasc Surg.* 2017;153:622-30.
2. Quader MA, Wolfe LG, Kasirajan V. Heart transplantation outcomes from cardiac arrest–resuscitated donors. *J Heart Lung Transplant.* 2013;32:1090-5.
3. Ali AA, Lim E, Thanikachalam M, Sudarshan C, White P, Parameshwar J, et al. Cardiac arrest in the organ donor does not negatively influence recipient survival after heart transplantation. *Eur J Cardiothorac Surg.* 2007;31:929-33.
4. Dhital KK, Iyer A, Connellan M, Chew HC, Gao L, Doyle A, et al. Adult heart transplantation with distant procurement and ex-vivo preservation of donor hearts after circulatory death: a case series. *Lancet.* 2015;385:2585-91.

<http://dx.doi.org/10.1016/j.jtcvs.2017.04.017>