Authors have nothing to disclose with regard to commercial support.

CONTRACTS WITH PEOPLE WHO INJECT DRUGS FOLLOWING VALVE SURGERY: UNREALISTIC AND MISGUIDED EXPECTATIONS

To the Editor:

We applaud Vlahakes' acknowledgment, in a recent editorial commentary,¹ of the importance of incorporating addiction medicine experts into the care of people with infective endocarditis caused by injection drug use. However, we take issue with his recommendations that surgeons pursue a contract with their patients who use injection drugs as motivation for them to remain clean.

The idea that a contract—an indenture between surgeon and patient—is enough to prevent illicit drug use is misguided. Substance use disorders have a complex pathophysiology, influenced heavily by comorbid psychiatric illness and socioeconomic factors. Treatment options are limited and often difficult to access. It is unrealistic to expect that a signature on a piece of paper, in a time of medical extremis, will insulate the patient against recurrent drug use. Even if the patient can grasp the gravity of prosthetic valve endocarditis, it is shortsighted to suggest that the potential for a disease will prevent use of drugs. Decades of neuropharmacology and neurobiology research demonstrate that drug dependence is defined by altered decision making. According to the fifth edition of the *Diagnostic and Statistical Manual of Mental Disorders*,² the essential feature of substance use disorder is that “the individual continues using the substance despite significant substance-related problems.”

Other behaviors, such as smoking cigarettes or caloric dietary decisions, can also cause recurrent cardiovascular disease. In our review of the literature, we have seen no editorials by surgeons on the importance of contracts with diabetics or smokers to avoid future behavioral transgressions. A smoker who comes in with angina following cardiac revascularization will not be refused treatment. A

diabetic with high sugars who has acute sternal osteomyelitis with retrosternal abscesses following cardiac surgery will not be refused debridement and reconstruction surgery.

Instead of contracting with patients to compel them to avoid injection drug use, a more useful contract is between surgeons and other inpatient clinicians with medical, psychiatric, and substance use-related services outside the hospital to assist patients with the transition from inpatient to posthospitalization care. As shown in a recent review of injection-drug-use-related endocarditis cases at 1 Boston hospital, fewer than 10% of cases were referred to medication-assisted treatment programs for buprenorphine or methadone.³ With limited posthospitalization support, it is not surprising that several patients with injection-drug-use-related endocarditis use illicit drugs again. There is room for improvement, and our focus should be on getting our patients the addiction-related services they need.

People who inject drugs represent a vulnerable, marginalized population, often with fragmented health care. Contracts dictating the availability of future life-saving care have no place in the clinician–patient relationship.

Alysse G. Wurcel, MD, MS^a

Sun Yu, MS^b

Manuel Pacheco, MD, FAPM, DFAPA^c

Kenneth Warner, MD^d

^aDivision of Geographic Medicine and Infectious Diseases

Department of Medicine

^cDepartment of Psychiatry

^dDivision of Cardiothoracic Surgery

Tufts Medical Center

^bTufts University School of Medicine

Boston, Mass

References

1. Vlahakes GJ. Consensus guidelines for the surgical treatment of infective endocarditis: the surgeon must lead the team. *J Thorac Cardiovasc Surg*. 2017;153:1259–60.
2. American Psychiatric Association. *DSM-5 Task Force. Diagnostic and statistical manual of mental disorders: DSM-5*. 5th ed. Washington: American Psychiatric Association; 2013.
3. Rosenthal ES, Karchmer AW, Theisen-Toupal J, Castillo RA, Rowley CF. Suboptimal addiction interventions for patients hospitalized with injection drug use-associated infective endocarditis. *Am J Med*. 2016;129:481–5.

<http://dx.doi.org/10.1016/j.jtcvs.2017.07.020>

The Editor welcomes submissions for possible publication in the Letters to the Editor section that consist of commentary on an article published in the Journal or other relevant issues. Authors should: • Include no more than 500 words of text, three authors, and five references. • Type with double-spacing. • See <http://jtcvs.ctsnetjournals.org/misc/fora.shtml> for detailed submission instructions. • Submit the letter electronically via jtcvs.editorialmanager.com. Letters commenting on an article published in the JTCVS will be considered if they are received within 6 weeks of the time the article was published. Authors of the article being commented on will be given an opportunity of offer a timely response (2 weeks) to the letter. Authors of letters will be notified that the letter has been received. Unpublished letters cannot be returned.

CONTRACT WITH THE PATIENT WITH INJECTION DRUG USE AND INFECTIVE ENDOCARDITIS: SURGEONS PERSPECTIVE

Reply to the Editor:

We read with interest both Vlahakes' editorial commentary¹ on the new American

Authors have nothing to disclose with regard to commercial support.

Association for Thoracic Surgery consensus guidelines for infective endocarditis (IE)² and Wurcel and colleagues' Letter to the Editor,³ "Contracts with People Who Inject Drugs Following Valve Surgery: Unrealistic and Misguided Expectations." The letter takes issue with the value of a contract with a patient who has substance use disorder (SUD) to remain clean and is directed more toward the Vlahakes editorial than toward the actual American Association for Thoracic Surgery guidelines. The wording of the recommendation in the guidelines is "Normal indications for surgery are reasonable to apply to patients who are intravenous drug users. Decision making must take the addiction into account, and management must include treatment of the addiction," a class IIa, level of evidence C, recommendation. The American Association for Thoracic Surgery guidelines do not mention using a contract, but rather they emphasize the utmost importance of addiction being "treated, mobilizing and taking advantage of all available resources and options for drug rehabilitation."

IE without treatment with antimicrobials and surgery, when needed, is almost uniformly fatal. We would consider surgery futile if the patient refused to be treated with antibiotics. Because SUD is also a fatal disorder, it is reasonable for the surgeon to require that the SUD be treated for surgery to be meaningful.

A recent study by our group showed that between 3 and 6 months after operation for IE, patients who inject drugs have a hazard of death or reoperation that is about 10 times higher than that of patients who do not inject drugs.⁴ The risk level associated with continued intravenous drug use is not comparable to that associated with such behaviors as smoking cigarettes or making caloric dietary decisions, as mentioned by Wurcel and colleagues.³ The reassurance that surgeons need and are entitled to is that the patient with SUD has been evaluated by an addiction medicine expert, addiction treatment is available and offered, and the patient has agreed to make a good faith effort to follow the treatment plan after the operation.

We do not believe that setting an expectation in writing for the achievement of addiction treatment goals is unreasonable. However, we do recognize that addiction treatment sometimes progresses irregularly. Slips can be managed and turned into learning experiences rather than being addressed punitively. The establishment of an open, accountable, and constructive therapeutic relationship at the center of treatment is essential. Although SUD inherently involves a loss of behavioral

control and relapses may occur, an experienced addiction treatment program works with a motivated (even though addicted) individual to develop new relapse prevention skills with time.⁵

Surgeons are not naïve, and we may hide behind "unrealistic and misguided expectations," but it is in good faith and with a sincere hope that all people with SUD-related IE will be able to take advantage of the second chance they get when we take care of their infections.

Syed T. Hussain, MD^a

Steven M. Gordon, MD^b

David W. Streem, MD^c

Eugene H. Blackstone, MD^{a,d}

Gösta B. Pettersson, MD, PhD^a

^aDepartment of Thoracic and Cardiovascular Surgery

Heart and Vascular Institute

^bDepartment of Infectious Disease

Medicine Institute

^cAlcohol and Drug Recovery Center

Neurological Institute

^dDepartment of Quantitative Health Sciences

Research Institute

Cleveland Clinic

Cleveland, Ohio

References

1. Vlahakes GJ. "Consensus guidelines for the surgical treatment of infective endocarditis": the surgeon must lead the team. *J Thorac Cardiovasc Surg.* 2017;153:1259-60.
2. Pettersson GB, Coselli JS, Hussain ST, Griffin B, Blackstone EH, Gordon SM, et al; AATS Surgical Treatment of Infective Endocarditis Consensus Guidelines Writing Committee. 2016 The American Association for Thoracic Surgery (AATS) consensus guidelines: surgical treatment of infective endocarditis: executive summary. *J Thorac Cardiovasc Surg.* 2017;153:1241-58.e29.
3. Wurcel AG, Yu S, Pacheco M, Warner K. Contracts with people who inject drugs following valve surgery: unrealistic and misguided expectations. *J Thorac Cardiovasc Surg.* 2017;154:2002.
4. Shrestha NK, Jue J, Hussain ST, Jerry JM, Pettersson GB, Menon V, et al. Injection drug use and outcomes after surgical intervention for infective endocarditis. *Ann Thorac Surg.* 2015;100:875-82.
5. McLellan AT, Schwarzlose J, for the Betty Ford Institute Consensus Panel. What is recovery? A working definition from the Betty Ford Institute. *J Subst Abuse Treat.* 2007;33:221-8.

<http://dx.doi.org/10.1016/j.jtcvs.2017.08.004>

**IS RIGID STERNAL
FIXATION REALLY
BETTER? WAS THE
CORRECT CONTROL
GROUP USED?**

To the Editor:

Allen and colleagues¹ report a nicely performed randomized, multicenter trial comparing sternal closure with rigid plate fixation to wire cerclage. Unfortunately, the criteria for the wire