^gDepartment of Radiation Oncology
University of California
Davis, Sacramento, Calif

^hDepartment of Radiation Oncology
Cleveland Clinic
Cleveland, Ohio

and ⁱDepartment of Radiation Oncology
Henry Ford Cancer Institute
Detroit, Mich

References

1. Schaheen LW, D'Cunha J. The slippery slope of nonoperative therapy in early-stage lung cancer. *J Thorac Cardiovasc Surg*. 2017;154:1121-2.
2. Swift D. Zapping tumor effective for stage I non-small cell lung cancer. May 19, 2015. Available at: <http://www.medpagetoday.com/HematologyOncology/LungCancer/51630>. Accessed September 25, 2017.
3. Videtic GM, Donington J, Giuliani M, Heinzerling J, Karas TZ, Kelsey CR, et al. Stereotactic body radiation therapy for early-stage non-small cell lung cancer: executive summary of an ASTRO Evidence-Based Guideline. *Pract Radiat Oncol*. 2017;7:295-301.
4. Moghanaki D, Chang JY. Is surgery still the optimal treatment for stage I non-small cell lung cancer? *Transl Lung Cancer Res*. 2016;5:183-9.
5. Samson P, Keogan K, Crabtree T, Colditz G, Broderick S, Puri V, et al. Interpreting survival data from clinical trials of surgery versus stereotactic body radiation therapy in operable stage I non-small cell lung cancer patients. *Lung Cancer*. 2017;103:6-10.
6. Chang JY, Senan S, Paul MA, Mehran RJ, Louie AV, Balter P, et al. Stereotactic ablative radiotherapy versus lobectomy for operable stage I non-small-cell lung cancer: a pooled analysis of two randomised trials. *Lancet Oncol*. 2015;16:630-7. Erratum in: *Lancet Oncol*. 2015;16:e427.
7. Morrison R, Deeley TJ, Cleland WP. The treatment of carcinoma of the bronchus: a clinical trial to compare surgery and supervoltage radiotherapy. *Lancet*. 1963;281:683-4.
8. Yerokun BA, Yang CJ, Gulack BC, Li X, Mulvihill MS, Gu L, et al. A national analysis of wedge resection versus stereotactic body radiation therapy for stage IA non-small cell lung cancer. *J Thorac Cardiovasc Surg*. 2017;154:675-86.e4.
9. D'Cunha J, Herndon JE II, Herzan DL, Patterson GA, Kohman LJ, Harpole DH, et al; Cancer and Leukemia Group B. Poor correspondence between clinical and pathologic staging in stage I non-small cell lung cancer: results from CALGB 9761, a prospective trial. *Lung Cancer*. 2005;48:241-6.
10. Boffa DJ, Kosinski AS, Paul S, Mitchell JD, Onaitis M. Lymph node evaluation by open or video-assisted approaches in 11,500 anatomic lung cancer resections. *Ann Thorac Surg*. 2012;94:347-53; discussion 353.
11. Heineman DJ, Ten Berge MG, Daniels JM, Versteegh MI, Marang-van de Mheen PJ, Wouters MW, et al. Clinical staging of stage I non-small cell lung cancer in the Netherlands—need for improvement in an era with expanding nonsurgical treatment options: data from the Dutch Lung Surgery Audit. *Ann Thorac Surg*. 2016;102:1615-21.
12. Louie AV, Palma DA, Dahele M, Rodrigues GB, Senan S. Management of early-stage non-small cell lung cancer using stereotactic ablative radiotherapy: controversies, insights, and changing horizons. *Radiother Oncol*. 2014;114:138-47.
13. Uematsu M, Shioda A, Suda A, Fukui T, Ozeki Y, Hama Y, et al. Computed tomography-guided frameless stereotactic radiotherapy for stage I non-small cell lung cancer: a 5-year experience. *Int J Radiat Oncol Biol Phys*. 2001;51:666-70.
14. Lagerwaard FJ, Versteegen NE, Haasbeek CJ, Slotman BJ, Paul MA, Smit EF, et al. Outcomes of stereotactic ablative radiotherapy in patients with potentially operable stage I non-small cell lung cancer. *Int J Radiat Oncol Biol Phys*. 2012;83:348-53.
15. Nagata Y, Hiraoka M, Shibata T, Onishi H, Kokubo M, Karasawa K, et al. Prospective trial of stereotactic body radiation therapy for both operable and inoperable T1N0M0 non-small cell lung cancer: Japan Clinical Oncology Group study JCOG0403. *Int J Radiat Oncol Biol Phys*. 2015;93:989-96.
16. Eriquchi T, Takeda A, Sanuki N, Tsurugai Y, Aoki Y, Oku Y, et al. Stereotactic body radiotherapy for operable early-stage non-small cell lung cancer. *Lung Cancer*. 2017;109:62-7.
17. Shibamoto Y, Hashizume C, Baba F, Ayakawa S, Miyakawa A, Murai T, et al. Stereotactic body radiotherapy using a radiobiology-based regimen for stage I

non-small-cell lung cancer: five-year mature results. *J Thorac Oncol*. 2015;10:960-4.

18. Onishi H, Shirato H, Nagata Y, Hiraoka M, Fujino M, Gomi K, et al. Stereotactic body radiotherapy (SBRT) for operable stage I non-small-cell lung cancer: can SBRT be comparable to surgery? *Int J Radiat Oncol Biol Phys*. 2011;81:1352-8.
19. Haque W, Szeja S, Tann A, Kalra S, Teh BS. Changes in treatment patterns and overall survival in patients with early-stage non-small cell lung cancer in the united states after the incorporation of stereotactic ablative radiation therapy: a population-based analysis. *Am J Clin Oncol*. January 14, 2016 [Epub ahead of print].

<http://dx.doi.org/10.1016/j.jtcvs.2017.09.001>

THE BURDEN OF PROOF

Reply to the Editor:

The erudite letter from the ASTRO Evidence-Based Guideline Committee for Stereotactic Body Radiation Therapy (SBRT) for Early-Stage Non-Small Cell Lung Cancer (ASTRO Group)¹ carries a title that all clinicians managing lung cancer could easily agree on. Conversely, the style of the letter is typical of a diplomatic effort that may not to stand the test of time because of the underlying biases of both radiation oncologists and thoracic surgeons.

Surgeons should be glad to learn that the ASTRO Group¹ agrees that the currently available evidence does not permit patients to draw crucial information regarding their own health management pathway independently from the intricacies of the literature. Better late than never—at this time, a balanced perspective toward which procedure is indicated is needed as clinicians clearly weigh the evidence in the literature differently depending on their specialty.²

In addition, thoracic surgeons keep on retrospectively analyzing large databases with propensity score matching to compare wedge and lobectomy with SBRT.¹ Although this statistical methodology is imperfect, radiation oncologists have also used the same methodology to the benefit of SBRT.³

Moreover, thoracic surgeons are appropriately not convinced by reading the SBRT literature, which is filled with failed randomized trials and related mathematical acrobacies.⁴ At the same time, there are diverse interpretations of SBRT—SBRT for peripheral,⁵ SBRT for central,⁶ and SBRT for ultracentral lung tumors.⁷ In fact, there is one SBRT approach for lung tumors larger than 5 cm (cT3) as opposed to smaller lesions⁸—a major leap “forward” compared with the “no-fly zone” concept.⁹ Of course, thoracic surgeons adapt the extent of resection as radiation oncologists modulate SBRT dosage in an attempt to strike a balance between the needs for oncologic completeness and reduced mortality and morbidity.⁹ If radiotherapists do what surgeons do, it means that they want to avoid similar problems¹⁰⁻¹³ that are underreported in the SBRT literature, with potentially harmful consequences to patients.^{14,15}

Author has nothing to disclose with regard to commercial support.

In the context of a multidisciplinary evaluation of a patient with lung cancer, the presence and the value of a thoracic surgeon are rightfully acknowledged by our Colleagues of the ASTRO Group.¹ In contrast, the trend toward indiscriminately referring to the literature of the same specialty over and over again—indeed very common among surgeons as well—inevitably emerges when an important statement is made by the ASTRO Group,¹ such as “we have reached a point in time when surgery is not the only treatment choice.” This statement is supported neither by a multicentric, prospective and randomized trial nor by a rigorous propensity score matching analysis of a retrospective series, but by an expert review article.^{1,16}

An excessively partisan approach, from whichever side it originates, should not contribute to hiding a fundamental truth. By current standards—that is, the adoption of the TNM international staging system for lung cancer and the assessment of the conventional 5-year posttreatment results—surgery for early-stage lung cancer is associated with outcomes that are unsurpassed by other treatments and reflect several hundred thousand patients operated on worldwide in the last decades.¹⁷ If the superiority of surgery is to be questioned, the burden of proof thus rests upon the shoulders of radiation oncologists. To avoid dissipating time, financial, and human resources with other useless trials on the value of SBRT in the treatment of early lung cancer independently run by radiation oncologists with a minor surgical participation, the collaboration with surgeons becomes mandatory and requires reciprocal—and unbiased—knowledge and scrutiny of the available evidence finally to throw our mutual, heavy cultural ballast overboard for the benefit of our patients. The time has finally come for a coordinated and cross-specialty effort involving all major professional societies on both sides and aiming at carefully designing a single large, international study conducted by thoracic surgeons and radiation oncologists with equal prerogatives.

Gaetano Rocco, MD, FRCSEd
Division of Thoracic Surgery
Thoracic Department
Istituto Nazionale Tumori
IRCCS-Fondazione G. Pascale
Naples, Italy

References

1. Moghanaki D, Simone CB II, Rimner A, Karas TZ, Donington J, Shirvani SM, et al; ASTRO Evidence-Based Guideline Committee for Stereotactic Body Radiation Therapy (SBRT) for Early-Stage Non-Small Cell Lung Cancer. The value of collaboration between thoracic surgeons and radiation oncologists while awaiting evidence in operable stage I non-small cell lung cancer. *J Thorac Cardiovasc Surg*. 2018;155:429-31.
2. Hopmans W, Damman OC, Porsius JT, Zwaan L, Senan S, Timmermans DR. Treatment recommendations by clinicians in stage I non-small cell lung cancer: a study of factors that influence the likelihood of accounting for the patient's preference. *Patient Educ Couns*. 2016;99:1808-13.
3. Shirvani SM, Jiang J, Chang JY, Welsh JW, Gomez DR, Swisher S, et al. Comparative effectiveness of 5 treatment strategies for early-stage non-small cell lung cancer in the elderly. *Int J Radiat Oncol Biol Phys*. 2012;84:1060-70.
4. Chang JY, Senan S, Paul MA, Mehran RJ, Louie AV, Balter P, et al. Stereotactic ablative radiotherapy versus lobectomy for operable stage I non-small-cell lung cancer: a pooled analysis of two randomised trials. *Lancet Oncol*. 2015;16:630-7.
5. Guckenberger M, Andratschke N, Dieckmann K, Hoogeman MS, Hoyer M, Hurkmans C, et al. ESTRO ACROP consensus guideline on implementation and practice of stereotactic body radiotherapy for peripherally located early stage non-small cell lung cancer. *Radiother Oncol*. 2017;124:11-7.
6. Park HS, Harder EM, Mancini BR, Decker RH. Central versus peripheral tumor location: influence on survival, local control, and toxicity following stereotactic body radiotherapy for primary non-small-cell lung cancer. *J Thorac Oncol*. 2015;10:832-7.
7. Modh A, Rimner A, Williams E, Foster A, Shah M, Shi W, et al. Local control and toxicity in a large cohort of central lung tumors treated with stereotactic body radiation therapy. *Int J Radiat Oncol Biol Phys*. 2014;90:1168-76.
8. Tekatli H, Haasbeek N, Dahele M, De Haan P, Verbakel W, Bongers E, et al. Outcomes of hypofractionated high-dose radiotherapy in poor-risk patients with “ultracentral” non-small cell lung cancer. *J Thorac Oncol*. 2016;11:1081-9.
9. Rocco G. Narcissus, the beam, and lung cancer. *J Thorac Cardiovasc Surg*. 2016;152:338-43.e3.
10. Haseltine JM, Rimner A, Gelblum DY, Modh A, Rosenzweig KE, Jackson A, et al. Fatal complications after stereotactic body radiation therapy for central lung tumors abutting the proximal bronchial tree. *Pract Radiat Oncol*. 2016;6:e27-33.
11. Din SU, Williams EL, Jackson A, Rosenzweig KE, Wu AJ, Foster A, et al. Impact of fractionation and dose in a multivariate model for radiation-induced chest wall pain. *Int J Radiat Oncol Biol Phys*. 2015;93:418-24.
12. Nambu A, Onishi H, Aoki S, Tominaga L, Kuriyama K, Araya M, et al. Rib fracture after stereotactic radiotherapy for primary lung cancer: prevalence, degree of clinical symptoms, and risk factors. *BMC Cancer*. 2013;13:68.
13. Kim SS, Song SY, Kwak J, Ahn SD, Kim JH, Lee JS, et al. Clinical prognostic factors and grading system for rib fracture following stereotactic body radiation therapy (SBRT) in patients with peripheral lung tumors. *Lung Cancer*. 2013;79:161-6.
14. Lo AC, Liu M, Chan E, Lund C, Truong PT, Loewen S, et al. The impact of peer review of volume delineation in stereotactic body radiation therapy planning for primary lung cancer: a multicenter quality assurance study. *J Thorac Oncol*. 2014;9:527-33.
15. Oskan F. The quality of toxicity reporting and the story of the lung SBRT “no-fly zone” *Int J Radiat Oncol Biol Phys*. 2015;92:514-5.
16. Moghanaki D, Chang JY. Is surgery still the optimal treatment for stage I non-small cell lung cancer? *Transl Lung Cancer Res*. 2016;5:183-9.
17. Boily G, Filion É, Rakovich G, Kopec N, Tremblay L, Samson B, et al; Comité de l'évolution des pratiques en oncologie. Stereotactic ablative radiation therapy for the treatment of early-stage non-small-cell lung cancer: CEPO review and recommendations. *J Thorac Oncol*. 2015;10:872-82.

<http://dx.doi.org/10.1016/j.jtcvs.2017.09.065>