Authors have nothing to disclose with regard to commercial support.

4. Vogelmeier CF, Criner GJ, Martinez FJ, Anzueto A, Barnes PJ, Bourbeau J, et al. Global strategy for the diagnosis, management, and prevention of chronic obstructive lung disease 2017 report. GOLD executive summary. *Am J Respir Crit Care Med*. 2017;195:557-82.

<https://doi.org/10.1016/j.jtcvs.2018.10.087>

TAKING UP THE CHRONIC PULMONARY OBSTRUCTIVE DISEASE GAUNTLET

Reply to the Editor:

The letter to editor by Medalion and colleagues provides important additional insights in regard to the article by Viceconte and colleagues.¹ The authors have identified potential perioperative confounding factors, especially pain, which could influence forced expiratory volume in 1 second and forced vital capacity assessment the time points studied. This is perhaps a more likely scenario on the second postoperative day; however, it is unclear whether pain would have sufficient impact on postoperative day 5 and beyond, as patients do tend to report lesser pain issues over time.² In future studies, in addition to the inclusion of a more comprehensive measure of pulmonary function, we would recommend the use of a valid measure of pain to coincide with postoperative assessments.

The authors of the letter to editor have previously reported long-term mortality in patients with chronic pulmonary obstructive disease to be lower several years after cardiac surgery.³ This serves to highlight the need to longitudinal follow-up in prospective studies for this patient population. Further, the inclusion of data on pre- and postoperative frailty would provide valuable insight of additive morbidity and mortality in patients severe chronic pulmonary obstructive disease and decreased left ventricular ejection fraction after cardiac surgery.^{4,5} Finally, although the traditional metrics of survival and rehospitalization are essential endpoints, we and others have also advocated for the use of longitudinal health-related quality of life assessments as a key patient-centered outcome, particularly in the older, frail adult patient.^{4,5} What the “ideal” long-term health-related quality of life tool(s) that is accurate and clinically feasible, however, requires further study.

We support the statement by Medalion and Katz on the timeliness of this paper as a “call to arms” on the need to

Dr Arora has received an unrestricted educational grant from Pfizer Canada, Inc, and honoraria from Mallinckrodt Pharmaceuticals and Abbott Nutrition for work unrelated to the manuscript. All other authors have nothing to disclose with regard to commercial support.

develop more detailed enhanced recovery protocols for these complex patients. We would further also advocate for the ongoing close collaboration between the pulmonologist and cardiac surgeon to extend beyond perioperative planning and include the transition to the community phase of the patient journey in this vulnerable population.

Zubair Luqman, MD
Rakesh C. Arora, MD, PhD, FRCSC
Department of Surgery
Max Rady College of Medicine
University of Manitoba
Winnipeg, Manitoba, Canada
Cardiac Sciences Program
St Boniface Hospital
Winnipeg, Manitoba, Canada

References

1. Viceconte M, Rocco IS, Pauletti HO, Vidotto M, Arena R, Gomes WJ, et al. Chronic obstructive pulmonary disease severity influences outcomes after off-pump coronary artery bypass. *J Thorac Cardiovasc Surg*. 2018;156:1554-61.
2. Matute-Jiménez P, Roca-Obrador J, Valero-Castell R, Gomar-Sancho C. Pain locations in the postoperative period after cardiac surgery: chronology of pain and response to treatment (abstract). *J Cardiothorac Vasc Anesth*. 2017;31:S45.
3. Medalion B, Katz MG, Cohen AJ, Hauptman E, Sasson L, Schachner A. Long-term beneficial effect of coronary artery bypass grafting in patients with COPD. *Chest*. 2004;125:56-62.
4. Yanagawa B, Graham MM, Afilalo J, Hassan A, Arora RC. Frailty as a risk predictor in cardiac surgery: beyond the eyeball test. *J Thorac Cardiovasc Surg*. September 5, 2018 [Epub ahead of print].
5. Lytwyn J, Stammers AN, Kehler DS, Jung P, Alexander B, Hiebert BM, et al. The impact of frailty on functional survival in patients 1 year after cardiac surgery. *J Thorac Cardiovasc Surg*. 2017;154:1990-9.

<https://doi.org/10.1016/j.jtcvs.2018.12.017>

IT'S THE LUNGS, NOT THE PUMP: CHRONIC OBSTRUCTIVE PULMONARY DISEASE SEVERITY AND OUTCOMES AFTER OFF-PUMP CORONARY

ARTERY BYPASS GRAFTING

Reply to the Editor:

In this issue of the *Journal*, Medalion and Katz¹ made some observations on the recent study by Viceconte and colleagues,² in which they examined the effects of the

Author has nothing to disclose with regard to commercial support.

severity of chronic obstructive pulmonary disease (COPD) on postoperative outcomes after off-pump coronary artery bypass (OPCAB).

I agree with Medalion and Katz¹ that there is no advantage of using OPCAB to significantly improve clinical outcomes in patients with COPD. Of note, findings from major clinical trials and retrospective studies indicate either noninferiority or superiority of on-pump coronary artery bypass compared with OPCAB.³ Furthermore, whether OPCAB would yield better outcomes in patients with severe COPD is debatable. In a substudy of the Randomized On/Off Bypass trial,⁴ patients with COPD who underwent OPCAB had more intraoperative complications and similar 30-day or 1-year outcomes when compared with those who underwent on-pump coronary artery bypass. Moreover, O'Boyle and colleagues⁵ reported that OPCAB had no impact on long-term survival in patients with COPD as defined by the Global Initiative for Lung Disease (GOLD).

Medalion and Katz¹ make the point that spirometry results alone do not give a clear picture of the clinical severity of the illness, and other clinical factors should be considered. Currently, as per the GOLD report,^{6,7} 3 criteria must be met for a COPD diagnosis: (1) a postbronchodilator forced expiratory volume in 1 second/forced vital capacity ratio less than 0.70, suggesting persistent airflow limitation; (2) appropriate symptoms, dyspnea, cough, and sputum production; and (3) significant exposure to noxious stimuli. In their study, Viceconte and colleagues² reported that their patients had been diagnosed with COPD and that the GOLD criteria were used to assess disease severity. Although symptoms are important and screening spirometry is not recommended in the absence of symptoms, uniformity across studies requires the use of a standard definition of disease severity. Historically, the definition of COPD has varied significantly, which has made meaningful interpretation and comparison between studies impractical. The GOLD criteria have been updated to account for disease severity; however, that does not undermine the importance of spirometric findings.

In the study by Viceconte and colleagues,² spirometry was conducted preoperatively and on postoperative days 2 and 5. The authors noted a correlation between severe COPD based on the GOLD criteria and impaired pulmonary function on postoperative days 2 and 5, with an increased risk of composite outcomes with worsening

COPD severity. I agree with Medalion and Katz¹ that pain, fluid overload, and systemic inflammation resulting from surgical manipulation affect the measurement of early postoperative spirometric findings. It is likely these early impairments in spirometry, although temporary, are critically important and contribute significantly to poor outcomes. However, it is unknown whether these immediate postoperative spirometry findings revert to preoperative values on longer follow-up or whether they affect long-term survival. Multiple studies have shown that COPD adversely affects survival after coronary artery bypass grafting; however, most clinicians believe that coronary artery bypass grafting provides survival benefits over medical management in such patients.

Medalion and Katz¹ rightly suggest that the study by Viceconte and colleagues² would be stronger had they performed a more comprehensive evaluation of other common postoperative complications, including readmissions, mortality, and survival data. Perhaps a longer-term follow-up study would address this issue.

No grading system is perfect, and the GOLD criteria have limitations; however, the criteria help achieve uniformity in results across studies, and their use will aid in standardizing the definition of COPD.

Shuab Omer, MD

Division of Cardiothoracic Surgery

Baylor College of Medicine

Houston, Tex

Michael E. DeBakey VA Medical Center

Houston, Tex

References

1. Medalion B, Katz MG. Chronic obstructive pulmonary disease severity influences outcomes after off-pump coronary artery bypass. *J Thorac Cardiovasc Surg.* 2019; 157:e191-2.
2. Viceconte M, Rocco IS, Pauletti HO, Vidotto M, Arena R, Gomes WJ, et al. Chronic obstructive pulmonary disease severity influences outcomes after off-pump coronary artery bypass. *J Thorac Cardiovasc Surg.* 2018;156: 1554-61.
3. Lazar HL. Should off-pump coronary artery bypass grafting be abandoned? *Circulation.* 2013;128:406-13.
4. Almassi GH, Shroyer AL, Collins JF, Hattler B, Bishawi M, Baltz JH, et al. Chronic obstructive pulmonary disease impact upon outcomes: the veterans affairs randomized on/off bypass trial. *Ann Thorac Surg.* 2013; 96:1302-9.
5. O'Boyle F, Mediratta N, Chalmers J, Al-Rawi O, Mohan K, Shaw M, et al. Long-term survival of patients with pulmonary disease undergoing coronary artery bypass surgery. *Eur J Cardiothorac Surg.* 2013;43:697-703.
6. Mirza S, Clay RD, Koslow MA, Scanlon PD. COPD guidelines: a review of the 2018 GOLD report. *Mayo Clin Proc.* 2018;93:1488-502.
7. Vogelmeier CF, Criner GJ, Martinez FJ, Anzueto A, Barnes PJ, Bourbeau J, et al. Global strategy for the diagnosis, management, and prevention of chronic obstructive lung disease 2017 report. GOLD executive summary. *Am J Respir Crit Care Med.* 2017;195:557-82.

<https://doi.org/10.1016/j.jtcvs.2018.10.123>