Dr Rocha has nothing to disclose with regard to commercial support. Dr Tam is supported by a CIHR Fellowship, Canada. Dr Fremes is supported by the Bernard S. Goldman Chair in Cardiovascular Surgery Toronto, Ontario, Canada.

artery bypass grafting in patients with versus without preoperative β -blocker therapy. No differences in mortality, morbidity, or length of stay were observed between groups. On the basis of the lack of early β -blockade benefit in coronary artery bypass grafting cases, Schubert and associates² evaluated the role of β -blocker therapy in patients requiring surgical aortic valve replacement (SAVR), in which its potential benefit was less certain. They compared the short-term outcomes of 4592 propensity score-matched patients undergoing isolated SAVR with versus without preoperative β -blockade, from 18 cardiac surgery centers across the Commonwealth of Virginia during a period of 14 years.² In their comparison, preoperative β -blocker use was associated with increased rates of postoperative renal failure requiring dialysis, transfusion, atrial fibrillation, and cardiac arrest, and with a longer intensive care unit stay. The investigators concluded that β -blocking agents should not be used as a quality metric in SAVR.

In their letter to the Editor, Mazzeffi and colleagues³ appropriately highlight their concerns with the conclusions of the study of Schubert and colleagues.² Limitations of administrative database studies include data granularity and, in this case, a paucity of information regarding the duration, indication, dosage, and type of β -blocker therapy. Mazzeffi and colleagues³ have highlighted the differences in pharmacology related to different β -blockers, which may have an important effect on outcome. Importantly, the retrospective nature of the data inevitably introduces confounding into the analysis. Despite the use of propensity score matching to balance multiple preoperative variables, there may still be residual unmeasured confounders in the results after matching. For instance, the longer aortic crossclamp time in the group of patients receiving β -blockers, despite similar incidence of aortic annular enlargement, suggests that the procedures in this group may have been more complex. Increased cardiopulmonary bypass time has been shown to be an independent risk factor for postoperative complications, such as renal failure requiring dialysis and transfusion.^{4,5} Moreover, the higher rate of postoperative atrial fibrillation, despite β -blockade, may likely reflect a potentially sicker group of patients. Echocardiographic data, on risk factors for atrial fibrillation, such as left ventricular or atrial dimensions, were not available.⁶

After propensity score matching, 19 comparisons were performed, which poses a risk for multiplicity and type I error. Common strategies to address multiplicity are the Bonferroni correction or false discovery rate. These techniques could have been considered when performing such a high number of comparisons. Nevertheless, this hypothesis-generating study adds important data to the cardiac surgery literature and sets the stage for a randomized, controlled trial, which may properly address the question of whether patients undergoing SAVR should receive preoperative β -blockers.

Rodolfo V. Rocha, MD^a

Derrick Y. Tam, MD^{a,b}

Stephen E. Fremes, MD, MSc^{a,b}

^aDivision of Cardiac Surgery

Department of Surgery

Schulich Heart Centre

Sunnybrook Health Sciences Centre

University of Toronto

Toronto, Ontario, Canada

^bInstitute of Health Policy, Management and Evaluation

University of Toronto

Toronto, Ontario, Canada

References

1. LaPar DJ, Crosby IK, Kron IL, Kern JA, Fonner E Jr, Rich JB, et al. Preoperative beta-blocker use should not be a quality metric for coronary artery bypass grafting. *Ann Thorac Surg*. 2013;96:1539-44; discussion 1544-5.
2. Schubert S, Hawkins R, Mehaffey J, Fonner CE, Rich JB, Speir AM, et al. Preoperative β -blocker use correlates with worse outcomes in patients undergoing aortic valve replacement. *J Thorac Cardiovasc Surg*. 2019;158:1589-97.
3. Mazzeffi MA, Evans CF. β -Blockers in aortic valve surgery: beta safe or sorry? *J Thorac Cardiovasc Surg*. 2019. In press.
4. Chertow GM, Lazarus JM, Christiansen CL, Cook EF, Hammermeister KE, Grover F, et al. Preoperative renal risk stratification. *Circulation*. 1997;95:878-84.
5. Scott BH, Seifert FC, Glass PS, Grimson R. Blood use in patients undergoing coronary artery bypass surgery: impact of cardiopulmonary bypass pump, hematocrit, gender, age, and body weight. *Anesth Analg*. 2003;97:958-63, table of contents.
6. Tiwari S, Schirmer H, Jacobsen BK, Hopstock LA, Nyrnes A, Heggelund G, et al. Association between diastolic dysfunction and future atrial fibrillation in the Tromsø Study from 1994 to 2010. *Heart*. 2015;101:1302-8.

<https://doi.org/10.1016/j.jtcvs.2019.05.076>

REPLY: BETA-BLOCKER HOUSE REVISITED!

Reply to the Editor:

Mazzeffi and Evans¹ raise several red flags regarding the conclusions offered by Schubert and colleagues.²

As I mentioned in my own comment,³ the duration and selectivity of the preoperative beta-blocker were not documented. The percentage of patients receiving preoperative beta-blocker therapy solely as a quality metric was also not noted. It is likely that a majority of patients received nonselective beta-blockers, a factor that

Author has nothing to disclose with regard to commercial support.

Authors have nothing to disclose with regard to commercial support.

can influence specific postreplacement outcomes. Mazzeffi and Evans¹ remind us that these caveats are reasons why preoperative beta-blocker therapy might not be responsible for the increased rates of atrial fibrillation, transfusion, and renal dialysis seen after aortic valve replacement.

My impression is that preoperative beta-blocker therapy was a marker for sicker patients undergoing aortic valve replacement. Even in the propensity-matched cohorts, the patients taking a beta-blocker medication had an increased incidence of coronary artery disease, prior coronary artery bypass grafting procedure, renal dialysis, and cerebral vascular accidents.

Patients undergoing transcatheter aortic valve replacement during the study period were at moderate to high risk and consequently likely to have been receiving a beta-blocker medication. Those patients were excluded from the propensity-matched aortic valve replacement cohorts and potentially influenced the authors' conclusions.

A carefully planned study controlling the duration and selectivity of the preoperative beta-blocker medication will be required before we can conclude that adding beta-blocker therapy to elective aortic valve replacement protocols affects outcomes.

Frank A. Baciewicz, Jr, MD
Division Cardiothoracic Surgery
Department of Surgery
Wayne State University School of Medicine
Detroit, Mich

References

1. Mazzeffi M, Evans C. Beta-blockers in aortic valve surgery: beta safe or sorry? *J Thorac Cardiovasc Surg.* 2020;159:e41.
2. Schubert SA, Hawkins RB, Mehaffey JH, et al. Preoperative beta-blocker use correlates with worse outcomes in patients undergoing aortic valve replacement. *J Thorac Cardiovasc Surg.* 2019;158:1589-97.
3. Baciewicz F. All members of Beta Blocker House on probation! *J Thorac Cardiovasc Surg.* 2019;158:1598-9.

<https://doi.org/10.1016/j.jtcvs.2019.06.010>

REPLY: BETA OR WORSE: MORE WORK NEEDED TO DETERMINE BENEFIT OR HARM IN AORTIC VALVE SURGERY

Reply to the Editor:

The letter by Mazzeffi and colleagues¹ regarding the

article by Shubert and associates² highlights and adds to the areas in need of further qualification and confirmation that we described in our editorial.³ Although we did not bring up this important statistical point, there were enough points of comparison in this database analysis that there was likely a type I error that led to the increased incidence of postoperative atrial fibrillation in the patients who received preoperative β -blocker therapy. We also pointed out that there were no data regarding whether patients who were receiving a β -blocker preoperatively had it continued postoperatively, and we agree with the lack of information regarding the type of β -blocker used, which may have had a significant effect on the outcomes.⁴ We pointed out that 27% of the patients had moderate to severe aortic insufficiency and that Shubert and associates² should have verified that the heart rates were kept at 70 beats/min or faster before the operation to prevent the patient from entering the operation with compromised ventricles.⁵ We emphasized that when preoperative β -blockers are continued postoperatively, the dose and drug type should be chosen on an individual basis, and adequate heart rate and blood pressure should be maintained to prevent renal dysfunction, the need for transfusion, and new-onset atrial fibrillation. Finally, we agree that more studies addressing this question are needed to guide clinical practice.

Vincent Conti, MD
Gal Levy, MD
Department of Surgery
University of Texas Medical Branch at Galveston
Galveston, Tex

References

1. Mazzeffi MA, Evans CF. Beta blockers in aortic valve surgery: beta safe or sorry? *J Thorac Cardiovasc Surg.* 2020;159:e42-3.
2. Schubert SA, Hawkins RB, Mehaffey JH, Chancellor WZ, Fonner CE, Speir AM, et al. Preoperative β -blocker use correlates with worse outcomes in patients undergoing aortic valve replacement. *J Thorac Cardiovasc Surg.* 2019;158:1589-97.
3. Levy G, Conti VR. Commentary: preoperative β -blockers—one size does not fit all when considering management. *J Thorac Cardiovasc Surg.* 2019;158:1600-1.
4. Kertai MD, Esper SA, Akushevich I, Voora D, Ginsburg GS, Stafford-Smith M, et al. Preoperative CYP2D6 metabolism-dependent beta-blocker use and mortality after coronary artery bypass grafting surgery. *J Thorac Cardiovasc Surg.* 2014;147:1368-75.e3.
5. Sampat U, Varadarajan P, Turk R, Kamath A, Khandhar S, Pai RG. Effect of beta-blocker therapy on survival in patients with severe aortic regurgitation results from a cohort of 756 patients. *J Am Coll Cardiol.* 2009;54:452-7.

<https://doi.org/10.1016/j.jtcvs.2019.05.027>