FRACS (Department of Cardiothoracic Surgery, The Perth Children's Hospital, Perth, Western Australia, Australia).

References

- Lillehei CW, Cohen M, Warden HE, Varco RL. The direct-vision intracardiac correction of congenital anomalies by controlled cross circulation; results in thirty-two patients with ventricular septal defects, tetralogy of Fallot, and atrioventricular communis defects. *Surgery*. 1955;38:11-29.
- Pan G, Song L, Zhou X, Zhao J. Complete atrioventricular septal defect: comparison of modified single-patch technique with two-patch technique in infants. *J Card Surg*. 2014;29:251-5.
- Yildirim O, Avsar M, Ozyuksel A, Akdemir M, Zeybek C, Demiroglu S, et al. Modified single versus double-patch technique for the repair of complete atrioventricular septal defect. *J Card Surg*. 2015;30:595-600.
- Ugaki S, Khoo NS, Ross DB, Rebecka IM, Adatia I. Modified single-patch compared with two-patch repair of complete atrioventricular septal defect. *Ann Thorac Surg*. 2014;97:666-71.
- Wilcox BR, Jones DR, Frantz EG, Brink LW, Henry GW, Mill MR, et al. Anatomically sound, simplified approach to repair of "complete" atrioventricular septal defect. *Ann Thorac Surg*. 1997;64:487-93.
- Nicholson IA, Nunn GR, Sholler GF, Hawker RE, Cooper SG, Lau KC, et al. Simplified single patch technique for the repair of atrioventricular septal defect. *J Thorac Cardiovasc Surg*. 1999;118:642-6.
- Backer CL, Stewart RD, Mavroudis C. What is the best technique for repair of complete atrioventricular canal? *Semin Thorac Cardiovasc Surg*. 2007;19:249-57.
- Fong LSW, Winlaw DS, Orr Y. Is the modified single-patch repair superior to the double-patch repair of complete atrioventricular septal defects? *Interact Cardiovasc Thorac Surg*. 2019;28:427-31.
- Atz AM, Hawkins JA, Lu M, Cohen MS, Colan SD, Jagers J, et al. Surgical management of complete atrioventricular septal defect: associations with surgical technique, age, and trisomy 21. *J Thorac Cardiovasc Surg*. 2011;141:1371-9.
- Backer CL, Stewart RD, Bailliard F, Kelle AM, Webb CL, Mavroudis C. Complete atrioventricular canal: comparison of modified single-patch technique with two-patch technique. *Ann Thorac Surg*. 2007;84:2038-46.
- Harmandar B, Aydemir NA, Karaci AR, Sasmazel A, Saritas T, Bilal MS, et al. Results for surgical correction of complete atrioventricular septal defect: associations with age, surgical era, and technique. *J Card Surg*. 2012;27:745-53.
- Li D, Fan Q, Iwase T, Hirata Y, An Q. Modified single-patch technique versus two-patch technique for the repair of complete atrioventricular septal defect: a meta-analysis. *Pediatr Cardiol*. 2017;38:1456-64.
- Nunn GR. Atrioventricular canal: modified single patch technique. *Semin Thorac Cardiovasc Surg Pediatr Card Surg Annu*. 2007;28:31.
- McMurry TL, Hu Y, Blackstone EH, Kozower BD. Propensity scores: methods, considerations, and applications in the Journal of Thoracic and Cardiovascular Surgery. *J Thorac Cardiovasc Surg*. 2015;150:14-9.
- Winger DG, Nason KS. Propensity-score analysis in thoracic surgery: when, why, and an introduction to how. *J Thorac Cardiovasc Surg*. 2016;151:1484-7.
- Austin PC. An introduction to propensity score methods for reducing the effects of confounding in observational studies. *Multivar Behav Res*. 2011;46:399-424.
- Austin PC. Optimal caliper widths for propensity-score matching when estimating differences in means and differences in proportions in observational studies. *Pharm Stat*. 2011;150:61.
- Ginde S, Lam J, Hill GD, Cohen S, Woods RK, Mitchell ME, et al. Long-term outcomes after surgical repair of complete atrioventricular septal defect. *J Thorac Cardiovasc Surg*. 2015;150:369-74 [Erratum in *J Thorac Cardiovasc Surg*. 2016;152:292].
- Jonas RA, Mora B. Individualized approach to repair of complete atrioventricular canal: selective use of the traditional single-patch technique versus the Australian technique. *World J Pediatr Congenit Heart Surg*. 2010;1:78-86.
- Xie O, Brizard CP, d'Udekem Y, Galati JC, Kelly A, Yong MS, et al. Outcomes of repair of complete atrioventricular septal defect in the current era. *Eur J Cardiothorac Surg*. 2014;45:610-7.
- Bakhtiary F, Takacs J, Cho MY, Razek V, Dahnert I, Doenst T. Long-term results after repair of complete atrioventricular septal defect with two-patch technique. *Ann Thorac Surg*. 2010;89:1239-43.
- Birim O, van Gameren M, de Jong PL, Witsenburg M, van Osch-Gevers L, Bogers AJ. Outcome after reoperation for atrioventricular septal defect repair. *Interact Cardiovasc Thorac Surg*. 2009;9:83-7.
- Malhotra SP, Lacour-Gayet F, Mitchell MB, Clarke DR, Dines ML, Campbell DN. Reoperation for left atrioventricular valve regurgitation after atrioventricular septal defect repair. *Ann Thorac Surg*. 2008;86:147-51.
- Kanani M, Elliott M, Cook A, Juraszek A, Devine W, Anderson RH. Late incompetence of the left atrioventricular valve after repair of atrioventricular septal defects: the morphologic perspective. *J Thorac Cardiovasc Surg*. 2006;132:640-6. 646.e641-643.
- Crawford FA Jr. Repair of complete atrioventricular septal defects "single patch" technique. *Oper Tech Thorac Cardiovasc Surg*. 2004;9(3 Special Issue):221-32.
- Al-Hay AA, MacNeill SJ, Yacoub M, Shore DF, Shinebourne EA. Complete atrioventricular septal defect, Down syndrome, and surgical outcome: risk factors. *Ann Thorac Surg*. 2003;75:412-21.
- Wetter J, Sinzobahamvya N, Blaschczok C, Brecher AM, Gravinghoff LM, Schmaltz AA, et al. Closure of the zone of apposition at correction of complete atrioventricular septal defect improves outcome. *Eur J Cardiothorac Surg*. 2000;17:146-53.
- Litwin SB, Tweddell JS, Mitchell ME, Mussatto KA. The double patch repair for complete atrioventricular communis. *Semin Thorac Cardiovasc Surg Pediatr Card Surg Annu*. 2007;21-7.
- Backer CL, Eltayeb O, Monge MC, Wurlitzer KC, Hack MA, Boles LH, et al. Modified single patch: are we still worried about subaortic stenosis? *Ann Thorac Surg*. 2015;99:1671-5.
- Najm HK, Coles JG, Endo M, Stephens D, Rebecka IM, Williams WG, et al. Complete atrioventricular septal defects: results of repair, risk factors, and freedom from reoperation. *Circulation*. 1997;96(9 Suppl). II-311-5.
- Hoohenkerk GJ, Bruggemans EF, Rijlaarsdam M, Schoof PH, Koolbergen DR, Hazekamp MG. More than 30 years' experience with surgical correction of atrioventricular septal defects. *Ann Thorac Surg*. 2010;90:1554-61.

Key Words: complete atrioventricular septal defect, complete AV canal, atrioventricular septal defect, AVSD

Discussion

Dr David P. Bichell (Nashville, Tenn).

Dr Fong, thank you for the compelling and potentially practice-changing technique that we have all seen, but many of us, myself included, have been hesitant to fully adopt. Even leading up to this presentation, taking a casual poll among my colleagues, there are many who will not fully embrace the single-patch technique as you described as the Graham Nunn technique out of fear of producing a substrate for left ventricular outflow tract obstruction. You beautifully addressed that by propensity matching and have somewhat answered the question of whether patient selection is the reason.

I have a few questions about whether or not there could be some selection bias. You beautifully propensity matched for size of ventricular septal defect (VSD), which frankly is the common concern in going ahead with this technique on every patient. From a personal perspective, there are many large VSDs that are quite shallow, but when I look up toward the aortic valve and see a deep VSD that is scooped out, that's where I kind of back up and put a patch in, because it looks like it's going to crowd the subaortic area.

So your propensity matching has looked at size of VSD but not depth of VSD, and added to that is the fact that at your institution there are still quite a few who are 1 method versus the other. What are some of the choices that go into that decision? Is it depth of VSD?

Dr Laura S. Fong (*Sydney, Australia*).

Ideally we would have propensity matched the depth of VSD, but unfortunately not all the preoperative echocardiography data for each patient were available, so we were unable to measure that for each patient. It is possible that there is some selection bias; however,

with propensity matching the aim was to reduce that.

In our data set we did have centers that preferentially used the modified single patch in all cases, and we didn't show a difference between the modified single patch and double patch outcome.

Dr Bichell. It's interesting to see that there is a difference in the treatment of the zone of apposition from 1 technique to another, and you have addressed that a little bit as an institutional preference. But a very interesting upshot of that is that when the zone of apposition is not closed and the cleft is not closed there does not appear to be a difference in reintervention rates in excellent 20-year follow-up, so it leaves me wondering if we have to close the zone of apposition. What are your thoughts about that?

Dr Fong. We would have to approach that cautiously, because in our results, although there is no difference between the double patch and modified single patch, with the modified-single-patch group, almost all had complete zone of apposition closure compared with only 56% in the double-patch group. So I don't think based on our data we can say that you can just leave it open, but obviously in the hands of some surgeons who don't close the zone of apposition in all cases, they do still have good long-term results.

Dr Bichell. So, the real question here is would you recommend the single patch technique for all-comers or is there a reason to be selective?

I am interested in a show of hands of those who are here today: Who applies this for all patients? But first your comment about that.

Dr Fong. Our data suggest that a surgeon can choose either technique with the expectation the patient will have equal long-term results. Having not performed any myself, I can't really vouch for either.

Dr Bichell. So how many people in this room consistent for all-comers will use the Graham Nunn single patch technique regardless of VSD?

(Show of hands).

Dr Bichell. And how many not?

(Show of hands).

Dr Bichell. Okay. Well, I think this beautifully presented data and the propensity matching and the length of follow-up will probably result in a different ratio of the show of hands if we ask this question next year.