References

- Blasberg JD, Seder CW, Levenson G, Shan Y, Maloney JD, Macke RA. Video-assisted thoracoscopic lobectomy for lung cancer: current practice patterns and predictors of adoption. *Ann Thorac Surg*. 2016;102:1854-62.
- Giambrone GP, Smith MC, Wu X, Gaber-Baylis LK, Bhat AU, Zabih R, et al. Variability in length of stay after uncomplicated pulmonary lobectomy: is length of stay a quality metric or a patient metric? *Eur J Cardiothorac Surg*. 2016;49:e65-71.
- Rosen JE, Salazar MC, Dharmarajan K, Kim AW, Detterbeck FC, Boffa DJ. Length of stay from the hospital perspective: practice of early discharge is not associated with increased readmission risk after lung cancer surgery. *Ann Surg*. 2017;266:383-8.
- Kumar A, Asaf BB. Robotic thoracic surgery: the state of the art. *J Minim Access Surg*. 2015;11:60-7.
- Paul S, Altorki NK, Sheng S, Lee PC, Harpole DH, Onaitis MW, et al. Thoracoscopic lobectomy is associated with lower morbidity than open lobectomy: a propensity-matched analysis from the STS database. *J Thorac Cardiovasc Surg*. 2010;139:366-78.
- Fiore JF Jr, Bejjani J, Conrad K, Niculiseanu P, Landry T, Lee L, et al. Systematic review of the influence of enhanced recovery pathways in elective lung resection. *J Thorac Cardiovasc Surg*. 2016;151:708-15.e6.
- Madani A, Fiore JF Jr, Wang Y, Bejjani J, Sivakumaran L, Mata J, et al. An enhanced recovery pathway reduces duration of stay and complications after open pulmonary lobectomy. *Surgery*. 2015;158:899-908; discussion -10.
- Cerfolio RJ, Pickens A, Bass C, Katholi C. Fast-tracking pulmonary resections. *J Thorac Cardiovasc Surg*. 2001;122:318-24.
- Tovar EA. One-day admission for major lung resections in septuagenarians and octogenarians: a comparative study with a younger cohort. *Eur J Cardiothorac Surg*. 2001;20:449-54.
- Tovar EA, Roethe RA, Weissig MD, Lloyd RE, Patel GR. One-day admission for lung lobectomy: an incidental result of a clinical pathway. *Ann Thorac Surg*. 1998;65:803-6.
- Towe CW, Khil A, Ho VP, Perry Y, Argote-Greene L, Wu KM, et al. Early discharge after lung resection is safe: 10-year experience. *J Thorac Dis*. 2018;10:5870-8.
- Tong BC, Kim S, Kosinski A, Onaitis MW, Boffa DJ, Habib RH, et al. Penetration, completeness, and representativeness of the Society of Thoracic Surgeons General Thoracic Surgery Database for Lobectomy. *Ann Thorac Surg*. 2019;107:897-902.
- Kozower BD, Sheng S, O'Brien SM, Liptay MJ, Lau CL, Jones DR, et al. STS database risk models: predictors of mortality and major morbidity for lung cancer resection. *Ann Thorac Surg*. 2010;90:875-83.
- Seder CW, Raymond DP, Wright CD, Gaissert HA, Chang AC, Clinton S, et al. The Society of Thoracic Surgeons General Thoracic Surgery Database 2017 Update on Outcomes and Quality. *Ann Thorac Surg*. 2017;103:1378-83.
- Durheim MT, Collard HR, Roberts RS, Brown KK, Flaherty KR, King TE Jr, et al. Association of hospital admission and forced vital capacity endpoints with survival in patients with idiopathic pulmonary fibrosis: analysis of a pooled cohort from three clinical trials. *Lancet Respir Med*. 2015;3:388-96.
- Khan NA, Quan H, Bugar JM, Lemaire JB, Brant R, Ghali WA. Association of postoperative complications with hospital costs and length of stay in a tertiary care center. *J Gen Intern Med*. 2006;21:177-80.
- Van Haren RM, Mehran RJ, Mena GE, Correa AM, Antonoff MB, Baker CM, et al. Enhanced recovery decreases pulmonary and cardiac complications after thoracotomy for lung cancer. *Ann Thorac Surg*. 2018;106:272-9.
- Brunelli A, Thomas C, Dinesh P, Lumb A. Enhanced recovery pathway versus standard care in patients undergoing video-assisted thoracoscopic lobectomy. *J Thorac Cardiovasc Surg*. 2017;154:2084-90.
- Rogers LJ, Bleetman D, Messenger DE, Joshi NA, Wood L, Rasburn NJ, et al. The impact of enhanced recovery after surgery (ERAS) protocol compliance on morbidity from resection for primary lung cancer. *J Thorac Cardiovasc Surg*. 2018;155:1843-52.
- Harboe KM, Bardram L. The quality of cholecystectomy in Denmark: outcome and risk factors for 20,307 patients from the national database. *Surg Endosc*. 2011;25:1630-41.
- Roslyn JJ, Binns GS, Hughes EF, Saunders-Kirkwood K, Zinner MJ, Cates JA. Open cholecystectomy. A contemporary analysis of 42,474 patients. *Ann Surg*. 1993;218:129-37.
- Schauer PR, Ikramuddin S, Gourash W, Ramanathan R, Luketich J. Outcomes after laparoscopic Roux-en-Y gastric bypass for morbid obesity. *Ann Surg*. 2000;232:515-29.
- Waydia S, Gunawardene A, Gilbert J, Cota A, Finlay IG. 23-hour/next day discharge post-laparoscopic Roux-en-Y gastric bypass (LRYGB) surgery is safe. *Obes Surg*. 2014;24:2007-10.
- Gopaldas RR, Bakaeen FG, Dao TK, Walsh GL, Swisher SG, Chu D. Video-assisted thoracoscopic versus open thoracotomy lobectomy in a cohort of 13,619 patients. *Ann Thorac Surg*. 2010;89:1563-70.
- Kirby TJ, Mack MJ, Landreneau RJ, Rice TW. Lobectomy—video-assisted thoracic surgery versus muscle-sparing thoracotomy. A randomized trial. *J Thorac Cardiovasc Surg*. 1995;109:997-1002.
- Flores RM, Park BJ, Dycoco J, Aronova A, Hirth Y, Rizk NP, et al. Lobectomy by video-assisted thoracic surgery (VATS) versus thoracotomy for lung cancer. *J Thorac Cardiovasc Surg*. 2009;138:11-8.
- Allen MS, Wood DE, Hawkinson RW, Harpole DH, McKenna RJ, Walsh GL, et al. Prospective randomized study evaluating a biodegradable polymeric sealant for sealing intraoperative air leaks that occur during pulmonary resection. *Ann Thorac Surg*. 2004;77:1792-801.
- Malapert G, Hanna HA, Pages PB, Bernard A. Surgical sealant for the prevention of prolonged air leak after lung resection: meta-analysis. *Ann Thorac Surg*. 2010;90:1779-85.
- Refai M, Brunelli A, Salati M, Pompili C, Xiume F, Sabbatini A. Efficacy of anterior fissureless technique for right upper lobectomies: a case-matched analysis. *Eur J Cardiothorac Surg*. 2011;39:1043-6.
- Brown LM. "Moving right along" after lung resection, but the data suggest "not so fast" *J Thorac Cardiovasc Surg*. 2016;151:715-6.

Key Words: length of stay, patient discharge, postoperative complications, risk factors, thoracic surgery

Discussion

Dr Benjamin D. Kozower (St. Louis, Mo). Congratulations to you and your team for performing a thoughtful and hypothesis-generating study. The thoracic surgery community continues to face increased pressure to perform these complicated anatomic lung resections in a minimally invasive fashion

with extremely low complication rates and to further reduce our resource use. Sending a patient home on postoperative day 1 after a lobectomy or segmentectomy would have been unheard of when the STS thoracic database was started almost 20 years ago. Your results demonstrate a median length of stay of 4 days, about a 3.6% rate of discharge on postoperative day 1 for the entire series, and the hospital variation for early discharge on postoperative day 1 is impressive, and I think not so unexpected. I have 3 questions for you.

First, did you examine the discharge location for these patients and whether or not they were discharged with a chest tube in place? Both of these variables are included in the STS database, and I think it is important.

Dr Christopher W. Towe (Cleveland, Ohio). Yes, we did look at that, and 99% of patients in the day 1 group were discharged to home. The STS doesn't have a category for the hotel across the street. I think there is some gut reaction that perhaps this is happening, but taking the data as they are, the majority go home. Twelve were discharged to a

rehab facility, which is surprising to me. But generally they are going home.

In terms of chest tube rate, there were 6% of patients discharged with a chest tube in the day 1 group. This was significantly less than the normal discharge group, which was 7.5%, and I think is a perplexing problem. I would certainly rather keep a patient in the hospital for a day and get the tube out, for a variety of reasons, really, but the data are what they are.

Dr Kozower. My second question is, your data also demonstrate that a patient was more likely to be discharged on day 1 after a segmentectomy than a lobectomy. That surprised me and has not been my own personal experience. As I thought about it, I was concerned. Are some of these segmentectomies in our database glorified wedge resections? So another thing that could be looked at and potentially something you could do in the future, when you looked at these segmentectomies, were they really adequate lung cancer operations that actually sample N1 and N2 nodes?

Dr Towe. It is an excellent question, and I have thought about it, too, in that in an unadjusted way you would hope that segmentectomies might balance in poor lung function, but in the adjusted analysis, there was still a relationship. So I am surprised that the prevalence of some segmentectomies is occurring like that. My hypothesis is that they are probably small tumors in the tip of the superior segment or things like that and they are highly selected cases.

When I take a step back from the data, the thing that strikes me the most is operative time per hour, and my guess is that it's the easiest segments, not the ones where you are sort of chiseling through the hilum to find a perforating branch of an artery that are the ones that are the best candidates to go home.

Dr Kozower. How will you and your team use these results going forward and do you have any plans to study this, particularly from the patient perspective?

Dr Towe. There are a lot of questions that this study generates, at least for us. How can we increase use of day 1 discharge among the low performers? In other words, what can the low performers learn from the high performers? A rising tide lifts all of us, right? So, we would like to initiate a national study of which aspects of an ERAS protocol at the high performers could be instituted elsewhere.

We would also like to look at which aspects of patient selection could be used to tailor patients into different ERAS protocols. Ultimately, a 1-size-fits-all ERAS protocol doesn't make any sense. The patients who were at higher risk in the 2008 Wright study for 14-day length of stays should get a different protocol than the patients who are going home tomorrow. They shouldn't all be treated the same.

Your question about patient-centered outcomes is also interesting. We would love to know more about what patients want out of their hospitalizations. Currently we assume it's to go home, but I think that has a lot to do with building the expectations or engendering a sense that going home is the safest thing for them. As Phil Linden alluded to earlier, early preoperative education so that expectations are met.

Unidentified Speaker. I think you actually were alluding to a question that I had regarding the early presentation, too, from Dr Park. So obviously preoperative education is important in setting expectations, but what do you do postoperatively? How do you educate the patients and the families? What protocols do you have in place to prevent patients from being readmitted?

Dr Towe. At our institution, I think the realization that day 1 discharge is safe has shed new light on the opportunity for certain patients to do that. I think there are protocols floating around in each of our minds about chest tubes being left in for 2 days or on suction for day 1 and water seal for day 2. So, what we are trying to do is shift that whole expectation.

With that also comes things like phone calls to patients after they are discharged, staying in communication and not letting them feel like being discharged is like breaking up with someone, but the relationship continues after they go home so that they are not left in the lurch. Those communications can head off a readmission or a problem like a late complication more so than keeping someone in the hospital and watching them very closely.

Unidentified Speaker. What is your postoperative follow-up time?

Dr Towe. We routinely see patients within 3 weeks of discharge. It depends on, sort of, the clinic day to operating room day gap. But we tend to see patients at 2 weeks, for pneumonectomy patients, again, at 6 weeks, and then after that sort of on an as-needed basis.

Dr Susan D. Moffatt-Bruce (*Columbus, Ohio*). Would you alter your protocol should you adopt a postoperative day number 1 discharge from 3 weeks to maybe 7 days?

Dr Towe. These data would suggest that the patients who had day 1 discharge probably don't need follow-up. They are probably going to do fine whether you see them or not. So the concept is that I think we still believe that we need to keep these patients very close to the chest, but I think the irony, of course, is that they are the patients who are least likely to have problems. We have preselected them in a way that not only are they the healthiest people but also we have done the simplest operation we can do on them.

Dr Dirk Van Raemdonck (*Leuven, Belgium*). In the patients who were discharged on day 1, do you have any idea from the database whether they underwent operation in the morning or in the afternoon, because there is a difference of 12 hours.

Dr Towe. We were forced to calculate the number of midnights between admission and discharge. I know some hospitals would calculate this based on a 24-hour cycle. So there is some variability in what that means based on time of surgery versus time of discharge, and that's a potential limitation but within the realm of what I think makes sense.

Dr Moffatt-Bruce. The Chair's prerogative: Dr Kneuert, do you have any comments? You have

been my partner who has been sending some of his patients home on postoperative day number 1. Any additional comments as to what you do differently to keep these patients safe and still have a low readmission rate?

Dr Peter Jens Achim Kneuert (*Columbus, Ohio*). I would echo what the other attendees have said. Setting expectations is probably the most important thing going forward. It does have to do with the approach. I think the smaller the incisions and the less long the operation is, the better the chance

we have for a fast recovery. We should all strive to continue to decrease the patient's length of stay.

Readers who found these articles interesting may also like to read the following papers found in recent and future issues of our sister publications, *Seminars in Thoracic and Cardiovascular Surgery* and *Operative Techniques in Thoracic and Cardiovascular Surgery*!

Thoracic: Lung Cancer

STCVS: Subxiphoid Uniportal VATS for Thymic and Combined Mediastinal and Pulmonary Resections – A Two-Year Experience. Abu-Akar F, Gonzalez-Rivas D, Yang C, Lin L, Wu L, Jiang L. *Semin Thorac Cardiovasc Surg.* 2019;31(3):614-619.

Commentary: The Ongoing Debate on the Ideal Approach to Thymectomy. Figueroa PU. *Semin Thorac Cardiovasc Surg.* 2019;31(3):620-621.