Commentary: Shaggy aorta in thoracoabdominal aortic aneurysm repair, an insidiously growing threat

Tohru Asai, MD, PhD

From the Department of Cardiovascular Surgery, Juntendo University, Tokyo, Japan.

Disclosures: Author has nothing to disclose with regard to commercial support.

Received for publication Aug 13, 2019; revisions received Aug 13, 2019; accepted for publication Aug 14, 2019; available ahead of print Sept 19, 2019.

Address for reprints: Tohru Asai, MD, PhD, Department of Cardiovascular Surgery, Juntendo University, Faculty of Medicine, 2-1-1, Hongo, Bunkyo-ku, Tokyo 113-8421, Japan (E-mail: t.asai.jj@juntendo.ac.jp).

J Thorac Cardiovasc Surg 2020;160:898

0022-5223/\$36.00

Copyright © 2019 Published by Elsevier Inc. on behalf of The American Association for Thoracic Surgery

<https://doi.org/10.1016/j.jtcvs.2019.08.072>

“Shaggy” aorta is a diffuse atherosclerotic disease, first reported by Hollier and colleagues¹ in 1991. The associated spontaneous visceral embolization is common, and its mortality is high with nonoperative and operative therapies. In aging populations, this potentially fatal disease is insidiously growing.

In this issue of *The Journal of Thoracic and Cardiovascular Surgery*, Yokawa and colleagues² in Kobe University report their experience of open repair for thoracoabdominal aortic aneurysm (TAAA) using segmental staged aortic clamping, and evaluate the effect of shaggy aorta. They define shaggy aorta as atheroma thickness of at least 5 mm with irregular atheroma surface, seen using enhanced computed tomography. They uniquely classified 251 patients into 3 groups: dissection aneurysm, degenerative aneurysm without shaggy aorta, and degenerative aneurysm with shaggy aorta. In their TAAA repair group, the incidence of shaggy aorta was 14.3%. Even in their experienced hands, the shaggy aorta group had significantly worse mortality (33.3%) and significantly worse incidence of spinal cord injury (27.8%). The composite outcomes including acute renal failure were also worse.

Open TAAA repair has been the most extensive and challenging operation for cardiovascular surgeons for years. The Kobe University group mostly uses segmental staged aortic clamping, even with careful intraoperative epiaortic scanning. However, as they commented, deep hypothermic circulatory arrest^{3,4} should be considered in higher-risk shaggy aorta patients, especially when the proximal descending aorta is involved. To minimize risks of organ embolization and spinal cord injury during the repair is especially crucial with shaggy aorta. Endovascular TAAA repair for shaggy aorta also carries the risks of diffuse organ

Tohru Asai, MD, PhD

Central Message

Shaggy aorta carries high risks for organ embolization and spinal cord injury in thoracoabdominal aortic aneurysm repair.

See Article page 889.

embolization. The definition of shaggy aorta, in terms of anatomy and pathology, including ways to distinguish highest-risk regions and how best to approach them, and the possibility of other treatments need to be further examined in future. However, this report should be noted for our readers to take extreme care to evaluate such aortic disease preoperatively, as not just dissection or degenerative, but also with or without shaggy aorta, for safe choice of operative techniques.

References

1. Hollier LH, Kazmier FJ, Ochsner J, Bowen JC, Procter CD. “Shaggy” aorta syndrome with atheromatous embolization to visceral vessels. *Ann Vasc Surg.* 1991;5: 439-44.
2. Yokawa K, Ikeno Y, Henmi S, Yamanaka K, Okada K, Okita Y. Impact of shaggy aorta on outcomes of open thoracoabdominal aortic aneurysm repair. *J Thorac Cardiovasc Surg.* 2020;160:889-97.e1.
3. Kouchoukos NT, Kulik A, Castner CF. Outcomes after thoracoabdominal aortic aneurysm repair using hypothermic circulatory arrest. *J Thorac Cardiovasc Surg.* 2013;145:S139-41.
4. Tanaka H, Minatoya K, Sasaki H, Seike Y, Itonaga T, Oda T, et al. Recent thoracoabdominal aortic repair outcomes using moderate-to-deep hypothermia combined with targeted reconstruction of the Adamkiewicz artery. *Interact Cardiovasc Thorac Surg.* 2015;20:605-10, discussion: 610.