Dr Cerfolio discloses past relationships with C-SATS, Intuitive Surgical, Myriad G, and TEGO Corporation, and he is the president of ROLO-7 Consulting Firm, and the author of 2 books on leadership.

**REPLY FROM THE
AUTHORS:
ACCOUNTABILITY AND
LEADERSHIP
CHANGES CULTURE—
YES, EVEN FOR LYMPH
NODES**

Reply to the Editor:

The authors have raised several important and interesting points that are undoubtedly true. In general, we agree with their comments; our view on this issue is quite different and cuts to the concept of leadership. And to the uninitiated, this editorial might seem harsh, but our experience suggests that this is the best way to drive excellence and change culture in a health care system. The quality of patient care is the single most valued and important element of health care. Lymph node resection during lung cancer surgery is a quality metric of the surgeon not of the pathologist. Our view is summarized by the phrase, “Leaders take accountability for all variables and for all metrics or surrogates of quality that are under our control—great leaders inspire those around them to make the entire process under their control. We don’t point fingers or cast blame on others nor do we blur lines of attribution. Rather, we willingly accept all the variables that influence our outcomes even those that initially might not seem to be in our purview.” There can be no higher goal than providing our patients the best care possible, and this means optimizing any metrics that might reflect or be surrogates of quality patient care.

This takes us to thoracic lymph nodes. If the number of lymph nodes removed and/or the number of lymph nodes stations assessed is a surrogate of quality (and it is) then why would we as leaders NOT take full accountability for it? The removal of mediastinal and hilar thoracic lymph nodes and how they are labeled during cancer surgery is a perfect example of our philosophy.

Yes, it is true that many surgeons “remove the paratracheal lymph nodes en bloc”¹ but the labeling of them as such is not the pathologist’s fault. It is the surgeon’s fault. The specimens should be labeled as, for example, “3 lymph nodes removed from station 4R.” Likewise, those removed from the 2R station should be separately labeled, as for example, “5 lymph nodes removed from station 2R.”

Yes, it is true that surgeons often take the 5 and 6 lymph nodes out en block and count them as a packet. Again, the

surgeon and his or her team should label it as, “4 lymph nodes removed from station 5,” etc.

Yes, it is true that many pathologists don’t look at N1 lymph nodes in the resected specimen. These are our metrics of the quality of our surgery. We as leaders should accept the responsibility. When a lobectomy or segmentectomy specimen is resected the surgeon has every opportunity to remove all of the lymph nodes that he or she might have included in that specimen on a back table and label them properly as described previously.

We have been using this technique since 1996. It is simple. It works. It completely prevents all of the problems articulated in the editorial. And you only need 1 pathologic slip for all of the lymph nodes from each station and then enumerate how many total lymph nodes that were removed from each station. If the surgeon truly believes there is value in en bloc resection for which we are unaware of any data to support, then they can simply dissect the packet out on the back table and send it to pathology properly labeled. The pathologist cannot determine where a 2R becomes a 4R. And they are not to blame for the poorly labeled surgical specimen. Leaders do not blame their teammates, instead they inspire others to perform better and take accountability.

The unfortunate and pervasive history of poorly performed and inadequately labeled lymph node dissection for patients with lung cancer has diminished the true value of lung cancer resection surgery compared with ablative therapies such as stereotactic radiosurgery, radiofrequency ablation, and now robotic bronchoscopic ablation of peripheral nodules.

Highly functional teams that provide their patients the best of care, which every patient deserves every day, is marked by great leaders. And even lymph nodes can teach us about leadership accountability and excellence.

*Robert J. Cerfolio, MD, MBA
Department of Thoracic Surgery
New York University
New York, NY*

Reference

1. Kneuert PJ, Merritt RE. Lymph node stations counts in retrospective studies—important end point, but interpret with caution. *J Thorac Cardiovasc Surg.* 2020;159:e145.

<https://doi.org/10.1016/j.jtcvs.2019.09.005>

**REPLY FROM THE
AUTHORS: THE
QUANTIFICATION OF
NODAL DISEASE HAS
PROGNOSTIC
RELEVANCE**

Reply to the Editor:

The letter by Drs Kneuert and Merritt describing the limitations encountered in retrospective studies to count

Authors have nothing to disclose with regard to commercial support.

the number of nodal stations explored in lung cancer surgery helps us understand why it is difficult to achieve the international standards of complete resection, and we cannot agree more with them.¹ Sooner or later, the categories of the N component of the tumor, node, and metastases (TNM) classification of lung cancer will be on the basis of the quantification of nodal disease in addition to or instead of the anatomic location of the involved lymph nodes. If this innovation is included in future editions of the TNM classification of lung cancer, it will even be more important to precisely quantify nodal disease. The number of involved nodal zones, in the analyses performed for the 7th edition,² and the number of involved nodal stations, in the analyses for the 8th edition,³ showed that the number of involved zones and stations had prognostic implications. The number of involved lymph nodes and the lymph node ratio (ie, the number of involved lymph nodes divided by the total number of lymph nodes removed) also have prognostic relevance.⁴ If we are to quantify nodal disease in the future to determine the categories of the N component of the TNM classification, we certainly will have to be more accurate in performing systematic nodal dissection, in counting the number of involved lymph nodes, nodal zones, or nodal station, and in labeling the removed lymph nodes. Molnar⁵ designed a lymph node tray to facilitate labeling and handling of removed lymph nodes in the operating room and in the pathology department. Osarogiagbon and colleagues⁶ proposed the use of a specialized specimen collection kit for mediastinal lymph nodes that improved lymph node staging. If these devices are not used, the surgeon should identify individual lymph nodes and label them according to the lymph node map proposed by the International Association for the Study of Lung Cancer to facilitate their counting.⁷

The study by Drs Kneuert and Merritt and colleagues,⁸ and the reply from Drs Kneuert and Merritt to our comment¹ show that there is much room for improvement in intraoperative lymph node assessment. It is also clear that the intrinsic limitations of retrospective studies should be overcome in prospective studies, especially because the quantification of nodal disease, no doubt, will be used to define the anatomic extent of lung cancer.

Diego González-Rivas, MD, FECTS^{a,b}
Ramón Rami-Porta, MD, FECTS^{c,d}

^aDepartment of Thoracic Surgery
Shanghai Pulmonary Hospital
Tongji University School of Medicine
Shanghai, China

^bDepartment of Thoracic Surgery and Minimally Invasive

Thoracic Surgery Unit
Coruña Hospital
Coruña, Spain

^cDepartment of Thoracic Surgery
Hospital Universitari Mútua Terrassa
University of Barcelona
Terrassa, Spain

and ^dNetwork of Centers for Biomedical Research in
Respiratory Diseases Lung Cancer Group
Terrassa, Spain

References

1. Kneuert PJ, Merritt RE. Lymph node stations counts in retrospective studies—important end point, but interpret with caution. *J Thorac Cardiovasc Surg*. 2020;159:e145.
2. Rusch VW, Crowley J, Giroux DJ, Goldstraw P, Im JG, Tsuboi M, et al. The IASLC Lung Cancer Staging Project: proposals for the revision of the N descriptors in the forthcoming seventh edition of the TNM classifications for lung cancer. *J Thorac Oncol*. 2007;2:603-12.
3. Asamura H, Chansky K, Crowley J, Goldstraw P, Rusch VW, Vansteenkiste JF, et al. The International Association for the Study of Lung Cancer Lung Cancer Staging Project: proposals for the revisions of the N descriptors in the forthcoming 8th edition of the TNM classification for lung cancer. *J Thorac Oncol*. 2015;10:1675-84.
4. Rami-Porta R. Quantification of regional lymph node involvement in lung cancer. *Thorax*. 2011;66:271-2.
5. Molnar TF. A new device for the identification of lymph nodes at lung cancer surgery. *Eur J Cardiothorac Surg*. 2007;31:311-2.
6. Osarogiagbon RU, Miller LE, Ramirez RA, Wang CG, O'Brien TF, Yu X, et al. Use of a surgical specimen-collection kit to improve mediastinal lymph-node examination of resectable lung cancer. *J Thorac Oncol*. 2012;7:1276-82.
7. Rusch VW, Asamura H, Watanabe H, Giroux DJ, Rami-Porta R, Goldstraw P. The IASLC Lung Cancer Staging Project. A proposal for a new international lymph node map in the forthcoming seventh edition of the TNM classification for lung cancer. *J Thorac Oncol*. 2009;4:568-77.
8. Kneuert PJ, Cheufou DH, D'Souza DM, Mardanzai K, Abdel-Rasoul M, Theegarten D, et al. Propensity-score adjusted comparison of pathologic nodal upstaging by robotic, video-assisted thoracoscopic, and open lobectomy for non-small cell lung cancer. *J Thorac Cardiovasc Surg*. 2019;158:1457-66.

<https://doi.org/10.1016/j.jtcvs.2019.09.007>

PREOPERATIVE PULMONARY REHABILITATION To the Editor:

I enjoyed the recent article by Gravier and colleagues¹ in the June issue of *The Journal of Thoracic and Cardiovascular Surgery*, “Effect of Prehabilitation on Ventilatory Efficiency in Non-Small Cell Lung Cancer Patients: A Cohort Study.” Our group did extensive outpatient preoperative pulmonary rehabilitation in patients with lung cancer with forced expiratory volume in 1 second < 1.56 L/s who underwent subsequent lung resection. The study compared those receiving preoperative pulmonary rehabilitation with a second group without any preoperative pulmonary therapy.

The group receiving pulmonary rehabilitation improved their forced expiratory volume in 1 second significantly